

ELETRIPTAN HBR- eletriptan hbr tablet, film coated
Direct_Rx

Eletriptan HBR

Eletriptan hydrobromide tablets are indicated for the acute treatment of migraine with or without aura in adults.

Limitations of Use:

Use only if a clear diagnosis of migraine has been established. If a patient has no response to the first migraine attack treated with eletriptan hydrobromide tablets, reconsider the diagnosis of migraine before eletriptan hydrobromide tablets are administered to treat any subsequent attacks.

Eletriptan hydrobromide tablets are not intended for the prevention of migraine attacks.

Safety and effectiveness of eletriptan hydrobromide tablets have not been established for cluster headache.

The maximum recommended single dose is 40 mg.

In controlled clinical trials, single doses of 20 mg and 40 mg were effective for the acute treatment of migraine in adults. A greater proportion of patients had a response following a 40 mg dose than following a 20 mg dose [see CLINICAL STUDIES (14)].

If the migraine has not resolved by 2 hours after taking eletriptan hydrobromide tablets, or returns after transient improvement, a second dose may be administered at least 2 hours after the first dose. The maximum daily dose should not exceed 80 mg.

The safety of treating an average of more than 3 migraine attacks in a 30-day period has not been established.

20 mg Tablets: Orange colored, round, biconvex, film-coated tablets debossed with 'EL' on one side and '20' on the other side.

40 mg Tablets: Orange colored, round, biconvex, film-coated tablets debossed with 'EL' on one side and '40' on the other side.

Eletriptan hydrobromide tablets are contraindicated in patients with:

Ischemic coronary artery disease (CAD) (angina pectoris, history of myocardial infarction, or documented silent ischemia) or coronary artery vasospasm, including Prinzmetal's angina [see WARNINGS AND PRECAUTIONS (5.1)].

Wolff-Parkinson-White syndrome or arrhythmias associated with other cardiac accessory conduction pathway disorders [see WARNINGS AND PRECAUTIONS (5.2)].

History of stroke, transient ischemic attack (TIA), or history or current evidence of hemiplegic or basilar migraine because these patients are at a higher risk of stroke [see WARNINGS AND PRECAUTIONS (5.4)].

Peripheral vascular disease [see WARNINGS AND PRECAUTIONS (5.5)].

Ischemic bowel disease [see WARNINGS AND PRECAUTIONS (5.5)].

Uncontrolled hypertension [see WARNINGS AND PRECAUTIONS (5.8)].

Recent use (i.e., within 24 hours) of another 5-hydroxytryptamine₁ (5-HT₁) agonist, ergotamine-containing medication, or ergot-type medication such as dihydroergotamine (DHE) or methysergide [see DRUG INTERACTIONS (7.1)].

Hypersensitivity to eletriptan hydrobromide tablets (angioedema and anaphylaxis seen) [see WARNINGS AND PRECAUTIONS (5.9)].

Recent use (i.e., within at least 72 hours) of the following potent CYP3A4 inhibitors: ketoconazole, itraconazole, nefazodone, troleandomycin, clarithromycin, ritonavir, or nelfinavir [see DRUG INTERACTIONS (7.2) and CLINICAL PHARMACOLOGY (12.3)].

Eletriptan hydrobromide should only be used where a clear diagnosis of migraine has been established.

5.1 Myocardial Ischemia, Myocardial Infarction, and Prinzmetal's Angina

Eletriptan hydrobromide is contraindicated in patients with ischemic or vasospastic CAD. There have been rare reports of serious cardiac adverse reactions, including acute myocardial infarction, occurring within a few hours following administration of eletriptan hydrobromide. Some of these reactions occurred in patients without known CAD. Eletriptan hydrobromide may cause coronary artery vasospasm (Prinzmetal's angina), even in patients without a history of CAD.

Perform a cardiovascular evaluation in triptan-naïve patients who have multiple cardiovascular risk factors (e.g., increased age, diabetes, hypertension, smoking, obesity, strong family history of CAD) prior to receiving eletriptan hydrobromide. Do not use eletriptan hydrobromide if there is evidence of CAD or coronary artery vasospasm [see CONTRAINDICATIONS (4)]. For patients with multiple cardiovascular risk factors who have a

negative cardiovascular evaluation, consider administering the first eletriptan hydrobromide dose in a medically-supervised setting and performing an electrocardiogram (ECG) immediately following administration of eletriptan hydrobromide. For such patients, consider periodic cardiovascular evaluation in intermittent long-term users of eletriptan hydrobromide.

5.2 Arrhythmias

Life-threatening disturbances of cardiac rhythm including ventricular tachycardia and ventricular fibrillation leading to death have been reported within a few hours following the administration of 5-HT₁ agonists. Discontinue eletriptan hydrobromide if these disturbances occur. Eletriptan hydrobromide is contraindicated in patients with Wolff-Parkinson-White syndrome or arrhythmias associated with other cardiac accessory conduction pathway disorders [see CONTRAINDICATIONS (4)].

5.3 Chest, Throat, Neck and/or Jaw Pain/Tightness/Pressure

Sensations of tightness, pain, and pressure in the chest, throat, neck, and jaw commonly occur after treatment with eletriptan hydrobromide and are usually non-cardiac in origin. However, perform a cardiac evaluation if these patients are at high cardiac risk. Eletriptan hydrobromide is contraindicated in patients with CAD or Prinzmetal's variant angina [see CONTRAINDICATIONS (4)].

5.4 Cerebrovascular Events

Cerebral hemorrhage, subarachnoid hemorrhage, and stroke have occurred in patients treated with 5-HT₁ agonists, and some have resulted in fatalities. In a number of cases, it appears possible that the cerebrovascular events were primary, the 5-HT₁ agonist having been administered in the incorrect belief that the symptoms experienced were a consequence of migraine, when they were not.

Before treating headaches in patients not previously diagnosed as migraineurs, and in migraineurs who present with symptoms atypical of migraine, other potentially serious neurological conditions need to be excluded. Eletriptan hydrobromide is contraindicated in patients with a history of stroke or TIA [see CONTRAINDICATIONS (4)].

5.5 Other Vasospasm Reactions

Eletriptan hydrobromide may cause non-coronary vasospastic reactions, such as peripheral vascular ischemia, gastrointestinal vascular ischemia and infarction (presenting with abdominal pain and bloody diarrhea), and Raynaud's syndrome. In patients who experience symptoms or signs suggestive of a vasospastic reaction following the use of any 5-HT₁ agonist, rule out a vasospastic reaction before receiving additional eletriptan hydrobromide doses [see CONTRAINDICATIONS (4)].

5.6 Medication Overuse Headache

Overuse of acute migraine drugs (e.g. ergotamine, triptans, opioids, or combination of these drugs for 10 or more days per month) may lead to exacerbation of headache (medication overuse headache). Medication overuse headache may present as migraine-like daily headaches or as a marked increase in frequency of migraine attacks. Detoxification of patients, including withdrawal of the overused acute migraine drugs and treatment of withdrawal symptoms (which often includes a transient worsening of headache) may be necessary.

5.7 Serotonin Syndrome

Serotonin syndrome may occur with eletriptan hydrobromide, particularly during co-administration with selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), and monoamine oxidase (MAO) inhibitors [see DRUG INTERACTIONS (7.3)]. Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination), and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). The onset of symptoms usually occurs within minutes to hours of receiving a new or a greater dose of a serotonergic medication. Discontinue eletriptan hydrobromide if serotonin syndrome is suspected.

5.8 Increase in Blood Pressure

Significant elevation in blood pressure, including hypertensive crisis with acute impairment of organ systems, has been reported on rare occasions in patients treated with 5-HT₁ agonists, including patients without a history of hypertension. Monitor blood pressure in patients treated with eletriptan hydrobromide. Eletriptan hydrobromide is contraindicated in patients with uncontrolled hypertension [see CONTRAINDICATIONS (4)].

5.9 Anaphylactic/Anaphylactoid Reactions

There have been reports of anaphylaxis, anaphylactoid, and hypersensitivity reactions including angioedema in patients receiving eletriptan hydrobromide. Such reactions can be life threatening or fatal. In general, anaphylactic reactions to drugs are more likely to occur in individuals with a history of sensitivity to multiple

allergens. Eletriptan hydrobromide is contraindicated in patients with a history of hypersensitivity reaction to eletriptan hydrobromide [see CONTRAINDICATIONS (4)].

The following adverse reactions are described elsewhere in other sections of the prescribing information:

Myocardial ischemia and myocardial infarction, and Prinzmetal's angina [see WARNINGS AND PRECAUTIONS (5.2)]

Arrhythmias [see WARNINGS AND PRECAUTIONS (5.3)]

Chest, throat, neck, and/or jaw pain/tightness/pressure [see WARNINGS AND PRECAUTIONS (5.4)]

Cerebrovascular events [see WARNINGS AND PRECAUTIONS (5.4)]

Other vasospasm reactions [see WARNINGS AND PRECAUTIONS (5.5)]

Medication overuse headache [see WARNINGS AND PRECAUTIONS (5.6)]

Serotonin syndrome [see WARNINGS AND PRECAUTIONS (5.7)]

Increase in blood pressure [see WARNINGS AND PRECAUTIONS (5.8)]

Hypersensitivity reactions [see CONTRAINDICATIONS (4) and WARNINGS AND PRECAUTIONS (5.9)]

6.1 Clinical Trials Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

Among 4,597 patients who treated the first migraine headache with eletriptan hydrobromide in short-term placebo-controlled trials, the most common adverse reactions reported with treatment with eletriptan hydrobromide were asthenia, nausea, dizziness, and somnolence. These reactions appear to be dose-related.

In long-term open-label studies where patients were allowed to treat multiple migraine attacks for up to 1 year, 128 (8.3%) out of 1,544 patients discontinued treatment due to adverse reactions.

Table 1 lists adverse reactions that occurred in the subset of 5,125 migraineurs who received eletriptan doses of 20 mg, 40 mg and 80 mg or placebo in worldwide placebo-controlled clinical trials.

Only adverse reactions that were more frequent in a eletriptan hydrobromide treatment group compared to the placebo group with an incidence greater than or equal to 2% are included in Table 1.

Table 1: Adverse Reactions Incidence in Placebo-Controlled Migraine Clinical Trials: Reactions Reported by \geq 2% Patients Treated with Eletriptan Hydrobromide and More Than Placebo

Adverse Reaction Type

Placebo

(n=988)

Eletriptan Hydrobromide

20 mg

(n=431)

Eletriptan Hydrobromide 40 mg

(n=1774)

Eletriptan Hydrobromide

80 mg

(n=1932)

ATYPICAL SENSATIONS

Paresthesia

2%

3%

3%

4%

Flushing/feeling of warmth

2%

2%

2%

2%

PAIN AND PRESSURE SENSATIONS

Chest - tightness/pain/pressure

1%

1%

2%

4%

Abdominal -

pain/discomfort/stomach pain/ cramps/pressure

1%

1%

2%

2%

DIGESTIVE

Dry mouth

2%

2%

3%

4%

Dyspepsia

1%

1%

2%

2%

Dysphagia-throat tightness/difficulty swallowing

0.2%

1%

2%

2%

Nausea

5%

4%

5%

8%

NEUROLOGICAL

Dizziness

3%

3%

6%

7%

Somnolence

4%

3%

6%

7%

Headache

3%

4%

3%

4%

OTHER

Asthenia

3%

4%
5%
10%

The frequency of adverse reactions in clinical trials did not increase when up to 2 doses of eletriptan hydrobromide were taken within 24 hours. The incidence of adverse reactions in controlled clinical trials was not affected by gender, age, or race of the patients. Adverse reaction frequencies were also unchanged by concomitant use of drugs commonly taken for migraine prophylaxis (e.g., SSRIs, beta blockers, calcium channel blockers, tricyclic antidepressants), estrogen replacement therapy or oral contraceptives.

6.2 Postmarketing Experience

The following adverse reaction(s) have been identified during post approval use of eletriptan hydrobromide. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Neurological: seizure

Digestive: vomiting

7.1 Ergot-Containing Drugs Including Other 5-HT_{1B/1D} Agonists

Ergot-containing drugs have been reported to cause prolonged vasospastic reactions. Because these effects may be additive, use of ergotamine-containing or ergot-type medications (like dihydroergotamine [DHE] or methysergide) and eletriptan hydrobromide within 24 hours of each other is contraindicated. Concomitant use of other 5-HT₁ agonists within 24 hours of eletriptan hydrobromide treatment is contraindicated [see CONTRAINDICATIONS (4)].

7.2 CYP3A4 Inhibitors

Potent CYP3A4 inhibitors significantly increase the exposure of eletriptan hydrobromide. Eletriptan hydrobromide should not be used within at least 72 hours of treatment with potent CYP3A4 inhibitors [see CONTRAINDICATIONS (4) and CLINICAL PHARMACOLOGY (12.3)].

7.3 Selective Serotonin Reuptake Inhibitors/Serotonin and Norepinephrine Reuptake Inhibitors and Serotonin Syndrome

Cases of serotonin syndrome have been reported during co-administration of triptans and SSRIs, SNRIs, TCAs and MAO inhibitors [see WARNINGS AND PRECAUTIONS (5.7)].

8.1 Pregnancy

Risk Summary

Available human data on the use of eletriptan hydrobromide in pregnant women are not sufficient to draw conclusions about drug-associated risk for major birth defects and miscarriage. In animal studies, oral administration of eletriptan during pregnancy or throughout pregnancy and lactation was associated with developmental toxicity (decreased fetal and pup weights, increased incidences of fetal structural abnormalities, decreased pup viability) at clinically-relevant doses [see Data].

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. The reported rate of major birth defects among deliveries to women with migraine ranged from 2.2% to 2.9% and the reported rate of miscarriage was 17%, which were similar to rates reported in women without migraine.

Clinical Considerations

Disease-Associated Maternal and/or Embryo/Fetal Risk: Several studies have suggested that women with migraine may be at increased risk of preeclampsia and gestational hypertension during pregnancy.

Data

Human Data

A study using linked data from the Medical Birth Registry of Norway to the Norwegian Prescription Database compared pregnancy outcomes in women who redeemed prescriptions for triptans during pregnancy, as well as a migraine disease comparison group who redeemed prescriptions for triptans before pregnancy only. Of the 189 women who redeemed prescriptions for eletriptan during the first trimester, 4 (2.1%) had infants with major congenital malformations, while for the 174 women who redeemed prescriptions for eletriptan before, but not during, pregnancy, 11 (6.3%) had infants with major congenital malformations. Methodological limitations of this study, including small size of the eletriptan population and infrequent events, do not allow for thorough characterization of risk.

Animal Data

When pregnant rats were administered eletriptan (0, 10, 30, or 100 mg/kg/day) during the period of organogenesis, fetal weights were decreased and the incidences of vertebral and sternbral variations were increased at 100 mg/kg/day (approximately 12 times the maximum recommended human dose [MRHD] of 80 mg/day on a mg/m² basis). The 30 and 100 mg/kg/day doses were also maternally toxic, as evidenced by decreased maternal body weight gain during gestation. The no-effect dose for adverse effects on embryofetal development in rats was 30 mg/kg/day, which is approximately 4 times the MRHD on a mg/m² basis.

When eletriptan (0, 5, 10, or 50 mg/kg/day) was orally administered to pregnant rabbits throughout organogenesis, fetal weights were decreased at 50 mg/kg/day. The incidences of fused sternbrae and vena cava deviations were increased at all doses. Maternal toxicity was not evident at any dose. A no-effect dose for adverse effects on embryofetal development in rabbits was not established; the lowest dose tested (5 mg/kg/day) is similar to the MRHD on a mg/m² basis.

Oral administration of eletriptan (0, 5, 15, or 50 mg/kg/day) to female rats throughout pregnancy and lactation resulted in a decrease in offspring viability and body weight at the highest dose tested. The no-effect dose for adverse effects on pre- and postnatal development in rats (15 mg/kg/day) is approximately 2 times the MRHD on a mg/m² basis.

8.2 Lactation

Risk Summary

Eletriptan is excreted in human milk. There are no data on the effects of eletriptan on the breastfed infant or the effects of eletriptan on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for eletriptan hydrobromide and any potential adverse effects on the breastfed child from eletriptan hydrobromide or from the underlying maternal condition. Infant exposure can be minimized by avoiding breastfeeding for 24 hours after treatment.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

The efficacy of eletriptan hydrobromide tablets (40 mg) in patients 11 to 17 was not established in a randomized, placebo-controlled trial of 274 adolescent migraineurs [see CLINICAL STUDIES (14)]. Adverse reactions observed were similar in nature to those reported in clinical trials in adults. Postmarketing experience with other triptans includes a limited number of reports that describe pediatric patients who have experienced clinically serious adverse reactions that are similar in nature to those reported rarely in adults. Long-term safety of eletriptan was studied in 76 adolescent patients who received treatment for up to one year. A similar profile of adverse reactions to that of adults was observed. The long-term safety of eletriptan in pediatric patients has not been established.

8.5 Geriatric Use

Blood pressure was increased to a greater extent in elderly subjects than in young subjects. The pharmacokinetic disposition of eletriptan in the elderly is similar to that seen in younger adults [see CLINICAL PHARMACOLOGY (12.3)]. In clinical trials, there were no apparent differences in efficacy or the incidence of adverse reactions between patients under 65 years of age and those 65 and above.

8.6 Hepatic Impairment

The effect of severe hepatic impairment on eletriptan hydrobromide metabolism has not been evaluated. Eletriptan hydrobromide is not recommended for use in patients with severe hepatic impairment [see CLINICAL PHARMACOLOGY (12.3)].

The elimination half-life of eletriptan is about 4 hours [see CLINICAL PHARMACOLOGY (12.3)], therefore monitoring of patients after overdose with eletriptan should continue for at least 20 hours or longer while symptoms or signs persist.

There is no specific antidote to eletriptan.

It is unknown what effect hemodialysis or peritoneal dialysis has on the serum concentration of eletriptan.

Eletriptan hydrobromide tablets contain eletriptan hydrobromide, which is a selective 5-hydroxytryptamine 1B/1D (5-HT_{1B/1D}) receptor agonist. Eletriptan hydrobromide is chemically designated as (R)-3-[(1-Methyl-2-pyrrolidinyl)methyl]-5-[2-(phenylsulfonyl)ethyl]-1H-indole monohydrobromide, and it has the following chemical structure:
[Chemical Structure]

The molecular formula is C₂₂H₂₆N₂O₂S · HBr, representing a molecular weight of 463.43. Eletriptan hydrobromide is an off-white to brown color powder that is freely soluble in methanol and very slightly soluble in water.

Each eletriptan hydrobromide tablet for oral administration contains 24.23 or 48.46 mg of eletriptan hydrobromide equivalent to 20 mg or 40 mg of eletriptan, respectively. Each tablet also contains the inactive ingredients croscarmellose sodium, FD&C yellow 6 aluminum lake, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, titanium dioxide and triacetin.

12.1 Mechanism of Action

Eletriptan binds with high affinity to 5-HT_{1B}, 5-HT_{1D} and 5-HT_{1F} receptors, has modest affinity for 5-HT_{1A}, 5-HT_{1E}, 5-HT_{2B} and 5-HT₇ receptors.

Migraines are likely due to local cranial vasodilatation and/or to the release of sensory neuropeptides (vasoactive intestinal peptide, substance P and calcitonin gene-related peptide) through nerve endings in the trigeminal system. The therapeutic activity of eletriptan hydrobromide for the treatment of migraine headache is thought to be due to the agonist effects at the 5-HT_{1B/1D} receptors on intracranial blood vessels (including the arterio-venous anastomoses) and sensory nerves of the trigeminal system which result in cranial vessel constriction and inhibition of pro-inflammatory neuropeptide release.

12.3 Pharmacokinetics

Absorption: Eletriptan is well absorbed after oral administration with peak plasma levels occurring approximately 1.5 hours after dosing to healthy subjects. In patients with moderate to severe migraine the median T_{max} is 2.0 hours. The mean absolute bioavailability of eletriptan is approximately 50%. The oral pharmacokinetics are slightly more than dose-proportional over the clinical dose range. The AUC and C_{max} of eletriptan are increased by approximately 20 to 30% following oral administration with a high fat meal. Eletriptan hydrobromide can be taken with or without food.

Distribution: The volume of distribution of eletriptan following IV administration is 138L. Plasma protein binding is moderate and approximately 85%.

Metabolism: The N-demethylated metabolite of eletriptan is the only known active metabolite. This metabolite causes vasoconstriction similar to eletriptan in animal models. Though the half-life of the metabolite is estimated to be about 13 hours, the plasma concentration of the N-demethylated metabolite is 10 to 20% of parent drug and is unlikely to contribute significantly to the overall effect of the parent compound.

In vitro studies indicate that eletriptan is primarily metabolized by cytochrome P-450 enzyme CYP3A4 [see CONTRAINDICATIONS (4) and DRUG INTERACTIONS (7.2)].

Elimination: The terminal elimination half-life of eletriptan is approximately 4 hours. Mean renal clearance (CLR) following oral administration is approximately 3.9 L/h. Non-renal clearance accounts for about 90% of the total clearance.

Specific Populations

Age: The pharmacokinetics of eletriptan are generally unaffected by age. Blood pressure was increased to a greater extent in elderly subjects than in young subjects [see USE IN SPECIFIC POPULATIONS (8.5)]. The pharmacokinetic disposition of eletriptan in the elderly is similar to that seen in younger adults.

There is a statistically significant increased half-life (from about 4.4 hours to 5.7 hours) between elderly (65 to 93 years of age) and younger adult subjects (18 to 45 years of age) [see USE IN SPECIFIC POPULATIONS (8.5)].

Gender: The pharmacokinetics of eletriptan are unaffected by gender.

Race: A comparison of pharmacokinetic studies run in western countries with those run in Japan has indicated an approximate 35% reduction in the exposure of eletriptan in Japanese male volunteers compared to western males. Population pharmacokinetic analysis of two clinical studies indicates no evidence of pharmacokinetic differences between Caucasians and non-Caucasian patients.

Menstrual Cycle: In a study of 16 healthy females, the pharmacokinetics of eletriptan remained consistent throughout the phases of the menstrual cycle.

Renal Impairment: There was no significant change in clearance observed in subjects with mild, moderate or severe renal impairment, though blood pressure elevations were observed in this population [see WARNINGS AND PRECAUTIONS (5.8)].

Hepatic Impairment: Subjects with mild or moderate hepatic impairment demonstrated an increase in both AUC (34%) and half-life. The C_{max} was increased by 18%. No dose adjustment is necessary in subjects with mild or moderate hepatic impairment. The effects of severe hepatic impairment on eletriptan metabolism have not been evaluated [see USE IN SPECIFIC POPULATIONS (8.6)].

Drug Interaction Studies

CYP3A4 inhibitors: In vitro studies have shown that eletriptan is metabolized by the CYP3A4 enzyme. A clinical study demonstrated about a 3-fold increase in C_{max} and about a 6-fold increase in the AUC of eletriptan when combined with ketoconazole. The half-life increased from 5 hours to 8 hours and the T_{max} increased from 2.8 hours to 5.4 hours. Another clinical study demonstrated about a 2-fold increase in C_{max} and about a 4-fold increase in AUC when erythromycin was co-administered with eletriptan. It has also been shown that co-administration of verapamil and eletriptan yields about a 2-fold increase in C_{max} and about a 3-fold increase in AUC of eletriptan, and that co-administration of fluconazole and eletriptan yields about a 1.4-fold increase in C_{max} and about a 2-fold increase in AUC of eletriptan.

Eletriptan hydrobromide is contraindicated within at least 72 hours of treatment with the following potent CYP3A4 inhibitors: ketoconazole, itraconazole, nefazodone, troleandomycin, clarithromycin, ritonavir and nelfinavir. Eletriptan hydrobromide should not be used within 72 hours with drugs that have demonstrated potent CYP3A4 inhibition [see CONTRAINDICATIONS (4)].

Propranolol: The C_{max} and AUC of eletriptan were increased by 10 and 33%, respectively, in the presence of propranolol. No interactive increases in blood pressure were observed. No dosage adjustment appears to be needed for patients taking propranolol.

The effect of eletriptan on other drugs: The effect of eletriptan on enzymes other than cytochrome P450 has not been investigated. In vitro human liver microsome studies suggest that eletriptan has little potential to inhibit CYP1A2, 2C9, 2E1 and 3A4 at concentrations up to 100 µM. While eletriptan has an effect on CYP2D6 at high concentration, this effect should not interfere with metabolism of other drugs when eletriptan is used at recommended doses. There is no in vitro or in vivo evidence that clinical doses of eletriptan will induce drug metabolizing enzymes. Therefore, eletriptan is unlikely to cause clinically important drug interactions mediated by these enzymes.

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Eletriptan was administered to rats and mice in the diet for 104 weeks. In rats, the incidence of testicular interstitial cell adenomas was increased at the high dose of 75 mg/kg/day, but not at 15 mg/kg/day, a dose associated with plasma exposures (AUC) approximately 2 times that in humans at the MRHD of 80 mg/day. In mice, the incidence of hepatocellular adenomas was increased at the high dose of 400 mg/kg/day, but not a dose of 90 mg/kg/day, associated with plasma AUC approximately 7 times that in humans at the MRHD.

Mutagenesis

Eletriptan was negative in in vitro (bacteria reverse mutation (Ames), mammalian cell gene mutation (CHO/HGPRT), chromosomal aberration assay in human lymphocytes) and in vivo (mouse micronucleus) assays.

Impairment of Fertility

In a fertility and early embryonic development study, eletriptan (50, 100, or 200 mg/kg/day) was orally administered to male and female rats prior to and throughout mating and continuing in females to implantation. Plasma exposures (AUC) were 4, 8 and 16 times in males and 7, 14 and 28 times in females, respectively, that in humans at the MRHD. Prolongation of the estrous cycle and decreases in the number of corpora lutea, implants, and viable fetuses per dam were observed at 200 mg/kg/day. Male fertility parameters were not affected.

The efficacy of eletriptan hydrobromide in the acute treatment of migraines was evaluated in eight randomized, double-blind placebo-controlled studies. All eight studies used 40 mg. Seven studies evaluated an 80 mg dose and two studies included a 20 mg dose.

In all eight studies, randomized patients treated their headaches as outpatients. Seven studies enrolled adults and one study enrolled adolescents (age 11 to 17). Patients treated in the seven adult studies were predominantly female (85%) and Caucasian (94%) with a mean age of 40 years (range 18 to 78). In all studies, patients were instructed to treat a moderate to severe headache. Headache response, defined as a reduction in headache severity from moderate or severe pain to mild or no pain, was assessed up to 2 hours after dosing. Associated symptoms such as nausea, vomiting, photophobia and phonophobia were also assessed.

Maintenance of response was assessed for up to 24 hours post dose. In the adult studies, a second dose of eletriptan hydrobromide or other medication was allowed 2 to 24 hours after the initial treatment for both persistent and recurrent headaches. The incidence and time to use of these additional treatments were also recorded.

In the seven adult studies, the percentage of patients achieving headache response 2 hours after treatment was significantly greater among patients receiving eletriptan hydrobromide at all doses compared to those

who received placebo. The two-hour response rates from these controlled clinical studies are summarized in Table 2.

Table 2: Percentage of Patients with Headache Response (Mild or No Headache) 2 Hours Following Treatment

Placebo
Eletriptan Hydrobromide
20 mg
Eletriptan Hydrobromide
40 mg
Eletriptan Hydrobromide
80 mg
Study 1
23.8%
54.3%*
65.0%*
77.1%*
(n=126)
(n=129)
(n=117)
(n=118)
Study 2
19.0%
NA
61.6%*
64.6%*
(n=232)
(n=430)
(n=446)
Study 3
21.7%
47.3%*
61.9%*
58.6%*
(n=276)
(n=273)
(n=281)
(n=290)
Study 4
39.5%
NA
62.3%*
70.0%*
(n=86)
(n=175)
(n=170)
Study 5
20.6%
NA
53.9%*
67.9%*
(n=102)
(n=206)
(n=209)
Study 6
31.3%
NA
63.9%*
66.9%*
(n=80)
(n=169)
(n=160)
Study 7
29.5%

NA
57.5%*
NA
(n=122)
(n=492)

* p value < 0.05 vs placebo

NA - Not Applicable

Comparisons of the performance of different drugs based upon results obtained in different clinical trials are never reliable. Because studies are generally conducted at different times, with different samples of patients, by different investigators, employing different criteria and/or different interpretations of the same criteria, under different conditions (dose, dosing regimen, etc.), quantitative estimates of treatment response and the timing of response may be expected to vary considerably from study to study.

The estimated probability of achieving an initial headache response within 2 hours following treatment is depicted in Figure 1.

Figure 1: Estimated Probability of Initial Headache Response Within 2 Hours*

[figure1]

*Figure 1 shows the Kaplan-Meier plot of probability over time of obtaining headache response (no or mild pain) following treatment with eletriptan. The plot is based on 7 placebo-controlled, outpatient trials in adults providing evidence of efficacy (Studies 1 through 7). Patients not achieving headache response or taking additional treatment prior to 2 hours were censored at 2 hours.

For patients with migraine-associated photophobia, phonophobia, and nausea at baseline, there was a decreased incidence of these symptoms following administration of eletriptan hydrobromide as compared to placebo.

Two to 24 hours following the initial dose of study treatment, patients were allowed to use additional treatment for pain relief in the form of a second dose of study treatment or other medication. The estimated probability of taking a second dose or other medications for migraine over the 24 hours following the initial dose of study treatment is summarized in Figure 2.

Figure 2: Estimated Probability of Taking a Second Dose/Other Medication Over the 24 Hours Following the First Dose*

[figure2]

*This Kaplan-Meier plot is based on data obtained in 7 placebo-controlled trials in adults (Studies 1 through 7). Patients were instructed to take a second dose of study medication as follows: a) in the event of no response at 2 hours (studies 2 and 4 to 7) or at 4 hours (study 3); b) in the event of headache recurrence within 24 hours (studies 2 to 7). Patients not using additional treatments were censored at 24 hours. The plot includes both patients who had headache response at 2 hours and those who had no response to the initial dose. It should be noted that the protocols did not allow re-medication within 2 hours post dose.

The efficacy of eletriptan hydrobromide was unaffected by the duration of attack, gender or age of the patient, relationship to menses, or concomitant use of estrogen replacement therapy/oral contraceptives or frequently used migraine prophylactic drugs.

In a single study in adolescents (n=274), there were no statistically significant differences between treatment groups. The headache response rate at 2 hours was 57% for both eletriptan hydrobromide 40 mg tablets and placebo.

Eletriptan hydrobromide tablets containing 20 mg or 40 mg eletriptan (base) as the hydrobromide salt.

Eletriptan Hydrobromide Tablets 20 mg are orange colored, round, biconvex, film-coated tablets debossed with 'EL' on one side and '20' on the other side. They are supplied as follows:

Carton of 6 (1 X 6) Unit-dose Tablets NDC 59651-104-69

Eletriptan Hydrobromide Tablets 40 mg are orange colored, round, biconvex, film-coated tablets debossed with 'EL' on one side and '40' on the other side. They are supplied as follows:

Carton of 6 (1 X 6) Unit-dose Tablets NDC 72189-517-06

Carton of 12 (2 X 6) Unit-dose Tablets NDC 59651-105-93

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

See FDA Approved Patient Labeling (Patient Information)

Myocardial Ischemia and/or Infarction, Prinzmetal's Angina, Other Vasospastic Reactions, and Cerebrovascular Events

Inform patients that eletriptan hydrobromide may cause serious cardiovascular adverse reactions such as myocardial infarction or stroke, which may result in hospitalization and even death. Although serious cardiovascular reactions can occur without warning symptoms, instruct patients to be alert for the signs and symptoms of chest pain, shortness of breath, weakness, slurring of speech, and instruct them to ask for medical advice when observing any indicative sign or symptoms. Instruct patients to seek medical advice if they have symptoms of other vasospastic reactions [see WARNINGS AND PRECAUTIONS (5.1, 5.2, 5.4, 5.5, and 5.8)].

Anaphylactic/Anaphylactoid Reactions

Inform patients that anaphylactic/anaphylactoid reactions have occurred in patients receiving eletriptan hydrobromide. Such reactions can be life threatening or fatal. In general, anaphylactic reactions to drugs are more likely to occur in individuals with a history of sensitivity to multiple allergens [see CONTRAINDICATIONS (4)].

Medication Overuse Headache

Inform patients that use of drugs to treat acute migraines for 10 or more days per month may lead to an exacerbation of headache, and encourage patients to record headache frequency and drug use (e.g., by keeping a headache diary) [see WARNINGS AND PRECAUTIONS (5.6)].

Serotonin Syndrome

Inform patients about the risk of serotonin syndrome with the use of eletriptan hydrobromide or other triptans, particularly during combined use with selective serotonin reuptake inhibitors (SSRIs) or serotonin and norepinephrine reuptake inhibitors (SNRIs) [see WARNINGS AND PRECAUTIONS (5.7) and DRUG INTERACTIONS (7.3)].

Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during therapy [see USE IN SPECIFIC POPULATIONS (8.1)].

Lactation

Inform patients to notify their healthcare provider if they are breastfeeding or plan to breastfeed [see USE IN SPECIFIC POPULATIONS (8.2)].

Patient Information

Eletriptan Hydrobromide Tablets

(EL-e-TRIP-tan HYE-droe-BROE-mide)

Please read this information before you start taking eletriptan hydrobromide tablets and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

What is the most important information I should know about eletriptan hydrobromide tablets?

Eletriptan hydrobromide tablets can cause serious side effects, including:

Heart attack and other heart problems. Heart problems may lead to death.

Stop taking eletriptan hydrobromide tablets and get emergency medical help right away if you have any of the following symptoms of a heart attack:

discomfort in the center of your chest that lasts for more than a few minutes, or that goes away and comes back

chest pain or chest discomfort that feels like an uncomfortable heavy pressure, squeezing, fullness, or pain

shortness of breath with or without chest discomfort

breaking out in a cold sweat

nausea or vomiting

feeling lightheaded

Eletriptan hydrobromide tablets are not for people with risk factors for heart disease unless a heart exam is done and shows no problem. You have a higher risk for heart disease if you:

have high blood pressure

have high cholesterol levels

smoke
are overweight
have diabetes
have a family history of heart disease
are a female who has gone through menopause
are a male over age 40

Serotonin syndrome. Serotonin syndrome is a serious and life-threatening problem that can happen in people taking eletriptan hydrobromide tablets, especially if eletriptan hydrobromide tablets are taken with antidepressant medicines called selective serotonin reuptake inhibitors (SSRIs) or serotonin and norepinephrine reuptake inhibitors (SNRIs).

Ask your doctor or pharmacist for a list of these medicines if you are not sure.

Call your doctor right away if you have any of the following symptoms of serotonin syndrome:

mental changes such as seeing things that are not there (hallucinations), agitation, or coma
fast heartbeat
changes in blood pressure
high body temperature
tight muscles
trouble walking
nausea, vomiting, or diarrhea

What are eletriptan hydrobromide tablets?

Eletriptan hydrobromide tablets are a prescription medicine used to treat acute migraine headaches with or without aura in adults.

Eletriptan hydrobromide tablets are for people who have been diagnosed with migraine headaches.

Eletriptan hydrobromide tablets are not used to prevent or decrease the number of migraine headaches you have.

It is not known if eletriptan hydrobromide tablets are safe and effective to treat cluster headaches.

It is not known if eletriptan hydrobromide tablets are safe and effective in children.

Who should not take eletriptan hydrobromide tablets?

Do not take eletriptan hydrobromide tablets if you:

have heart problems or a history of heart problems
have or have had a stroke or problems with your blood circulation
have hemiplegic or basilar migraines. If you are not sure if you have these types of migraines, ask your doctor
have narrowing of the blood vessels in your legs, arms, stomach, or kidney (peripheral vascular disease)
have ischemic bowel disease
have uncontrolled high blood pressure

have taken any of the following medicines in the last 24 hours:

other "triptans" or triptan combination products such as:

almotriptan (Axert®)
frovatriptan (Frova®)
naratriptan (Amerge®)
rizatriptan (Maxalt®)
sumatriptan and naproxen sodium, (Treximet®)
zolmitriptan (Zomig®)

sumatriptan (Imitrex®)

ergotamines such as:

Bellergal-S®

Cafergot®

Ergomar®

Wigraine®

dihydroergotamines such as:

D.H.E. 45® or Migranal® or methysergide (Sansert®)

have taken the following medicines within the last 72 hours:

ketoconazole (Nizoral®)

itraconazole (Sporanox®)

nefazodone (Serzone®)
troleandomycin (TAO®)
clarithromycin (Biaxin®)
ritonavir (Norvir®)
nelfinavir (Viracept®)

are allergic to eletriptan or any of the ingredients in eletriptan hydrobromide tablets. See the end of this leaflet for a complete list of ingredients in eletriptan hydrobromide tablets.

What should I tell my doctor before taking eletriptan hydrobromide tablets?

Before you take eletriptan hydrobromide tablets, tell your doctor if you:

have heart problems or family history of heart problems or stroke
have high blood pressure
have high cholesterol
have diabetes
smoke
are overweight
are a female who has gone through menopause
have kidney problems
have liver problems
are pregnant or plan to become pregnant. It is not known if eletriptan hydrobromide will harm your unborn baby.
are breastfeeding or plan to breastfeed. Eletriptan hydrobromide passes into your breast milk and may harm your baby. Talk to your doctor about the best way to feed your baby if you take eletriptan hydrobromide tablets.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Your doctor will decide if you can take eletriptan hydrobromide tablets with your other medicines. Eletriptan hydrobromide tablets and other medicines may affect each other causing side effects.

Especially tell your doctor if you take anti-depressant medicines called:

selective serotonin reuptake inhibitors (SSRIs)
serotonin and norepinephrine reuptake inhibitors (SNRIs)

Ask your doctor or pharmacist for a list of these medicines if you are not sure.

Know the medicines you take. Keep a list of them to show your doctor or pharmacist when you get a new medicine.

How should I take eletriptan hydrobromide tablets?

Take eletriptan hydrobromide tablets exactly as your doctor tells you to take it.
Your doctor will tell you how much eletriptan hydrobromide tablets to take and when to take it.
Your doctor may change your dose if needed. Do not change your dose without first talking to your doctor.
Take 1 eletriptan hydrobromide tablet as soon as you feel a migraine coming on.
If you do not get any relief after your first eletriptan hydrobromide tablet, do not take a second tablet without first talking with your doctor.
If your headache comes back or you only get some relief from your headache, you can take a second tablet 2 hours after the first tablet.
Do not take more than a total of 80 mg of eletriptan hydrobromide tablets in a 24-hour period.
If you take too much eletriptan hydrobromide, call your doctor or go to the nearest hospital emergency room right away.
You should write down when you have headaches and when you take eletriptan hydrobromide tablets so you can talk to your doctor about how well eletriptan hydrobromide tablets are working for you.

What should I avoid while taking eletriptan hydrobromide tablets?

Eletriptan hydrobromide tablets can cause dizziness, weakness, or drowsiness. If you have these symptoms, do not drive a car, use machinery, or do anything where you need to be alert.

What are the possible side effects of eletriptan hydrobromide tablets?

Eletriptan hydrobromide tablets may cause serious side effects. See "What is the most important information I should know about eletriptan hydrobromide tablets?"

These serious side effects include:

changes in color or sensation in your fingers and toes (Raynaud's syndrome)

stomach and intestinal problems (gastrointestinal and colonic ischemic events). Symptoms of gastrointestinal and colonic ischemic events include:

- sudden or severe stomach pain
- stomach pain after meals
- weight loss
- nausea or vomiting
- constipation or diarrhea
- bloody diarrhea
- fever

problems with blood circulation to your legs and feet (peripheral vascular ischemia). Symptoms of peripheral vascular ischemia include:

- cramping and pain in your legs or hips
- feeling of heaviness or tightness in your leg muscles
- burning or aching pain in your feet or toes while resting
- numbness, tingling, or weakness in your legs
- cold feeling or color changes in 1 or both legs or feet

medication overuse headaches. Some people who take too many eletriptan hydrobromide tablets may have worse headaches (medication overuse headache). If your headaches get worse, your doctor may decide to stop your treatment with eletriptan hydrobromide tablets.

The most common side effects of eletriptan hydrobromide tablets include:

- dizziness
- nausea
- weakness
- tiredness
- drowsiness

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of eletriptan hydrobromide tablets. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

How should I store eletriptan hydrobromide tablets?

Store eletriptan hydrobromide tablets at room temperature between 20° to 25°C (68° to 77°F).

General information about the safe and effective use of eletriptan hydrobromide tablets

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use eletriptan hydrobromide tablets for a condition for which it was not prescribed. Do not give eletriptan hydrobromide tablets to other people, even if they have the same symptoms you have. It may harm them.

This Patient Information summarizes the most important information about eletriptan hydrobromide tablets. If you would like more information about eletriptan hydrobromide tablets, talk with your doctor. You can ask your doctor or pharmacist for information on eletriptan hydrobromide tablets that is written for health professionals.

For more information call Aurobindo Pharma USA, Inc. at 1-866-850-2876 or go to www.aurobindousa.com.

What are the ingredients in eletriptan hydrobromide tablets?

Active ingredient: eletriptan hydrobromide

Inactive ingredients: croscarmellose sodium, FD&C yellow 6 aluminum lake, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, titanium dioxide and triacetin.

The brands listed are the trademarks of their respective owners and are not trademarks of the Aurobindo Pharma Limited.

Distributed by:

Aurobindo Pharma USA, Inc.
279 Princeton-Hightstown Road
East Windsor, NJ 08520

Manufactured by:

Aurobindo Pharma Limited
Hyderabad-500 038, India

This Patient Information has been approved by the U.S. Food and Drug Administration.

Revised: 04/2020

ELETRIPTAN HBR

eletriptan hbr tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:72189-517(NDC:59651-105)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ELETRIPTAN HYDROBROMIDE (UNII: M41W832TA3) (ELETRIPTAN - UNII:22QO09B8KI)	ELETRIPTAN	40 mg

Inactive Ingredients

Ingredient Name	Strength
TRIACETIN (UNII: XHX3C3X673)	
MICROCRYSTALLINE CELLULOSE 112 (UNII: X7XJ6RM9Q2)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND ((biconvex))	Size	8mm
Flavor		Imprint Code	EL;40
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72189-517-06	6 in 1 CARTON; Type 0: Not a Combination Product	10/11/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210708	10/11/2023	

Labeler - Direct_Rx (079254320)

Registrant - Direct_Rx (079254320)

Establishment

Name	Address	ID/FEI	Business Operations
Direct_Rx		079254320	relabel(72189-517)

Revised: 10/2023

Direct_Rx