

SODIUM CITRATE 4% W/V ANTICOAGULANT- trisodium citrate dihydrate injection, solution

Terumo BCT, Ltd

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP safely and effectively. See full prescribing information for SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP.

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP

Sterile Fluid

Polyvinyl Chloride (PVC) Bag

Initial U.S. Approval: 1978

----- **INDICATIONS AND USAGE** -----

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP is intended for use only for the anticoagulation of whole blood as part of automated apheresis procedures. (1)

----- **DOSAGE AND ADMINISTRATION** -----

- SODIUM CITRATE 4% ANTICOAGULANT SOLUTION USP is added to tubing sets during apheresis procedures. (2)
- SODIUM CITRATE 4% ANTICOAGULANT SOLUTION USP may only be used with apheresis devices. For instructions on the use of the solution see the apheresis device operator's manual. (2.1)
- Follow the directions for connecting the SODIUM CITRATE 4% ANTICOAGULANT SOLUTION USP bag to the apheresis system. (2.2)

----- **DOSAGE FORMS AND STRENGTHS** -----

- 250 mL sterile fluid in a PVC bag. (3)

----- **CONTRAINDICATIONS** -----

- DO NOT INFUSE SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP DIRECTLY TO THE DONOR. (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Verify that the SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP has been securely attached to the Anticoagulant (AC) line on the system tubing set. Use aseptic technique throughout all procedures to ensure donor safety and quality. (5)

----- **ADVERSE REACTIONS** -----

Citrate reactions or toxicity may occur with the infusion and return of blood containing citrate anticoagulant. The recipient of the blood containing citrate should be monitored for the signs and symptoms of citrate toxicity. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Terumo BCT, Inc. at 1-877-339-4228 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **USE IN SPECIFIC POPULATIONS** -----

ANTICOAGULANT SOLUTION has not been studied in controlled clinical trials with specific populations. (7)

Revised: 8/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information

2.2 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
6 ADVERSE REACTIONS
8 USE IN SPECIFIC POPULATIONS
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

16 HOW SUPPLIED/STORAGE AND HANDLING

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION is intended for use only for the anticoagulation of whole blood as part of automated apheresis procedures. *[See Dosage and Administration (2).]*

2 DOSAGE AND ADMINISTRATION

2.1 General Dosing Information

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP is added to tubing sets during apheresis procedures. The solution is connected the tubing set in an apheresis collection. The recommended dose is determined by the apheresis device and metered into the tubing set by the apheresis device. It is not intended for direct intravenous infusion.

For instructions on the use of the solution with the apheresis device and tubing set, see the device operator's manual.

2.2 Administration

- Ensure solution is the SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP and is within the expiration date.
- Inspect the bag. Do not use if the container is damaged, leaking or if there is any visible sign of deterioration.
- Use only if solution is clear and free of particulate matter.
- Protect from sharp objects.

Directions for connecting the SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP bag to the apheresis device.

At the prompt to connect anticoagulant to the apheresis device tubing set:

1. Remove the overwrap by pulling down at the notch, and remove the SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP bag.
2. Before use, perform the following checks *[See Warnings and Precautions (5).]*:
 - Check for leaks by gently squeezing the bag. If leaks are found, discard the bag.
 - Ensure that the solution is the SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP and is within the expiration date.
 - Inspect the solution in good light. Bags showing cloudiness, haze, or particulate matter should not be used.
3. Remove the protective cap from the port on the bag.

4. Connect the bag to the apheresis device tubing set using aseptic technique and hang the solution.
5. Proceed according to the apheresis device operator's manual.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever the solution and container permit.

3 DOSAGE FORMS AND STRENGTHS

250 mL SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP is a sterile solution in a PVC bag. Each 100 mL contains: Sodium Citrate (dihydrate) 4.0 g; and 100 mL Water for Injection, (pH adjusted with citric acid).

4 CONTRAINDICATIONS

DO NOT INFUSE SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP DIRECTLY TO THE DONOR.

5 WARNINGS AND PRECAUTIONS

- Verify that the SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP has been securely attached to the Anticoagulant (AC) line on the system tubing set. Use aseptic technique throughout all procedures to ensure donor safety and quality.
- Do not reuse. Discard unused or partially used solution bags.

6 ADVERSE REACTIONS

Citrate reactions or toxicity may occur with the infusion of blood products to patients and return of blood containing citrate anticoagulant to donors. The recipient of the blood containing citrate should be monitored for the signs and symptoms of citrate toxicity. The signs and symptoms of citrate toxicity begin with paresthesia, a "tingling" sensation around the mouth or in the extremities, followed by severe reactions that are characterized by hypotension and possible cardiac arrhythmia. Citrate toxicity may occur more frequently in patients who are hypothermic, have impaired liver or renal function, or have low calcium levels because of an underlying disease.

8 USE IN SPECIFIC POPULATIONS

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP has not been adequately studied in controlled clinical trials with specific populations.

11 DESCRIPTION

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP is designed to be metered by an apheresis device in apheresis procedures, to prevent platelet activation and coagulation as blood moves throughout the extracorporeal unit (tubing set) in an apheresis procedure.

The solution is sterile and non-pyrogenic, and it contains no bacteriostatic or antimicrobial agents.

The formulas of the active ingredients are provided in Table 1.

Table 1: Active Ingredients

Ingredients	Molecular Formula	Molecular Weight
Sodium Citrate Dihydrate	$C_6H_9Na_3O_9$	294.10

Water for Injection	H ₂ O	18.00
---------------------	------------------	-------

Each 100 mL of SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP contains: Sodium Citrate (dihydrate) 4.0 g; and 100mL Water for Injection (pH adjusted with citric acid).

The PVC bag is not made with natural rubber latex.

The bag is made from a multilayered film. It contains materials that have been tested to demonstrate the suitability of the container for storing pharmaceutical solutions. The bag is nontoxic and biologically inert. The bag-solution unit is a closed system and is not dependent upon entry of external air during administration. The bag is overwrapped to provide protection from the physical environment and to provide an additional moisture barrier when necessary.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP acts as an extracorporeal anticoagulant by binding the free calcium in the blood. Calcium is a necessary co-factor to several steps in the clotting cascade. The following ingredients are key components of the solution:

- Citric acid for pH regulation
- Sodium Citrate anticoagulant

This solution has no pharmacological effect.

16 HOW SUPPLIED/STORAGE AND HANDLING

SODIUM CITRATE 4% W/V ANTICOAGULANT SOLUTION USP is a clear solution supplied in sterile and non-pyrogenic PVC bags. The bags are packaged 30 bags per case.

SIZE	CATALOG NUMBER	NDC NUMBER
250 mL	40881	14537-881-25

STORAGE

Up to 25 °C.
Protect from freezing.

Issued: August 2017

Manufactured by
Terumo BCT, Inc.
Lakewood, CO 80215

PRINCIPAL DISPLAY PANEL - 250 mL Bag Carton Label

Sodium Citrate 4% w/v
Anticoagulant Solution USP

Catalog # 40881
30 x 250 mL units
NDC 14537-881-25

For use only with automated apheresis procedures.

See apheresis device operator's manual for complete instructions.
Read the package insert before application.

Sterile. Non-pyrogenic. Sterilized with steam.

Caution: Do not use unless the solution is clear and the container is intact.
Single use container. Discard any unused product. Not for direct intravenous infusion. Rx Only.

Recommended storage:

Up to 25 °C.

Protect from freezing.

Each 100 mL contains:

Sodium Citrate Dihydrate

4.0 g

Water for Injection to

(pH adjusted with citric acid)

100 mL

Approximate Millimoles:

Sodium Citrate

13.8

Manufactured by Terumo BCT, Inc.

10811 W. Collins Ave., Lakewood CO 80215, USA

777962-240

TERUMOBCT

Lot

Expiry Date

**Sodium Citrate 4% w/v
Anticoagulant Solution USP**

Catalog # 40881

30 x 250 mL units

NDC 14537-881-25

For use only with automated apheresis procedures.

See apheresis device operator's manual for complete instructions.

Read the package insert before application.

Sterile. Non-pyrogenic. Sterilized with steam.

Caution: Do not use unless the solution is clear and the container is intact.

Single use container. Discard any unused product. Not for direct

intravenous infusion. Rx Only.

Recommended storage:

Up to 25 °C.

Protect from freezing.

Each 100 mL contains:

Sodium Citrate Dihydrate 4.0 g

Water for Injection to 100 mL

(pH adjusted with citric acid)

Approximate Millimoles:

Sodium Citrate 13.8

Manufactured by Terumo BCT, Inc.

10811 W. Collins Ave., Lakewood CO 80215, USA

777962-240

TERUMOBCT

Lot

Expiry Date

SODIUM CITRATE 4% W/V ANTICOAGULANT

trisodium citrate dihydrate injection, solution

Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:14537-881	
Route of Administration	INTRAVENOUS			
Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K) (Anhydrous Citric Acid - UNII:XF417D3PSL)	Anhydrous Citric Acid	4 g in 100 mL		
Inactive Ingredients				
Ingredient Name	Strength			
water (UNII: 059QF0KO0R)				
Citric Acid monohydrate (UNII: 2968PHW8QP)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:14537-881-25	30 in 1 CARTON		
1		250 mL in 1 BAG; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	BA125608	06/26/2018		

Labeler - Terumo BCT, Ltd (233649834)

Establishment

Name	Address	ID/FEI	Business Operations
Terumo BCT, Ltd		233649834	MANUFACTURE(14537-881) , ANALYSIS(14537-881) , STERILIZE(14537-881) , LABEL(14537-881)

Revised: 7/2018

Terumo BCT, Ltd