

LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet
Zee Medical Inc

Loperamide Hydrochloride

Drug Facts

Active ingredient (in each caplet)

Loperamide Hydrochloride USP, 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert

Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert

Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store between 20° - 25°C (68° - 77°F)
- see side panel for lot number and expiration date
- **TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**

Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

Questions?

Call toll-free Monday-Friday 8:30 am to 5 pm EST at **1800-406-7984**.

Distributed by
ZEE MEDICAL, INC.,Irvine, CA 92606

PRINCIPAL DISPLAY PANEL - 2 mg Caplet Blister Pack Carton

COMPARE TO THE
ACTIVE INGREDIENT IN
IMODIUM[®]A-D**

ZEE[®]

Loperamide Hydrochloride
Tablets USP, 2 mg

Anti-Diarrheal

See New Warnings
and Directions

12 Caplets*

Each caplet (*capsule-shaped tablet)
contains Loperamide Hydrochloride USP, 2 mg

Controls the Symptoms
of Diarrhea

5197521

Expiration Date:

NON VARNISH

Lot No.:

All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Imodium® A-D.



Loperamide Hydrochloride Tablets USP, 2 mg Anti-Diarrheal

COMPARE TO THE ACTIVE INGREDIENT IN IMODIUM® A-D**



12 Caplets*

Each caplet (*capsule-shaped tablet) contains Loperamide Hydrochloride USP, 2 mg

Controls the Symptoms of Diarrhea

See New Warnings and Directions

Drug Facts

Active ingredient (in each caplet) Loperamide Hydrochloride USP, 2 mg

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings
Allergy alert Do not use if you have ever had a rash or other allergic reaction to loperamide hydrochloride

Heart alert Taking more than directed can cause serious heart problems or death

Drug Facts (continued)

Do not use if you have bloody or black stool

Ask a doctor before use if you have
■ ever
■ mucus in the stool

Ask a doctor or pharmacist before use if you are
■ a history of abnormal heart rhythm
■ a history of liver disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product dizziness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children, in case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

Questions? Call toll-free Monday-Friday 8:30 am to 5 pm EST at 1-800-406-7984.

Drug Facts (continued)

children 9-11 years 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours (60-95 lbs)

children 6-8 years 1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours (48-59 lbs)

children 2-5 years ask a doctor (34 to 47 lbs)

children under 2 years do not use (up to 33 lbs)

Other information

■ store between 20° - 25° C (68° - 77° F)

■ see side panel for lot number and expiration date

■ **TAMPER EVIDENT: THIS PRODUCT IS PROTECTED WITH SEALED BUSTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**


Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hydroxymethylcellulose, pregelatinized starch

24 hours

Keep the carton. It contains important information.

Distributed by ZEE MEDICAL, INC., Irvine, CA 92606 ZEE MEDICAL.COM



6 90689 01457 5



5197521

Loperamide Hydrochloride Tablets USP, 2 mg

Anti-Diarrheal



1457

Rev 0419

LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:35418-123
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77T135393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	green	Score	2 pieces
Shape	CAPSULE (shaped tablet)	Size	9mm
Flavor		Imprint Code	123
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35418-123-	3 in 1 CARTON	01/16/2014	

1	12	2 III 1 CARTON	01/16/2014	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA074091		01/16/2014	

Labeler - Zee Medical Inc (009645623)

Registrant - Ranbaxy Pharmaceuticals Inc (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
OHM LABORATORIES INC.		051565745	manufacture(35418-123)

Revised: 1/2024

Zee Medical Inc