

EQUATE ORASOL- 20% benzocaine gel
WAL-MART STORES INC.

Maximum Strength Equate Orasol

Drug Facts

Active ingredient

Benzocaine 20%

Purpose

Oral Pain Reliever

Uses

- temporary relief of occasional minor irritation, pain and sore mouth

Warnings

Methemoglobinemia Warning: Use of this product may cause Methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops;

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do Not Use

- more than directed
- for more than 7 days unless told to do so by a dentist or doctor

Stop use and ask a doctor if

- swelling, rash or fever develops
- irritation, pain, or redness persists or worsens

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use tube if it is cut prior to opening
- cut open tip of tube on score mark
- use your fingertip or cotton applicator to apply a small pea-size amount of Oral Pain Relief Gel
- apply to affected area up to four times daily or as directed by a dentist or physician
- Adults and children 2 years of age and older: Apply to affected area
- Children under 12 years of age should be supervised in the use of this product
- Children under 2 years of age: Consult a doctor

Inactive ingredients

Benzyl Alcohol, Carbomer, D&C Yellow No 10, FD&C Blue No 1, FD&C Red No 40, Flavor, Glycerin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin

Other information

store at a controlled room temperature 68°- 77°F (20°-25°C)

Principal Display Panel

NDC 79903-966-09

Oral Pain Relief

MAXIMUM STRENGTH GEL

Benzocaine 20% NET WT. 0.33 OZ (9.35g)

equate™
EXPERT

NDC 79903-966-09

Compare to
Maximum
Strength
Anbesol® Gel
active
ingredient*

Instant Pain Relief

Toothaches Canker Sores
Gum Pain Denture Pain

MAXIMUM STRENGTH

Orasol

Benzocaine 20%
Oral Anesthetic Gel

NET WT 0.33 OZ (9.35g)

MAXIMUM STRENGTH

Orasol

Benzocaine 20%
Oral Anesthetic Gel

NET WT 0.33 OZ (9.35g)

LOT:
EXP:

567132



DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716
*This product is not manufactured or distributed by Foundation Consumer Brands, owner of the registered trademark Anbesol®.

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.



Drug Facts

Active ingredient Purpose
Benzocaine 20% Oral anesthetic

Uses for the temporary relief of pain associated with:
• toothache • canker sores • minor dental procedures
• minor injury of the mouth and gums caused by dentures or orthodontic appliances

Warnings METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.

Do not use • for teething • in children under 2 years of age

When using this product • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless told to do so by a dentist or doctor

Inactive ingredients
Benzyl Alcohol, Carboron, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, Flavor, Glycerin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin

Directions
• adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist
• children between 2 and 12 years of age: adult supervision should be given in use of this product
• children under 2 years of age: do not use

Other information • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors • store at a controlled room temperature 68°-77° (20°-25°C)

Keep out of the reach of children.
• irritation, pain, or redness persists or worsens
• swelling, rash or fever develops

Stop use and ask a doctor if
• sore mouth symptoms do not improve in 7 days

Drug Facts (continued)



MAXIMUM STRENGTH
Orasol
 Benzocaine 20%
 Oral Anesthetic Gel
 Instant Pain Relief

NDC 79903-966-09

NET WT
 0.33 OZ (9.35g)

Active ingredient Benzocaine 20% **Purpose** Oral anesthetic **Uses** for the temporary relief of pain associated with: • toothache • canker sores • minor dental procedures • minor injury of the mouth and gums caused by dentures or orthodontic appliances
Warnings **METHEMOGLOBINEMIA WARNING:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy **Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away. **Do not use** • for teething • in children under 2 years of age **When using this product** • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless told to do so by a dentist or doctor **Stop use and ask a doctor if** • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash or fever develops **Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist • children between 2 and 12 years of age: adult supervision should be given in use of this product • children under 2 years of age: do not use **Other information** • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors • store at a controlled room temperature 68°-77°F (20°-25°C) Questions? Call 1-888-287-1915

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

Not For Individual Retail Sale.

290131

734935

EQUATE ORASOL

20% benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-966
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOMER 940 (UNII: 4Q93RCW27E)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-966-09	1 in 1 CARTON	04/30/2024	
1		9.35 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	04/30/2024	

Labeler - WAL-MART STORES INC. (051957769)

Revised: 5/2024

WAL-MART STORES INC.