

MAGNESIUM CITRATE- major magnesium citrate oral solution - lemon liquid
Major Pharmaceuticals

Major Magnesium Citrate Oral Solution - Lemon

Active Ingredient

Magnesium citrate 1.745g per fl oz

Purpose

Saline Laxative

Use

for relief of occasional constipation (irregularity).
Generally produces bowel movement in 1/2 to 6 hours

Warnings

kidney disease
a magnesium restricted diet
abdominal pain, nausea, or vomiting
noticed a sudden change in bowel habits that persists over a period of 2 weeks
already used a laxative for a period longer than 1 week

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any other drug.
Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

STOP USE AND ASK A DOCTOR IF

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

IF PREGNANT OR BREAST-FEEDING

ask health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center right away.

Other Information

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

shake well before using

drink a full glass (8 ounces) of liquid with each dose

may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours

children under 2 years of age - ask a doctor

Other Information

each fl oz contains: magnesium 290 mg

each fl oz contains: sodium 1 mg

Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

MAJOR[®] NDC 0904-7418-44

Magnesium Citrate

Saline Laxative

Lemon Flavor

Alcohol content 0.025%
(From flavoring)

10 FL OZ (296 mL)

Drug Facts	Drug Facts (continued)
Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g Saline laxative	Store at temperatures between 46° and 86°F (8° and 30°C)
Uses: ■ For relief of occasional constipation (irregularity). ■ Generally produces bowel movement in ½ to 6 hours	Directions: ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses
Warnings: Ask a doctor before use if you have ■ Kidney disease ■ a magnesium restricted diet ■ Abdominal pain, nausea, or vomiting ■ Noticed a sudden change in bowel habits that persists over a period of 2 weeks ■ Already used a laxative for a period longer than 1 week	adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children under 2 years of age ask a doctor
Ask a doctor or pharmacist before use if you are ■ Taking any other drug. ■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.	Other Information: ■ each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.	Inactive ingredients: benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶	Questions? 1-833-551-0932

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN, OR SEPARATED FROM NECKRING

Rev. 08/23 M-184 Re-order No. 701382



3 09047 41844 2

Distributed by:
MAJOR[®] PHARMACEUTICALS
Indianapolis, IN 46268
Questions or comments?
Call (800) 616-2471
www.majorpharmaceuticals.com

MAGNESIUM CITRATE

major magnesium citrate oral solution - lemon liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7418
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 29.6 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
WATER (UNII: 059QF0KO0R)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7418-44	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/25/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	10/25/2023	

Labeler - Major Pharmaceuticals (191427277)