

**VAPOR ICE DAYTIME AND NITETIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin  
Meijer, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Meijer Distribution, Inc. Vapor Ice™ Daytime & Nitetime Cold & Flu Drug Facts**

**Active ingredients (in each 30 mL) - Nitetime**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 42 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

## Questions or comments?

**1-800-719-9260**

### **Active ingredients (in each 30 mL) - Daytime**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

### **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may

include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

**do not use more than directed**

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

DAYTIME & NITETIME COMBO PACK

Compare to Vicks® NyQuil® Severe+VapoCOOL™ and DayQuil® Severe+VapoCOOL™

active ingredients

meijer®

Compare to Vicks® DayQuil® Severe+VapoCOOL™ active ingredients

meijer®

MAXIMUM STRENGTH

NON-DROWSY

vapor ice®

daytime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Phenylephrine HCl

Nasal Decongestant

Dextromethorphan HBr

Cough Suppressant

Guaifenesin | Expectorant

SEVERE

ALCOHOL 10%

Compare to Vicks® NyQuil® Severe+VapoCOOL™ active ingredients

meijer®

MAXIMUM STRENGTH

vapor ice®

nighttime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Phenylephrine HCl

Nasal Decongestant

Doxylamine Succinate

Antihistamine

Dextromethorphan HBr

Cough Suppressant

SEVERE

ALCOHOL 10%

Multi-Symptom Relief

2 -12 FL OZ (355 mL) BOTTLES

TOTAL 24 FL OZ (1.5 pt) (710 mL)

meijer

vapor ice<sup>®</sup>  
daytime  
cold & flu

meijer

vapor ice<sup>®</sup>  
nitetime  
cold & flu

SEVERE

Multi-Symptom Relief

SEVERE

Multi-Symptom Relief

DAYTIME & NITETIME  
COMBO PACK

NDC 79481-2800-0

meijer

Compare to Vicks<sup>®</sup> NyQuil<sup>®</sup>  
Severe+ VapoCOOL<sup>™</sup>  
and DayQuil<sup>®</sup> Severe+  
VapoCOOL<sup>™</sup>  
active ingredients\*

meijer

Compare to Vicks<sup>®</sup>  
NyQuil<sup>®</sup> Severe+  
VapoCOOL<sup>™</sup>  
active ingredients\*

MAXIMUM  
STRENGTH

vapor ice<sup>®</sup>  
nitetime  
cold & flu

SEVERE

Multi-Symptom  
Relief

ALCOHOL 10%

meijer

Compare to Vicks<sup>®</sup>  
DayQuil<sup>®</sup> Severe+  
VapoCOOL<sup>™</sup>  
active ingredients\*

MAXIMUM  
STRENGTH  
NON-DROWSY

vapor ice<sup>®</sup>  
daytime  
cold & flu

Acetaminophen  
Pain Reliever | Fever Reducer  
Phenylephrine HCl  
Nasal Decongestant  
Dextromethorphan HBr  
Cough Suppressant  
Guaifenesin | Expectorant

meijer

Compare to Vicks<sup>®</sup>  
NyQuil<sup>®</sup> Severe+  
VapoCOOL<sup>™</sup>  
active ingredients\*

MAXIMUM  
STRENGTH

vapor ice<sup>®</sup>  
nitetime  
cold & flu

Acetaminophen  
Pain Reliever | Fever Reducer  
Phenylephrine HCl  
Nasal Decongestant  
Doxylamine Succinate  
Antihistamine  
Dextromethorphan HBr  
Cough Suppressant

SEVERE

ALCOHOL 10%

SEVERE

ALCOHOL 10%

Multi-Symptom Relief

2-12 FL OZ (355 mL) BOTTLES  
TOTAL 24 FL OZ (1.5 pt) (710 mL)

DO NOT USE IF PRINTED NECKBAND  
IS BROKEN OR MISSING

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com



WWW.MEIJER.COM/  
SATISFACTION

PID 4827306



7 08820 00682 9

28FD2 6E C2

Vapor Ice® Nighttime Severe

Drug Facts	Purpose
<b>Active ingredients (in each 30 mL)</b>	
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 more than 4,000 mg of acetaminophen in 24 hours  
 with other drugs containing acetaminophen  
 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 skin redness, blisters, rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 liver disease, heart disease, high blood pressure, thyroid disease, diabetes, glaucoma, cough that occurs with too much phlegm (mucus), a breathing problem such as emphysema or chronic bronchitis, trouble urinating due to an enlarged prostate gland, persistent or chronic cough such as occurs with smoking, asthma, or emphysema, a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**  
 taking sedatives or tranquilizers, taking the blood-thinning drug warfarin

**When using this product**  
 do not use more than directed, excitability may occur, especially in children, marked drowsiness may occur, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery, alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**  
 you get nervous, dizzy or sleepless, pain, nasal congestion, or cough gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

take only as directed - see Overdose warning	do not exceed 4 doses per 24 hrs
only use the dose cup provided	do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**  
 each 30 mL contains: sodium 42 mg. store at 20-25°C (68-77°F)

**Inactive ingredients** alcohol, anhydrous citric acid, DSC yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

**Questions or comments?** 1-800-719-8280

Vapor Ice® Daytime Severe

Drug Facts	Purpose
<b>Active ingredients (in each 30 mL)</b>	
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passages of bothersome mucus and make coughs more productive

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 more than 4,000 mg of acetaminophen in 24 hours  
 with other drugs containing acetaminophen  
 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 skin redness, blisters, rash  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
 with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 liver disease, heart disease, thyroid disease, diabetes, high blood pressure, trouble urinating due to an enlarged prostate gland, cough that occurs with too much phlegm (mucus), persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**  
 taking the blood-thinning drug warfarin

**When using this product do not use more than directed**

**Stop use and ask a doctor if**  
 you get nervous, dizzy or sleepless, pain, nasal congestion, or cough gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

take only as directed - see Overdose warning	do not exceed 4 doses per 24 hrs
only use the dose cup provided	do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**  
 each 30 mL contains: sodium 41 mg. store at 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** alcohol, anhydrous citric acid, DSC yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

**Questions or comments?** 1-800-719-8280

**meijer** Compare to Vicks® DayQuil® Severe+ VapoCOOL™ active ingredients\*

**MAXIMUM STRENGTH NON-DROWSY**

**vapor ice® daytime cold & flu**

**SEVERE Multi-Symptom Relief**

**ALCOHOL 10%**

**PARENTS:**  
 Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



\*These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks®, HyQuil®, and DayQuil®.

## VAPOR ICE DAYTIME AND NITETIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-2800
---------------------	----------------	---------------------------	----------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-2800-0	1 in 1 PACKAGE; Type 0: Not a Combination Product	09/23/2021	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

## Part 1 of 2

### VAPOR ICE NITETIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution

### Product Information

<b>Item Code (Source)</b>	NDC:79481-0590
---------------------------	----------------

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0590-0	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

## Part 2 of 2

### VAPOR ICE DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

## Product Information

Item Code (Source)	NDC:79481-0940
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0940-0	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/23/2021	

**Labeler** - Meijer, Inc. (006959555)

