

**DAY COLD AND FLU AND NIGHT SEVERE COLD AND FLU- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl, acetaminophen, dextromethorphan hbr, phenylephrine hcl, triprolidine hcl
TARGET CORPORATION**

726R Target Day Cold & Flu and Night Severe Cold & Flu Tablets

Drug Facts

Active ingredients (in each caplet)

Day Cold & Flu

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Active ingredients (in each caplet)

Night Severe Cold & Flu

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg
Triprolidine HCl 1.25 mg

Purposes

Day Cold & Flu

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Purposes

Night Severe Cold & Flu

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant
Antihistamine

Uses

Day Cold & Flu

temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- stuffy nose

- sinus congestion and pressure

temporarily reduces fever

helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Night Severe Cold & Flu

temporarily relieves these common cold and flu symptoms:

- cough
- nasal congestion
- minor aches and pains
- sore throat
- headache
- sneezing
- sinus congestion and pressure
- runny nose
- itching of the nose or throat
- itchy, watery eyes due to hay fever

temporarily reduces fever

controls cough to help you get to sleep

Warnings

Liver warning:This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert:Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (**Night Severe Cold & Flu only**)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (**Night Severe Cold & Flu only**)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Night Severe Cold & Flu only**)

When using this product

- do not use more than directed
- excitability may occur, especially in children (**Night Severe Cold & Flu only**)
- marked drowsiness may occur (**Night Severe Cold & Flu only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Night Severe Cold & Flu only**)
- avoid alcoholic drinks (**Night Severe Cold & Flu only**)
- use caution when driving a motor vehicle or operating machinery (**Night Severe Cold & Flu only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions**Day Cold & Flu**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period

- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Night Severe Cold & Flu

- **do not take more than directed (see Overdose warning)**
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients Day Cold & Flu

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Inactive ingredients Night Severe Cold & Flu

colloidal silicon dioxide, croscarmellose sodium, mica, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid, talc, titanium dioxide, yellow iron oxide

Questions or Comments?

Call **1-877-290-4008**



DAY COLD AND FLU AND NIGHT SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl, acetaminophen, dextromethorphan hbr, phenylephrine hcl, triprolidine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-343
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-343-64	4 in 1 CARTON	05/01/2024	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	4 BLISTER PACK	24 in 6
Part 2	4 BLISTER PACK	16 in 4

Part 1 of 2

DAY COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, coated

Product Information

Item Code (Source)	NDC:11673-347
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)

Product Characteristics

Color	red (Light red to red)	Score	no score
Shape	OVAL (Capsule shaped)	Size	19mm
Flavor		Imprint Code	D1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Part 2 of 2

NIGHT SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl, triprolidine hcl tablet, coated

Product Information

Item Code (Source)	NDC:11673-792
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	1.25 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MICA (UNII: V8A1AW0880)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow (Light golden yellow to Golden yellow)	Score	no score
Shape	OVAL (Capsule shaped)	Size	19mm
Flavor		Imprint Code	N1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(11673-343)