

DAYTIME AND NIGHTTIME SEVERE HONEY COLD AND FLU- daytime - acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl and nighttime - acetaminophen, dextromethorphan hbr, phenylephrine hcl, doxylamine succinate
TARGET CORPORATION

745L Target Daytime Severe Honey and Nighttime Severe Honey Cold & Flu Combo Pack

Nighttime Severe Honey Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks daily while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children	30 mL every 4 hrs
12 yrs & over	
children 6 to under	15 mL every 4 hrs
12 yrs	
children 4 to under	do not use unless
6 yrs	directed by a doctor
children under 4	do not use
yrs	

Other information

- each 15 mL contains: sodium 14 mg
- Store at no greater than 25°C and do not refrigerate.

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6,

flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-877-290-4008

Daytime Severe Honey Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks daily while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product,

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless

- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- Store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-877-290-4008



DAYTIME AND NIGHTTIME SEVERE HONEY COLD AND FLU

daytime - acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl and nighttime - acetaminophen, dextromethorphan hbr, phenylephrine hcl, doxylamine succinate kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-328

Packaging			
#	Item Code	Package Description	Marketing Start Date
1	NDC:11673-328-02	1 in 1 CARTON; Type 0: Not a Combination Product	05/01/2024

Quantity of Parts			
Part #	Package Quantity		Total Product Quantity
Part 1	1 BOTTLE		354 mL
Part 2	1 BOTTLE		354 mL

Part 1 of 2

DAYTIME SEVERE HONEY COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Item Code (Source)	NDC:11673-338
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
GLYCERIN (UNII: PDC6A3C0OX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Part 2 of 2	
NIGHTTIME SEVERE HONEY COLD AND FLU	
acetaminophen, dextromethrophan, doxylamine succinate, phenylephrine hcl liquid	

Product Information	
Item Code (Source)	NDC:11673-322
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

XANTHAN GUM (UNII: TTV12P4NEE)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	

Product Characteristics

Color	brown	Score	
Shape		Size	
Flavor	HONEY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(11673-328)

Revised: 10/2023

TARGET CORPORATION