

DRAMAMINE- dimenhydrinate tablet, chewable
Lil' Drug Store Products, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Dramamine® Chewable Formula

Drug Facts

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Use

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use for children under 2 years of age unless directed by a doctor

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a doctor before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

-
- take 1 to 2 chewable tablets every 4-6

adults and children 12 years and over	<p>hours</p> <ul style="list-style-type: none"> do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	<ul style="list-style-type: none"> give ½ to 1 chewable tablet every 6-8 hours do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	<ul style="list-style-type: none"> give ½ chewable tablet every 6-8 hours do not give more than 1-½ chewable tablets in 24 hours, or as directed by a doctor

Other information

- **Phenylketonurics:** contains phenylalanine 0.84 mg per tablet
- store at room temperature 68°-77°F (20°-25°C)
- **do not use if pouch is opened.**
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, aspartame, FD&C yellow 6 aluminum lake, flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified starch, sorbitol

Questions or comments ?

Call 1-800-382-7219

Repackaged and distributed by:

Lil' Drug Store Products, Inc., 1201 Continental Place NE, Cedar Rapids, IA 52402

PRINCIPAL DISPLAY PANEL - 2 Tablet Carton

#1

**Pharmacist
Recommended
Brand**

Dimenhydrinate
Tablets/Antiemetic

**CHEWABLE FORMULA
Dramamine®**

MOTION SICKNESS RELIEF

DUAL ACTION

**1. TREATS SYMPTOMS
ON THE SPOT**

**2. PREVENTS NAUSEA,
DIZZINESS & VOMITING**

**Convenient
Chewable Tablets**

**2 ORANGE FLAVORED
TABLETS (50 mg EACH)**

2 TABLETS

***Lil'*
Drug Store[®]**

#1 Pharmacist Recommended Brand

Dimenhydrinate Tablets/Antiemetic

CHEWABLE FORMULA

Dramamine®

MOTION SICKNESS RELIEF

DUAL ACTION

1. TREATS SYMPTOMS ON THE SPOT
2. PREVENTS NAUSEA, DIZZINESS & VOMITING



Convenient Chewable Tablets

2 ORANGE FLAVORED TABLETS (50 mg EACH)



2 TABLETS

CHEWABLE FORMULA
Dramamine®
MOTION SICKNESS RELIEF

CHEWABLE FORMULA
Dramamine®
MOTION SICKNESS RELIEF

2 TABLETS

2 TABLETS



DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!

Drug Facts

Drug Facts

Active ingredient (in each tablet)

Dimenhydrinate 50 mg.....Antiemetic

Purpose

Use for prevention and treatment of these symptoms associated with motion sickness: ■ nausea ■ vomiting ■ dizziness

Warnings

Do not use for children under 2 years of age unless directed by a doctor

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a doctor before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

LIFT PANEL FOR CONTINUED DRUG FACTS

TABLETS
2

MOTION SICKNESS RELIEF

Dramamine
CHEWABLE FORMULA

Drug Facts (continued)

Directions

- to prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

adults and children 12 years and over	■ take 1 to 2 chewable tablets every 4-6 hours ■ do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	■ give 1/2 to 1 chewable tablet every 6-8 hours ■ do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	■ give 1/2 chewable tablet every 6-8 hours ■ do not give more than 1-1/2 chewable tablets in 24 hours, or as directed by a doctor

Other information

- **Phenylketonurics:** contains phenylalanine 0.84 mg per tablet
- store at room temperature 68°-77°F (20°-25°C)
- **do not use if pouch is opened.**
- see bottom panel for lot number and expiration date

Inactive ingredients anhydrous citric acid, aspartame, FD&C yellow 6 aluminum lake, flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified starch, sorbitol

Questions or comments ? Call 1-800-382-7219

Product manufactured for:
 Medtech Products Inc., Irvington, NY 10533
 A Prestige Brands Company Made in USA

Repackaged and distributed by:
 Lil' Drug Store Products, Inc., 1201 Continental Place NE, Cedar Rapids, IA 52402
 1-877-507-6516 (M-F 8AM-4:30PM CST) www.lildrugstore.com 97091C-US-12-11



DRAMAMINE

dimenhydrinate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-9709(NDC:63029-902)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Dimenhydrinate (UNII: JB937PER5C) (Diphenhydramine - UNII:8GTS82S83M)	Dimenhydrinate	50 mg

Inactive Ingredients

Ingredient Name	Strength
anhydrous citric acid (UNII: XF417D3PSL)	
aspartame (UNII: Z0H242BBR1)	
FD&C yellow no. 6 (UNII: H77VEI93A8)	
aluminum oxide (UNII: LM26O6933)	
magnesium stearate (UNII: 70097M6I30)	
maltodextrin (UNII: 7CVR7L4A2D)	
sorbitol (UNII: 506T60A25R)	

Product Characteristics

Color	ORANGE	Score	2 pieces
Shape	ROUND (Circular Flat Faced Tablet)	Size	12mm
Flavor	ORANGE	Imprint Code	DRAMAMINE
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9709-1	1 in 1 CARTON		
1		2 in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part336	05/10/2010	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2012

Lil' Drug Store Products, Inc.