

**ANTI DIARRHEAL AND ANTI GAS- loperamide hydrochloride,  
simethicone tablet  
Walgreen Company**

-----  
**Walgreen Co. Anti-Diarrheal + Anti-Gas Drug Facts**

**Active ingredients (in each caplet)**

Loperamide HCl 2 mg

Simethicone 125 mg

**Purposes**

Anti-diarrheal

Anti-gas

**Uses**

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

- if you have bloody or black stool
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

## Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

## Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **do not use if printed blister unit is broken or torn**

## Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

**Questions or comments?**

**1-800-719-9260**

**Package/Label Principal Display Panel**

Walgreens

WALGREENS PHARMACIST RECOMMENDED

Compare to the active ingredients in Imodium® Multi-Symptom Relief

Anti-Diarrheal + Anti-Gas

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg / 125 mg

ANTI-DIARRHEAL / ANTI-GAS

Multi-Symptom Relief Caplet

- Relieves symptoms of diarrhea *plus*: Cramps & Pressure; Bloating; Gas

24 CAPLETS\*

(\*capsule-shaped tablets)

ACTUAL SIZE

Walgreens



# Anti-Diarrheal + Anti-Gas

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg / 125 mg  
ANTI-DIARRHEAL / ANTI-GAS

W3ORG1021-F2

Walgreens



Compare to the active ingredients in Imodium® Multi-Symptom Relief\*

NDC 0363-1830-62

# Anti-Diarrheal + Anti-Gas

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg / 125 mg  
ANTI-DIARRHEAL / ANTI-GAS

Multi-Symptom Relief Caplet

- Relieves symptoms of diarrhea plus: Cramps & Pressure; Bloating; Gas

**24** CAPLETS\*  
(\*capsule-shaped tablets)



ACTUAL SIZE

OR162 94 C2

CODE AREA

OPEN OTHER END

### Drug Facts

**Active ingredients (in each caplet)** Purposes  
Loperamide HCl 2 mg..... Anti-diarrheal  
Simethicone 125 mg..... Anti-gas

**Uses** relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

### Warnings

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.  
**Heart alert:** Taking more than directed can cause serious heart problems or death.

### Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

### Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- drink plenty of clear fluid to help prevent dehydration caused by diarrhea
- take only on an empty stomach, 1 hour before or 2 hours after a meal
- take with a full (8 oz) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### Other information

- each caplet contains: calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- do not use if printed blister unit is broken or torn



CONVENIENT RECYCLING TAB

### Drug Facts (continued)

**Inactive ingredients** croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, polyethylene glycol, stearic acid

Use only as directed. ©2021 Walgreen Co.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com  
©2021 Walgreen Co.

Our pharmacists recommend the Walgreens brand.  
We invite you to compare to national brands.  
\*This product is not manufactured or distributed by  
Johnson & Johnson Consumer Inc., distributor of  
Imodium® Multi-Symptom Relief.  
Guten Free

ITEM 179347 W10106-0222-L



## ANTI DIARRHEAL AND ANTI GAS

loperamide hydrochloride, simethicone tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-1830
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>ANHYDROUS DIBASIC CALCIUM PHOSPHATE</b> (UNII: L11K75P92J)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	LOR1;2;125
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1830-53	12 in 1 CARTON	10/10/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-1830-65	1 in 1 CARTON	10/10/2018	06/30/2021

2		30 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-1830-62	24 in 1 CARTON	06/25/2019	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	10/10/2018	

**Labeler** - Walgreen Company (008965063)

Revised: 5/2022

Walgreen Company