

ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet
Western Family Foods Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Western Family Acetaminophen Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain one or more of these ingredients

Principal Display Panel

See New Warnings

BUY ONE GET ONE FREE

TWO 100 COUNT BOTTLES INSIDE

TOTAL = 200 CAPLETS

EXTRA STRENGTH

ASPIRIN FREE CAPLETS

Acetaminophen

PAIN RELIEVER/FEVER REDUCER

FOR ADULTS

Gluten Free

COMPARE TO TYLENOL® active ingredient

Actual Size

100 CAPLETS – 500 mg EACH



LOT NO.
EXP.
: A5546 3U 05

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over	<ul style="list-style-type: none"> ■ do not take more than directed (see overdose warning) ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Acetaminophen
EXTRA STRENGTH
ASPIRIN FREE CAPLETS
PAIN RELIEVER / FEVER REDUCER FOR ADULTS

100 CAPLETS-500 mg EACH

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

12096-F-PER

0 15400 12096 8

Drug Facts (continued)

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients camauba wax, com starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol®.

Proudly Distributed by Western Family Foods, Inc.
P.O. Box 4057, Portland, OR 97208 USA
Copyright 2014 Marca Registrada
www.westernfamily.com

Acetaminophen
EXTRA STRENGTH
ASPIRIN FREE CAPLETS
PAIN RELIEVER / FEVER REDUCER FOR ADULTS

100 CAPLETS-500 mg EACH

ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55312-484
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg	

Inactive Ingredients	
Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	

STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE (caplet)	Size	16 mm
Flavor		Imprint Code	L484
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55312-484-71	1 in 1 CARTON		
1		50 in 1 BOTTLE		
2	NDC:55312-484-78	1 in 1 CARTON		
2		100 in 1 BOTTLE		
3	NDC:55312-484-90	500 in 1 BOTTLE		
4	NDC:55312-484-62	1 in 1 CARTON		
4		24 in 1 BOTTLE		
5	NDC:55312-484-52	10 in 1 VIAL		
6	NDC:55312-484-02	550 in 1 BOTTLE		
7	NDC:55312-484-82	2 in 1 CARTON		
7		100 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/14/2008	

Labeler - Western Family Foods Inc (192166072)

Revised: 1/2015

Western Family Foods Inc