

**HEALTH MART ADULT ASPIRIN- aspirin tablet, film coated**  
**Strategic Sourcing Services LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**McKesson Adult Aspirin Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily relieves minor aches and pains due to:

- menstrual pain
- headache
- toothache
- colds
- minor pain of arthritis
- muscle pain
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

if you have ever had an allergic reaction to any other pain reliever/fever reducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

### **If pregnant or breast feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

## **Inactive ingredients**

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

## **Questions?**

**1-800-719-9260**

## **Package/Label Principal Display Panel**

Compare to Bayer® Aspirin active ingredient

Adult Aspirin

Pain Reliever/Fever Reducer (NSAID)

Headaches

Everyday aches and pains

Safe pain relief plus lifesaving benefits

Talk to your doctor or other healthcare provider before using this product for your heart.

ACTUAL SIZE

Gluten Free

SEE NEW WARNING

100 Coated Tablets, 325 mg EACH

NDC 62011-0432-1

**HealthMart**  
Compare to Bayer® Aspirin active ingredient

**Adult**

**Aspirin**  
Pain Reliever/Fever Reducer (NSAID)  
Headaches • Everyday aches and pains  
Safe pain relief plus lifesaving benefits

ACTUAL SIZE GLUTEN FREE [SEE NEW WARNING](#)

**100** Coated Tablets, 325 mg EACH

**Drug Facts** (continued)  
get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor

**Drug Facts** (continued)  
Inactive ingredients: corn starch, dibasic calcium phosphate dihydrate, hypromellose, tbc, triacetin

**Questions?**  
1-800-719-9280

**Warnings**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor

: 37J78 W6 F2

NDC 62011-0432-1

**HealthMart**  
Compare to Bayer® Aspirin active ingredient

**Adult**

**Aspirin**  
Pain Reliever/Fever Reducer (NSAID)  
Headaches • Everyday aches and pains  
Safe pain relief plus lifesaving benefits

ACTUAL SIZE GLUTEN FREE [SEE NEW WARNING](#)

**100** Coated Tablets, 325 mg EACH

**Drug Facts** (continued)  
get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor

**Drug Facts** (continued)  
Inactive ingredients: corn starch, dibasic calcium phosphate dihydrate, hypromellose, tbc, triacetin

**Questions?**  
1-800-719-9280

**Warnings**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours; not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor

: 37J78 W6 F2

HEALTH MART ADULT ASPIRIN			
aspirin tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:62011-0432
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
TALC (UNII: 7SEV7J4R1U)			
TRIACETIN (UNII: XHX3C3X673)			
Product Characteristics			
Color	WHITE	Score	no score

<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	L37J
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0432-2	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/04/2020	03/04/2020
2	NDC:62011-0432-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	03/04/2020	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	03/04/2020	

**Labeler** - Strategic Sourcing Services LLC (116956644)