

TYLENOL EXTRA STRENGTH- acetaminophen tablet, film coated
Select Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Tylenol® Extra Strength

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning) adults and children 12 years and over
children under 12 years

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 6 hours while symptoms last ▪ do not take more than 6 caplets in 24 hours, unless directed by a doctor ▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if pouch is opened
- see above middle panel for lot number and expiration date

Inactive ingredients

carnauba wax¹, corn starch¹, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch¹, polyethylene glycol¹, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

1 contains one or more of these ingredients

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

PRINCIPAL DISPLAY PANEL - 2 Caplet Pouch Blister Pack

TYLENOL[®]

FOR ADULTS

Acetaminophen

Pain Reliever

Fever Reducer

Extra Strength

2 Caplets

Single-Pack

500 mg each



DO NOT USE IF PACKAGE OR POUCH IS OPENED

1711946017

Open Flap for **DRUG FACTS** →

READ THE LABEL Read all warnings and directions before use

TYLENOL
Acetaminophen 500 mg
Extra Strength

6 55708 11946 4

Single - Pack: 2 Caplets

Product Repackaged and Distributed with permission of manufacturer by:
Select Corporation, Carrollton, TX 75007

Drug Facts

Active ingredient (in each caplet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease →

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> take 2 caplets every 6 hours while symptoms last do not take more than 6 caplets in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)

Drug Facts (continued)

- do not use if pouch is opened
- see above middle panel for lot number and expiration date

Inactive ingredients

carnauba wax, corn starch, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, modified starch, polyethylene glycol, powdered cellulose, pregelatinized starch, propylene glycol, shellac, sodium starch glycolate, titanium dioxide

*contains one or more of these ingredients

Questions or comments?
call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Contains No Aspirin

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
Visit us at www.tylenol.com
or call toll-free 1-877-TYLENOL
(1-877-895-3665)

Care To Recycle™

1711946017

TYLENOL EXTRA STRENGTH			
acetaminophen tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52904-946(NDC:50580-449)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)		Acetaminophen	500 mg

Inactive Ingredients

Ingredient Name	Strength
carnauba wax (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C red no. 40 (UNII: WZB9127XOA)	
aluminum oxide (UNII: LMI26O6933)	
hypromellose, unspecified (UNII: 3NXW29V3WO)	
magnesium stearate (UNII: 70097M6I30)	
polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)	
powdered cellulose (UNII: SMD1X3XO9M)	
propylene glycol (UNII: 6DC9Q167V3)	
shellac (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
titanium dioxide (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	TYLENOL;500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52904-946-04	1 in 1 BLISTER PACK	03/01/1997	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:52904-946-05	2 in 1 BLISTER PACK	03/01/1997	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:52904-946-20	20 in 1 CARTON	03/01/1997	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:52904-946-25	25 in 1 CARTON	03/01/1997	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:52904-946-30	30 in 1 CARTON	03/01/1997	
5		2 in 1 POUCH; Type 0: Not a Combination Product		
6	NDC:52904-946-50	50 in 1 CARTON	03/01/1997	
6		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	03/01/1997	

Labeler - Select Corporation (053805599)

Revised: 3/2022

Select Corporation