# VIRCIN ADVANCED WART TREATMENT- salicylic acid cream Pure Source, LLC

-----

#### VIRCIN Advanced wart treatment

### **Active Ingredient**

Salicylic acid 17%; in collodion; contains alcohol

### **Purpose**

Wart Remover

#### Uses:

For the removal of plantar warts on the bottom of the foot. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness and the interruption of the footprint pattern. For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance of the surface.

### Warnings

on irritated skin, infected or reddened areas, moles, birthmarks, warts with hair growing from them, genital warts, warts on the face or mucous membrane. diabetes or poor blood circulation. Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes. Stop use and ask a doctor if discomfort persists. Cap tube tightly and store at room temperature away from heat. For external use only.Do not useAsk a doctor before use if you haveAvoid inhaling vapors.Flammable. Keep away from fire and flame.

### Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

#### **Directions:**

Wash affected area, may soak in warm water for 5 minutes. Dry area thoroughly. By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

### **Inactive Ingredients:**

BHT, Ethyl Pyruvate, Glycerin, Povidone Iodine, Propylene Glycol, Transcutol, Triethanolamine

### VIRCIN Advanced wart treatment 0.5oz/14g (65121-397-15)


**DIRECTIONS FOR USE**: By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

ACTIVE INGREDIENT: Salicylic acid 17%; in collodion; contains alcohol. INACTIVE INGREDIENTS: See Carton WARNINGS: For external use only. Do not use on irritated skin, infected or reddened areas, moles, birthmarks, warts with hair growing from them, genital warts, warts on the face or mucous membrane.

Ask a doctor before use if you have diabetes or poor blood circulation. Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes.

Flammable. Keep away from fire and flame. Cap tube tightly and store at room temperature away from heat. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center.

Distributed by Pedicis Research LLC, Radnor, PA 19087. Made in the USA





#### Purpose

Uses: For the removal of plantar warts on the bottom of the foot. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness and the interruption of the footprint pattern. For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance of the surface.

Warnings: For external use only. Do not use on irritated skin, infected or reddened areas, moles, birthmarks, warts with hair growing from them, genital warts, warts on the face or mucous membrane. Ask a doctor before use if you have diabetes or poor blood circulation. Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes. Avoid inhaling vapors. Stop use and ask a doctor if discomfort persists. Flammable. Keep away from fire and flame. Cap tube tightly and store at room temperature away from heat. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions." Wash affected area, may soak in warm water for 5 minutes. Dry area thoroughly. By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

Inactive ingrredients: BHT, Ethyl Pyruvate, Glycerin, Povidone Iodine, Propylene Glycol, Transcutol, Triethanolamine





Net Weight 0.5 oz / 14 gm

DIRECTIONS FOR USE: Wash affected area, may soak in warm water for 5 minutes. Dry area thoroughly. By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

Distributed by Pedicis Research LLC, Radnor, PA 19087. Made in the USA Questions: 1-800-748-6539

Patent Pending

#### VIRCIN ADVANCED WART TREATMENT

salicylic acid cream

#### **Product Information**

**Product Type** HUMAN OTC DRUG Item Code (Source) NDC:65121-397

**TOPICAL Route of Administration** 

### **Active Ingredient/Active Moiety**

**Ingredient Name Basis of Strength** Strength

SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ) SALICYLIC ACID 17 g in 100 g

#### **Inactive Ingredients**

Ingredient Name	Strength
INGRAGENT NAME	STEANATA
	Stichati

**BUTYLATED HYDROXYTOLUENE** (UNII: 1P9D0Z171K)

ETHYL PYRUVATE (UNII: 03098E010B)	
GLYCERIN (UNII: PDC6A3C0OX)	
POVIDONE-IODINE (UNII: 85H0HZU99M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TROLAMINE (UNII: 903K93S3TK)	

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:65121-397- 15	1 in 1 CARTON	02/10/2017			
1		14 g in 1 TUBE; Type 0: Not a Combination Product				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M028	06/12/2014		

## Labeler - Pure Source, LLC (080354456)

Establishment				
Name	Address	ID/FEI	Business Operations	
Pure Source, LLC		080354456	manufacture(65121-397)	

Revised: 11/2023 Pure Source, LLC