

**PROMETHAZINE WITH CODEINE- promethazine hydrochloride and codeine phosphate solution**  
**Lohxa**

-----  
**PROMETHAZINE WITH CODEINE ORAL SOLUTION**

**CV**

**(Promethazine Hydrochloride 6.25 mg/5 mL &**

**Codeine Phosphate 10 mg/5mL)**

**Rx only**

**WARNING: ULTRA-RAPID METABOLISM OF CODEINE AND OTHER RISK FACTORS FOR LIFE-THREATENING RESPIRATORY DEPRESSION IN CHILDREN and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS**

**Ultra-Rapid Metabolism of Codeine and Other Risk Factors for Life-Threatening Respiratory Depression in Children**

**Life-threatening respiratory depression and death have occurred in children who received codeine. Most of the reported cases occurred following tonsillectomy and/or adenoidectomy, and many of the children had evidence of being an ultra-rapid metabolizer of codeine due to a CYP2D6 polymorphism. Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated in children younger than 12 years of age and in children younger than 18 years of age following tonsillectomy and/or adenoidectomy (see CONTRAINDICATIONS). Avoid the use of Promethazine HCl and Codeine Phosphate Oral Solution in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of codeine. (See WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression).**

**Promethazine and Respiratory Depression in Children**

**Postmarketing cases of respiratory depression, including fatalities have been reported with use of promethazine in pediatric patients. Children may be particularly sensitive to the additive respiratory depressant effects when promethazine is combined with other respiratory depressants, including codeine. (See WARNINGS – Promethazine and Respiratory Depression in Children).**

**Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**

**Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death (see WARNINGS, PRECAUTIONS - Drug Interactions). Avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol.**

**DESCRIPTION**


Each 5 mL (one teaspoonful), for oral administration contains: Promethazine hydrochloride 6.25 mg;

codeine phosphate 10 mg in a flavored oral solution base with a pH between 4.7 and 5.2. Alcohol 7%.

Inactive Ingredients: Artificial raspberry flavor, ascorbic acid, citric acid anhydrous, D&C Red #33, dehydrated alcohol, edetate disodium, FD&C Blue #1, glycerin, liquid sugar, methylparaben, purified water, saccharin sodium, sodium benzoate, sodium citrate dihydrate and sodium propionate.


Codeine is one of the naturally occurring phenanthrene alkaloids of opium derived from the opium poppy; it is classified pharmacologically as a narcotic analgesic. Codeine phosphate may be chemically designated as 7,8-Didehydro-4,5 $\alpha$ -epoxy-3-methoxy-17-methylmorphinan-6 $\alpha$ -ol phosphate (1:1)(salt) hemihydrate.

The phosphate salt of codeine occurs as white, needle-shaped crystals or white crystalline powder. Codeine phosphate is freely soluble in water and slightly soluble in alcohol. It has a molecular weight of 406.37, a molecular formula of  $C_{18}H_{21}NO_3 \cdot H_3PO_4 \cdot \frac{1}{2}H_2O$ , and the following structural formula:



Promethazine hydrochloride, a phenothiazine derivative, is chemically designated as ( $\pm$ )-10-[2-(Dimethylamino)propyl] phenothiazine monohydrochloride.

Promethazine hydrochloride occurs as a white to faint yellow, practically odorless, crystalline powder which slowly oxidizes and turns blue on prolonged exposure to air. It is soluble in water and freely soluble in alcohol. It has a molecular weight of 320.88, a molecular formula of  $C_{17}H_{20}N_2S \cdot HCl$ , and the following structural formula:



## CLINICAL PHARMACOLOGY

### Codeine

Narcotic analgesics, including codeine, exert their primary effects on the central nervous system and gastrointestinal tract. The analgesic effects of codeine are due to its central action; however, the precise sites of action have not been determined, and the mechanisms involved appear to be quite complex. Codeine resembles morphine both structurally and pharmacologically, but its actions at the doses of codeine used therapeutically are milder, with less sedation, respiratory depression, and gastrointestinal, urinary, and pupillary effects. Codeine produces an increase in biliary tract pressure, but less than morphine or meperidine. Codeine is less constipating than morphine.

Codeine has good antitussive activity, although less than that of morphine at equal doses. It is used in preference to morphine, because side effects are infrequent at the usual antitussive dose of codeine.

Codeine in oral therapeutic dosage does not usually exert major effects on the cardiovascular system.

Narcotic analgesics may cause nausea and vomiting by stimulating the chemoreceptor trigger zone (CTZ); however, they also depress the vomiting center, so that subsequent doses are unlikely to produce vomiting. Nausea is minimal after usual oral doses of codeine.

Narcotic analgesics cause histamine release, which appears to be responsible for wheals or urticaria sometimes seen at the site of injection on parenteral administration. Histamine release may also produce dilation of cutaneous blood vessels, with resultant flushing of the face and neck, pruritus, and sweating.

Codeine and its salts are well absorbed following both oral and parenteral administration. Codeine is about 2/3 as effective orally as parenterally. Codeine is metabolized primarily in the liver by enzymes of the endoplasmic reticulum, where it undergoes O-demethylation, N-demethylation, and partial conjugation with glucuronic acid. The drug is excreted primarily in the urine, largely as inactive metabolites and small amounts of free and conjugated morphine. Negligible amounts of codeine and its metabolites are found in the feces.

Following oral or subcutaneous administration of codeine, the onset of analgesia occurs within 15 to 30 minutes and lasts for four to six hours.

The cough-depressing action, in animal studies, was observed to occur 15 minutes after oral administration of codeine, peak action at 45 to 60 minutes after ingestion. The duration of action, which is dose-dependent, usually did not exceed 3 hours.

### **Promethazine**

Promethazine is a phenothiazine derivative which differs structurally from the antipsychotic phenothiazines by the presence of a branched side chain and no ring substitution. It is thought that this configuration is responsible for its lack (1/10 that of chlorpromazine) of dopamine antagonist properties.

Promethazine is an H<sub>1</sub> receptor blocking agent. In addition to its antihistaminic action, it provides clinically useful sedative and antiemetic effects.

Promethazine is well absorbed from the gastrointestinal tract. Clinical effects are apparent within 20 minutes after oral administration and generally last four to six hours, although they may persist as long as 12 hours. Promethazine is metabolized by the liver to a variety of compounds; the sulfoxides of promethazine and N-demethylpromethazine are the predominant metabolites appearing in the urine.

### **INDICATIONS AND USAGE**

Promethazine HCl and Codeine Phosphate Oral Solution is indicated for the temporary relief of coughs and upper respiratory symptoms associated with allergy or the common cold.

### **CONTRAINDICATIONS**

Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated in all children younger than 12 years of age. (See **WARNINGS - Ultra- Rapid Metabolism of Codeine and Respiratory**

## **Depression).**

Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated for postoperative management in children younger than 18 years who have undergone tonsillectomy and/or adenoidectomy. (See **WARNINGS - Ultra- Rapid Metabolism of Codeine and Respiratory Depression**).

Codeine is contraindicated in patients with a known hypersensitivity to the drug.

Promethazine is contraindicated in comatose states, and in individuals known to be hypersensitive or to have had an idiosyncratic reaction to promethazine or to other phenothiazines.

Antihistamines and codeine are both contraindicated for use in the treatment of lower respiratory tract symptoms, including asthma

## **WARNINGS**

**(see Boxed Warnings)**

### **Ultra-Rapid Metabolism of Codeine and Other Risk Factors for Life-threatening Respiratory Depression in Children**

Life-threatening respiratory depression and death have occurred in children who received codeine. Codeine is subject to variability in metabolism based upon CYP2D6 genotype (described below), which can lead to an increased exposure to the active metabolite morphine. Based upon post-marketing reports, children younger than 12 years old appear to be more susceptible to the respiratory depressant effects of codeine, particularly if there are risk factors for respiratory depression. For example, many reported cases of death occurred in the post-operative period following tonsillectomy and/or adenoidectomy, and many of the children had evidence of being ultra-rapid metabolizers of codeine. Furthermore, children with obstructive sleep apnea who are treated with codeine for post-tonsillectomy and/or adenoidectomy pain may be particularly sensitive to its respiratory depressant effect.

Because of the risk of life-threatening respiratory depression and death:

- Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated in all children younger than 12 years of age. (See **CONTRAINDICATIONS** ).
- Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated for post-operative management in pediatric patients younger than 18 years of age following tonsillectomy and/or adenoidectomy. (See **CONTRAINDICATIONS** ).
- Avoid the use of Promethazine HCl and Codeine Phosphate Oral Solution in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of codeine. Risk factors include conditions associated with hypoventilation, such as postoperative status, obstructive sleep apnea, obesity, severe pulmonary disease, neuromuscular disease, and concomitant use of other medications that cause respiratory depression.
- When prescribing codeine for adolescents, healthcare providers should choose the lowest effective dose for the shortest period of time and inform patients and caregivers about these risks and the signs of morphine overdose. (See **OVERDOSAGE** ).

### *Nursing Mothers*

At least one death was reported in a nursing infant who was exposed to high levels of morphine in breast milk because the mother was an ultra-rapid metabolizer of codeine. Breastfeeding is not recommended during treatment with Promethazine HCl and Codeine Phosphate Oral Solution. (See **PRECAUTIONS – Nursing Mothers**).

### *CYP2D6 Genetic Variability: Ultra-rapid metabolizer*

Some individuals may be ultra-rapid metabolizers because of a specific CYP2D6 genotype (gene duplications denoted as \*1/\*1xN or \*1/\*2xN). The prevalence of this CYP2D6 phenotype varies widely

and has been estimated at 1 to 10% for Whites (European, North American), 3 to 4% for Blacks (African Americans), 1 to 2% for East Asians (Chinese, Japanese, Korean), and may be greater than 10% in certain racial/ethnic groups (i.e., Oceanian, Northern African, Middle Eastern, Ashkenazi Jews, Puerto Rican). These individuals convert codeine into its active metabolite, morphine, more rapidly and completely than other people. This rapid conversion results in higher than expected serum morphine levels. Even at labeled dosage regimens, individuals who are ultra-rapid metabolizers may have life-threatening or fatal respiratory depression or experience signs of overdose (such as extreme sleepiness, confusion, or shallow breathing). (See **OVERDOSAGE**). Therefore, individuals who are ultra-rapid metabolizers should not use Promethazine HCl and Codeine Phosphate Oral Solution.

### **Promethazine and Respiratory Depression in Children**

- *Children*

Postmarketing cases of respiratory depression, including fatalities have been reported with use of promethazine in pediatric patients. Concomitant administration with other respiratory depressants may increase the risk of respiratory depression. Children may be particularly sensitive to the additive respiratory depressant effects when promethazine is combined with other respiratory depressants, including codeine. See WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression for limitations on the use of Promethazine HCl and Codeine Phosphate Oral Solution in children.

Excessively large dosages of antihistamines, including promethazine hydrochloride, in pediatric patients may cause sudden death (see OVERDOSAGE).

- *Concomitant Conditions and Other Risk Factors*

Avoid use of promethazine in patients at risk for respiratory depression. Risk factors include conditions associated with hypoventilation, such as postoperative status, obstructive sleep apnea, obesity, severe pulmonary disease, neuromuscular disease, and concomitant use of other medications that cause respiratory depression. (See WARNINGS -Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants).

- **Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**

Concomitant use of opioids, including Promethazine HCl and Codeine Phosphate Oral Solution, with benzodiazepines, or other CNS depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Because of these risks, avoid use of opioid cough medications in patients taking benzodiazepines, other CNS depressants, or alcohol (see **Precautions - Drug Interactions**).

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioids alone. Because of similar pharmacologic properties, it is reasonable to expect similar risk with concomitant use of opioid cough medications and benzodiazepines, other CNS depressants, or alcohol.

Advise both patients and caregivers about the risks of respiratory depression and sedation if Promethazine HCl and Codeine Phosphate Oral Solution is used with benzodiazepines, alcohol, or other CNS depressants (see **Precautions – Information for Patients**).

### **CNS Depression and Other CNS Effects**

- *CNS Depression*

Promethazine may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. The impairment may be amplified by concomitant use of other central-nervous-system depressants such as alcohol, sedatives/hypnotics (including barbiturates), narcotics, narcotic analgesics, general anesthetics, tricyclic antidepressants, and tranquilizers; therefore avoid use of Promethazine HCl and Codeine Phosphate Oral Solution in patients on these medications. (See PRECAUTIONS – Information for Patients and Drug Interactions).

- *Convulsions*

Convulsions have occurred with therapeutic doses and overdoses of promethazine hydrochloride in pediatric patients. Promethazine may lower seizure threshold. It should be used with caution in persons with seizure disorders or in persons who are using concomitant medications, such as narcotics or local anesthetics, which may also affect seizure threshold.

- *Dystonias*

Promethazine is a phenothiazine. Phenothiazines are associated with dystonic reactions. In pediatric patients who are acutely ill associated with dehydration, there is an increased susceptibility to dystonias with the use of promethazine HCl.

- **Head Injury And Increased Intracranial Pressure**

The respiratory depressant effects of narcotic analgesics and their capacity to elevate cerebrospinal fluid pressure may be markedly exaggerated in the presence of head injury, intracranial lesions, or a preexisting increase in intracranial pressure. Narcotics may produce adverse reactions which may obscure the clinical course of patients with head injuries.

### **Neuroleptic Malignant Syndrome**

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant

Syndrome (NMS) has been reported in association with promethazine HCl alone or in combination with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis and cardiac dysrhythmias).

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g. pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include 1) immediate discontinuation of promethazine HCl, antipsychotic drugs, if any, and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

Since recurrences of NMS have been reported with phenothiazines, the reintroduction of promethazine HCl should be carefully considered.

Double click here to open Word Application.

Write ordered list in to word document.

Close Word Application.

### **Pulmonary Consideration**

- *Asthma and Other Pulmonary Disease*

Narcotic analgesics or cough suppressants, including codeine, should not be used in patients with severe asthma or those with frequent asthma attacks (see **CONTRAINDICATIONS**).

Nor should they be used in acute febrile illness associated with productive cough or in chronic respiratory disease where interference with ability to clear the tracheobronchial tree of secretions would have a deleterious effect on the patient's respiratory function.

- *Unresponsive Cough*

Dosage of codeine should not be increased if cough fails to respond; an unresponsive cough should be reevaluated in 5 days or sooner for possible underlying pathology, such as foreign body or lower respiratory tract disease.

### **Cardiovascular Effects**

- *Hypotension*

Codeine may produce orthostatic hypotension in ambulatory patients.

### **Bone-Marrow Depression**

Promethazine should be used with caution in patients with bone-marrow depression. Leukopenia and agranulocytosis have been reported, usually when promethazine HCl has been used in association with other known marrow-toxic agents.

### **Constipation**

Codeine may cause or aggravate constipation.

### **Other Considerations**

Administration of promethazine has been associated with reported cholestatic jaundice.

Administration of codeine may be accomplished by histamine release and should be used with caution in atopic children.

## **PRECAUTIONS**

### **General**

Narcotic analgesics, including codeine, should be administered with caution and the initial dose reduced in patients with acute abdominal conditions, convulsive disorders, significant hepatic or renal impairment, fever, hypothyroidism, Addison's disease, ulcerative colitis, prostatic hypertrophy, in patients with recent gastrointestinal or urinary tract surgery, and in the very young or elderly or debilitated patients.

Promethazine should be used cautiously in persons with cardiovascular disease or with impairment of liver function.

### **Anticholinergic Effects**

Drugs having anticholinergic properties should be used with caution in patients with narrow-angle glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction, and bladder-neck obstruction.

### **Information for Patients**

Advise patients of the risks of respiratory depression and death with Promethazine HCl and Codeine Phosphate Oral Solution in children younger than 18 years of age. Advise patients that Promethazine HCl and Codeine Phosphate Oral Solution should not be used in children younger than 12 years of age or in a child younger than 18 years of age for treatment after tonsillectomy and/or adenoidectomy. Advise caregivers of children ages 12 to 18 receiving Promethazine HCl and Codeine Phosphate Oral Solution to monitor for signs of respiratory depression. (See **WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression**).

Advise women that breastfeeding is not recommended during treatment with Promethazine HCl and Codeine Phosphate Oral Solution. (See **WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression**).

Promethazine and codeine may cause marked drowsiness or may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery. Ambulatory patients should be told to avoid engaging in such activities until it is known that they do not become drowsy or dizzy from promethazine and codeine therapy. Pediatric patients should be supervised to avoid potential harm in bike riding or in other hazardous activities.

Inform patients and caregivers that potentially fatal additive effects may occur if Promethazine HCl and Codeine Phosphate Oral Solution is used with benzodiazepines or other CNS depressants, including alcohol. Because of this risk, patients should avoid concomitant use of Promethazine HCl and Codeine Phosphate Oral Solution with benzodiazepines or other CNS depressants, including alcohol (See **WARNINGS- Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**).

Patients should be advised to measure Promethazine HCl and Codeine Phosphate Oral Solution with an accurate measuring device. A household teaspoon is not an accurate measuring device and could lead to overdosage, especially when a half a teaspoon is measured. A pharmacist can recommend an appropriate measuring device and can provide instructions for measuring the correct dose.

Patients should be advised to report any involuntary muscle movements.

Avoid prolonged exposure to the sun.

Codeine, like other narcotic analgesics, may produce orthostatic hypotension in some ambulatory patients. Patients should be cautioned accordingly.

## **Drug Interactions**

Codeine

Codeine

In patients receiving MAO inhibitors, an initial small test dose is advisable to allow observation of any excessive narcotic effects or MAOI interaction.

The use of benzodiazepines, opioids, antihistamines, antipsychotics, anti-anxiety agents, or other CNS depressants (including alcohol) concomitantly with Promethazine HCl and Codeine Phosphate Oral Solution may cause an additive CNS depressant effect, profound sedation, respiratory depression, coma, and death and should be avoided. (See **WARNINGS- Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants**).

Promethazine

*Epinephrine:* Because of the potential for promethazine to reverse epinephrine's vasopressor effect, epinephrine should NOT be used to treat hypotension associated with promethazine overdose.

*Anticholinergics:* Concomitant use of other agents with anticholinergic properties should be undertaken with caution.

*Monoamine Oxidase Inhibitors (MAOI):* Drug interactions, including an increased incidence of extrapyramidal effects, have been reported when some MAOI and phenothiazines are used concomitantly.

## **Drug/Laboratory Test Interactions**

Because narcotic analgesics may increase biliary tract pressure, with resultant increase in plasma amylase or lipase levels, determination of these enzyme levels may be unreliable for 24 hours after a narcotic analgesic has been given.

The following laboratory tests may be affected in patients who are receiving therapy with promethazine hydrochloride:

*Pregnancy Tests:* Diagnostic pregnancy tests based on immunological reactions between HCG and anti-

HCG may result in false-negative or false-positive interpretations.

*Glucose Tolerance Test:* An increase in blood glucose has been reported in patients receiving promethazine.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term animal studies have not been performed to assess the carcinogenic potential of codeine or of promethazine, nor are there other animal or human data concerning carcinogenicity, mutagenicity, or impairment of fertility with these agents. Codeine has been reported to show no evidence of carcinogenicity or mutagenicity in a variety of test systems, including the micronucleus and sperm abnormality assays and the *Salmonella* assay. Promethazine was nonmutagenic in the *Salmonella* test system of Ames.

### **Pregnancy**

Teratogenic Effects

*Pregnancy Category C*

#### Codeine

A study in rats and rabbits reported no teratogenic effect of codeine administered during the period of organogenesis in doses ranging from 5 to 120 mg/kg. In the rat, doses at the 120-mg/kg level, in the toxic range for the adult animal were associated with an increase in embryo resorption at the time of implantation. In another study a single 100-mg/kg dose of codeine administered to pregnant mice reportedly resulted in delayed ossification in the offspring.

There are no studies in humans, and the significance of these findings to humans, if any, is not known.

#### Promethazine

Teratogenic effects have not been demonstrated in rat-feeding studies at doses of 6.25 and 12.5 mg/kg of promethazine HCl. These doses are from approximately 2.1 to 4.2 times the maximum recommended total daily dose of promethazine for a 50-kg subject depending upon the indication for which the drug is prescribed. Daily doses of 25 mg/kg intraperitoneally have been found to produce fetal mortality in rats.

Specific studies to test the action of the drug on parturition, lactation, and development of the animal neonate were not done, but a general preliminary study in rats indicated no effect on these parameters. Although antihistamines have been found to produce fetal mortality in rodents, the pharmacological effects of histamine in the rodent do not parallel those in man. There are no adequate and well-controlled studies of promethazine in pregnant women.

Animal reproduction studies have not been conducted with the drug combination – promethazine and codeine. It is not known whether this drug combination can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Promethazine HCl and Codeine Phosphate Oral Solution should be given to a pregnant woman only if clearly needed.

Promethazine HCl and Codeine Phosphate Oral Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nonteratogenic Effects

Dependence has been reported in newborns whose mothers took opiates regularly during pregnancy. Withdrawal signs include irritability, excessive crying, tremors, hyperreflexia, fever, vomiting, and diarrhea. Signs usually appear during the first few days of life.

Promethazine administered to a pregnant woman within two weeks of delivery may inhibit platelet aggregation in the newborn.

### **Labor and Delivery**

Narcotic analgesics cross the placental barrier. The closer to the delivery and the larger the dose used, the greater the possibility of respiratory depression in the newborn. Narcotic analgesics should be avoided during labor if delivery of a premature infant is anticipated. If the mother has received narcotic analgesics during labor, newborn infants should be observed closely for signs of respiratory depression. Resuscitation may be required (see **OVERDOSAGE**). Limited data suggests that use of promethazine hydrochloride during labor and delivery does not have an appreciable effect on the duration of labor or delivery and does not increase the risk of need for intervention in the newborn. The effect of promethazine and/or codeine on later growth and development of the newborn is unknown.

See Nonteratogenic Effects.

### **Nursing Mothers**

Codeine and its active metabolite, morphine, are present in human milk. There are published studies and cases that have reported excessive sedation, respiratory depression, and death (one case) in infants exposed to codeine via breast milk. Women who are ultra-rapid metabolizers of codeine achieve higher than expected serum levels of morphine, potentially leading to higher levels of morphine in breast milk that can be dangerous in their breastfed infants. In women with normal codeine metabolism (normal CYP2D6 activity), the amount of codeine secreted into human milk is low and dose-dependent. There is no information on the effects of the codeine on milk production. Because of the potential for serious adverse reactions, including excess sedation, respiratory depression, and death in a breastfed infant, advise patients that breastfeeding is not recommended during treatment with Promethazine HCl and Codeine Phosphate Oral Solution. (See **WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression**).

If infants are exposed to Promethazine HCl and Codeine Phosphate Oral Solution through breast milk, they should be monitored for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

It is not known whether promethazine is excreted in human milk.

### **Pediatric Use**

Life-threatening respiratory depression and death have occurred in children who received codeine. (See **WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression**).

Postmarketing cases of respiratory depression, including fatalities have been reported with use of promethazine in pediatric patients. (See **WARNINGS – Promethazine and Respiratory Depression in Children**). Because of the risk of life-threatening respiratory depression and death:

- Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated in all children younger than age 12 years of age. (See **CONTRAINDICATIONS** ).
- Promethazine HCl and Codeine Phosphate Oral Solution is contraindicated for post-operative management in pediatric patients younger than 18 years of age following tonsillectomy and/or adenoidectomy. (See **CONTRAINDICATIONS** ).
- Avoid the use of Promethazine HCl and Codeine Phosphate Oral Solution in adolescents 12 to 18 years of age who have other risk factors that may increase their sensitivity to the respiratory depressant effects of codeine. Risk factors include conditions associated with hypoventilation, such as postoperative status, obstructive sleep apnea, obesity, severe pulmonary disease, neuromuscular disease, and concomitant use of other medications that cause respiratory depression. (See **WARNINGS - Ultra-Rapid Metabolism of Codeine and Respiratory Depression** ).
- When prescribing codeine for adolescents, healthcare providers should choose the lowest effective dose for the shortest period of time and inform patients and caregivers about these risks and the signs of morphine overdose. (See **OVERDOSAGE** ).

### **Geriatric Use**

Clinical studies of Promethazine HCl and Codeine Phosphate Oral Solution did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

Sedating drugs may cause confusion and over-sedation in the elderly; elderly patients generally should be started on low doses of Promethazine Hydrochloride and Codeine Phosphate Oral Solution and observed closely.

## **ADVERSE REACTIONS**

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

You may also report side effects to Wockhardt USA, LLC at 1-800-445-4290.

For more information about Promethazine with Codeine Oral Solution, go to [www.wockhardtdusa.com](http://www.wockhardtdusa.com) or call Wockhardt USA, LLC at 1-800-346-6854.

### **Codeine**

*Central Nervous System:* CNS depression, particularly respiratory depression, and to a lesser extent circulatory depression; light-headedness, dizziness, sedation, euphoria, dysphoria, headache, transient hallucination, disorientation, visual disturbances, and convulsions.

*Cardiovascular:* Tachycardia, bradycardia, palpitation, faintness, syncope, orthostatic hypotension (common to narcotic analgesics).

*Gastrointestinal:* Nausea, vomiting, constipation, and biliary tract spasm. Patients with chronic ulcerative colitis may experience increased colonic motility; in patients with acute ulcerative colitis, toxic dilation has been reported.

*Genitourinary:* Oliguria, urinary retention, antidiuretic effect has been reported (common to narcotic analgesics).

*Allergic:* Infrequent pruritus, giant urticaria, angioneurotic edema, and laryngeal edema.

*Other:* Flushing of the face, sweating and pruritus (due to opiate-induced histamine release); weakness.

### **Promethazine**

*Central Nervous System:* Drowsiness is the most prominent CNS effect of this drug. Sedation, somnolence, blurred vision, dizziness, confusion, disorientation and extrapyramidal symptoms such as oculogyric crisis, torticollis, and tongue protrusion; lassitude, tinnitus, incoordination, fatigue, euphoria, nervousness, diplopia, insomnia, tremors, convulsive seizures, excitation, catatonic-like states, hysteria. Hallucinations have also been reported.

*Cardiovascular:* Increased or decreased blood pressure, tachycardia, bradycardia, faintness.

*Dermatologic:* Dermatitis, photosensitivity, urticaria.

*Hematologic:* Leukopenia, thrombocytopenia, thrombocytopenic purpura, agranulocytosis.

*Gastrointestinal:* Dry mouth, nausea, vomiting, jaundice.

*Respiratory:* Asthma, nasal stuffiness, respiratory depression (potentially fatal) and apnea (potentially fatal). (See **WARNINGS- Promethazine; Respiratory Depression**).

*Other:* Angioneurotic edema. Neuroleptic malignant syndrome (potentially fatal) has also been reported. (See **WARNINGS- Promethazine; Neuroleptic Malignant Syndrome**).

*Paradoxical Reactions:* Hyperexcitability and abnormal movements have been reported in patients following a single administration of promethazine HCl. Consideration should be given to the discontinuation of promethazine HCl and to the use of other drugs if these reactions occur. Respiratory depression, nightmares, delirium and agitated behavior have also been reported in some of these patients.

## **DRUG ABUSE AND DEPENDENCE**

### **Controlled Substance**

Promethazine HCl and Codeine Phosphate Oral Solution is a Schedule V Controlled Substance.

### **Abuse**

Codeine is known to be subject to abuse; however, the abuse potential of oral codeine appears to be quite low. Even parenteral codeine does not appear to offer the psychic effects sought by addicts to the same degree as heroin or morphine. However, codeine must be administered only under close supervision to patients with a history of drug abuse or dependence.

### **Dependence**

Psychological dependence, physical dependence, and tolerance are known to occur with codeine.

## **OVERDOSAGE**

### **Codeine**

Serious overdose with codeine is characterized by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, and sometimes bradycardia and hypotension. The triad of coma, pinpoint pupils, and respiratory depression is strongly suggestive of opiate poisoning. In severe overdosage, particularly by the intravenous route, apnea, circulatory collapse, cardiac arrest, and death may occur. Promethazine is additive to the depressant effects of codeine.

It is difficult to determine what constitutes a standard toxic or lethal dose. However, the lethal oral dose of codeine in an adult is reported to be in the range of 0.5 to 1 gram. Infants and children are believed to be relatively more sensitive to opiates on a body-weight basis. Elderly patients are also comparatively intolerant to opiates.

### **Promethazine**

Signs and symptoms of overdosage with promethazine range from mild depression of the central nervous system and cardiovascular system to profound hypotension, respiratory depression, unconsciousness and sudden death. Other reported reactions include hyperreflexia, hypertonia, ataxia, athetosis and extensor-plantar reflexes (Babinski reflex).

Stimulation may be evident, especially in children and geriatric patients. Convulsions may rarely occur. A paradoxical reaction has been reported in children receiving single doses of 75 mg to 125 mg orally, characterized by hyperexcitability and nightmares.

Atropine-like signs and symptoms - dry mouth, fixed dilated pupils, flushing, as well as gastrointestinal symptoms, may occur.

### **Treatment**

Treatment of overdosage with promethazine and codeine is essentially symptomatic and supportive. Only in cases of extreme overdosage or individual sensitivity do vital signs including respiration,

pulse, blood pressure, temperature, and EKG need to be monitored. Activated charcoal orally or by lavage may be given, or sodium or magnesium sulfate orally as a cathartic. Attention should be given to the reestablishment of adequate respiratory exchange through provision of a patent airway and institution of assisted or controlled ventilation. The narcotic antagonist, naloxone hydrochloride, may be administered when significant respiratory depression occurs with promethazine and codeine; any depressant effects of promethazine are not reversed with naloxone. Diazepam may be used to control convulsions. Avoid analeptics, which may cause convulsions. Acidosis and electrolyte losses should be corrected. A rise in temperature or pulmonary complications may signal the need for institution of antibiotic therapy.

Severe hypotension usually responds to the administration of norepinephrine or phenylephrine. EPINEPHRINE SHOULD NOT BE USED, since its use in a patient with partial adrenergic blockade may further lower the blood pressure.

Limited experience with dialysis indicates that it is not helpful.

## DOSAGE AND ADMINISTRATION

*It is important that Promethazine HCl and Codeine Phosphate Oral Solution is measured with an accurate measuring device (see PRECAUTIONS - Information for Patients). A household teaspoon is not an accurate measuring device and could lead to overdosage, especially when half a teaspoon is to be measured. It is strongly recommended that an accurate measuring device be used. A pharmacist can provide an appropriate device and can provide instructions for measuring the correct dose.*

**Promethazine HCl and Codeine Phosphate is contraindicated in pediatric patients less than 6 years of age, because the combination may cause fatal respiratory depression in this age population.**

The average effective dose is given in the following table:

Adults (12 years of age and over)	5 mL (1 teaspoonful) every 4 to 6 hours, not to exceed 30.0 mL in 24 hours.
-----------------------------------	---

## HOW SUPPLIED

Promethazine with Codeine Oral Solution contains promethazine hydrochloride 6.25 mg/5 mL, codeine phosphate 10 mg/5 mL and alcohol 7 percent.

**Single Use Pre-Filled Unit Dose Syringe.**

**Store at 20° – 25°C (68° – 77°F) [see USP Controlled Room Temperature].**

**Protect from light.**

**Rx Only**

**Distributed By:**

**Lohxa**

**Worcester, MA 01608**

REV. 02-18

## MEDICATION GUIDE

Promethazine (proe METH a zeen) with Codeine (KOE deen) Oral Solution, CV

(Promethazine Hydrochloride and Codeine Phosphate Oral Solution)

## **What is the most important information I should know about Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Promethazine Hydrochloride and Codeine Phosphate Oral Solution can cause children to stop breathing (respiratory depression) and cause death.
  - Do not give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to children younger than 12 years of age, or to children younger than 18 years of age after tonsillectomy or adenoidectomy surgery.
  - You should not give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to children between 12 to 18 years of age who have certain risk factors for breathing problems, including:
    - recent surgery
    - obstructive sleep apnea
    - are obese
    - severe lung problems
    - other medical conditions that can cause difficulty breathing, such as neuromuscular problems
    - taking other medicines that cause breathing problems
- You should not breastfeed during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution because it may cause your child to stop breathing and cause death.
- Severe drowsiness, breathing problems (respiratory depression), coma, and death can happen in adults and children who take Promethazine Hydrochloride and Codeine Phosphate Oral Solution with benzodiazepines, or other central nervous system depressants, including alcohol.
  - Avoid taking other medicines that can cause drowsiness or sleepiness, or drinking alcohol during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Ask your healthcare provider for a list of these medicines if you are not sure.
- Promethazine Hydrochloride and Codeine Phosphate Oral Solution can cause breathing problems (respiratory depression) and drowsiness.
  - Take Promethazine Hydrochloride and Codeine Phosphate Oral Solution exactly as prescribed by your healthcare provider. If you take the wrong dose of Promethazine Hydrochloride and Codeine Phosphate Oral Solution you could overdose and die. See the Medication Guide section called "How should I take Promethazine Hydrochloride and Codeine Phosphate Oral Solution?" for important information about how to take Promethazine Hydrochloride and Codeine Phosphate Oral Solution correctly.
  - Avoid driving a car or operating machinery during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution.
- Call your healthcare provider or get emergency medical help right away if anyone taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution, or your breastfeeding baby has any of the symptoms below:
  - increased sleepiness
  - confusion
  - difficulty breathing
  - shallow breathing
  - limpness
  - your baby has difficulty breastfeeding
- Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution in a safe place away from children. Accidental use by a child is a medical emergency and can cause death. If a child accidentally takes Promethazine Hydrochloride and Codeine Phosphate Oral Solution, get emergency medical help right away.

## **What is Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Promethazine Hydrochloride and Codeine Phosphate Oral Solution is a prescription medicine used

to temporarily treat cough and upper respiratory symptoms, including a stuffy nose (nasal congestion) that you can have with allergies or a common cold. Promethazine Hydrochloride and Codeine Phosphate Oral Solution contains 2 medicines, promethazine and codeine. Promethazine is an antihistamine. Codeine is a narcotic cough suppressant.

- **Promethazine Hydrochloride and Codeine Phosphate Oral Solution is a federal controlled substance (C-V) because it contains codeine that can be abused or lead to dependence.** Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution in a safe place to prevent misuse and abuse. Selling or giving away Promethazine Hydrochloride and Codeine Phosphate Oral Solution may harm others, and is against the law. Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines or street drugs
- Promethazine Hydrochloride and Codeine Phosphate Oral Solution is not for children under 12 years of age.

#### **Who should not take Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- **Do not** give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to children younger than 12 years of age.
- **Do not** give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to children younger than 18 years of any age after tonsillectomy or adenoidectomy surgery.
- **Do not** take Promethazine Hydrochloride and Codeine Phosphate Oral Solution if you are allergic to any of the ingredients in Promethazine Hydrochloride and Codeine Phosphate Oral Solution. See the end of this Medication Guide for a complete list of ingredients in Promethazine Hydrochloride and Codeine Phosphate Oral Solution.
- **Do not** take Promethazine Hydrochloride and Codeine Phosphate Oral Solution if you have asthma.
- **Do not** take Promethazine Hydrochloride and Codeine Phosphate Oral Solution if you have high blood pressure or narrowing of blood vessels to your legs or arms (peripheral vascular insufficiency).
- **Do not** take Promethazine Hydrochloride and Codeine Phosphate Oral Solution if you take a medicine called a Monoamine Oxidase Inhibitor (MAOI).

#### **Before you take Promethazine Hydrochloride and Codeine Phosphate Oral Solution, tell your healthcare provider about all of your medical conditions, including if you:**

- have a drug dependence
- have lung or breathing problems including chronic obstructive pulmonary disease (COPD), or sleep apnea.
- have had a head injury
- have pain in your stomach-area (abdomen)
- have a history of severe or persistent cough
- have prostate problems
- have problems with your urinary tract (urethral stricture) or difficulty urinating
- have had surgery recently or plan to have surgery
- drink alcohol
- have kidney or liver problems
- have thyroid problems including hypothyroidism
- have Addison's disease
- have glaucoma (increased pressure in eyes)
- have diabetes
- have seizures
- have heart problems or blood vessel (cardiovascular) problems including high blood pressure or peripheral vascular insufficiency See " **Who should not take Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**
- have blood disorders including low white blood cells
- are pregnant or plan to become pregnant. It is not known if Promethazine Hydrochloride and Codeine

Phosphate Oral Solution will harm your unborn baby. You and your healthcare provider should decide if you should take Promethazine Hydrochloride and Codeine Phosphate Oral Solution while you are pregnant.

- are breastfeeding or plan to breastfeed. Codeine Phosphate passes into your breast milk and may harm your baby. It is not known if Promethazine Hydrochloride can pass into your breast milk. You should not breastfeed during treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution with certain other medicines can cause side effects or affect how well Promethazine Hydrochloride and Codeine Phosphate Oral Solution or the other medicines work. Do not start or stop other medicines without talking to your healthcare provider. Especially tell your healthcare provider if you:

- take pain medicines such as narcotics
- take cold or allergy medicines that contain antihistamines or cough suppressants
- take medicines for mental illness (anti-psychotics, anti-anxiety)
- drink alcohol
- take medicines for depression, including monoamine oxidase inhibitors (MAOIs) or tricyclic antidepressants (TCAs)
- take medicines for stomach or intestinal problems
- take ergot containing medicines
- take medicines called alpha blockers
- take medicines called beta blockers
- take stimulants or uppers (amphetamines)

Ask your healthcare provider if you are not sure if you take one of these medicines.

### **How should I take Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Take Promethazine Hydrochloride and Codeine Phosphate Oral Solution exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much Promethazine Hydrochloride and Codeine Phosphate Oral Solution to take and when to take it. Do not change your dose without talking to your healthcare provider.
- Ask your pharmacist to give you a measuring device to help you measure the correct amount of Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Do not use a household teaspoon to measure your medicine. You may accidentally take too much.
- **Do not** take more than 30 mL of Promethazine Hydrochloride and Codeine Phosphate Oral Solution in 24 hours.
- If you take too much Promethazine Hydrochloride and Codeine Phosphate Oral Solution, call your healthcare provider or go to the nearest hospital emergency room right away.

### **What should I avoid while taking Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Promethazine Hydrochloride and Codeine Phosphate Oral Solution can cause you to be drowsy. Avoid driving a car or operating machinery during the treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Also, watch your child closely when riding a bike.
- Avoid drinking alcohol during the treatment with Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Drinking alcohol can increase your chances of having serious side effects.
- Avoid spending a long time in sunlight.

**What are the possible side effects of Promethazine Hydrochloride and Codeine Phosphate Oral Solution? Promethazine Hydrochloride and Codeine Phosphate Oral Solution may cause serious side effects, including:**

- See "What is the most important information I should know about Promethazine Hydrochloride and Codeine Phosphate Oral Solution?"
- **Breathing problems (respiratory depression)** which can lead to death. Call your healthcare provider or get emergency treatment right away if you have excessive sleepiness, shallow or slow breathing, or confusion.
- **Physical dependence or abuse.** Take Promethazine Hydrochloride and Codeine Phosphate Oral Solution exactly as your healthcare provider tells you to take it. Stopping Promethazine Hydrochloride and Codeine Phosphate Oral Solution suddenly could cause withdrawal symptoms.
- **Bowel problems** including constipation or stomach pain.
- Increased intracranial pressure.
- **Feeling dizzy, faint or light-headed, especially when you stand up (orthostatic hypotension)** . Light-headedness or fainting may happen when getting up too quickly from a sitting or lying position.
- **Neuroleptic malignant syndrome (NMS) which can lead to death.** Tell you healthcare provider right away if you have any of the following symptoms of NMS:
  - high fever
  - stiff muscles
  - confusion
  - sweating
  - changes in pulse, heart rate, and blood pressure

**The most common side effects of Promethazine Hydrochloride and Codeine Phosphate Oral Solution include:**

- sleepiness
- confusion
- dryness of mouth, nose, throat
- nausea and vomiting
- difficulty urinating
- trouble breathing
- high blood pressure
- restlessness
- fast heart beat

These are not all the possible side effects of Promethazine Hydrochloride and Codeine Phosphate Oral Solution. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA- 1088.

**How should I store Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

- Store Promethazine Hydrochloride and Codeine Phosphate Oral Solution at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution in a tightly closed, child-resistant container and out of the light.
- Safely throw away medicine that is out of date or no longer needed.
- **Keep Promethazine Hydrochloride and Codeine Phosphate Oral Solution and all medicines out of the reach of children.**

**General information about the safe and effective use of Promethazine Hydrochloride and Codeine Phosphate Oral Solution.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not

use Promethazine Hydrochloride and Codeine Phosphate Oral Solution for a condition for which it was not prescribed. Do not give Promethazine Hydrochloride and Codeine Phosphate Oral Solution to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about Promethazine Hydrochloride and Codeine Phosphate Oral Solution that is written for health professionals.

**What are the ingredients in Promethazine Hydrochloride and Codeine Phosphate Oral Solution?**

**Active ingredients:** promethazine hydrochloride and codeine phosphate

**Inactive ingredients:** Artificial raspberry flavor, ascorbic acid, citric acid anhydrous, D&C Red #33, dehydrated alcohol, edetate disodium, FD&C Blue #1, glycerin, liquid sugar, methylparaben, purified water, saccharin sodium, sodium benzoate, sodium citrate dihydrate and sodium propionate.

**Manufactured For:**

Wockhardt USA, LLC

Parsippany, NJ 07054

**Manufactured By:**

Morton Grove Pharmaceuticals, Inc.

Morton Grove, IL 60053

26606P

Rev. 11-17

For more information, go to [www.wockhardtusa.com](http://www.wockhardtusa.com) or call Wockhardt USA, LLC at 1-800-346-6854.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

**PRINCIPAL DISPLAY PANEL**

NDC 70166-555-05

PROMETHAZINE WITH CODEINE ORAL SOLUTION

(Promethazine Hydrochloride 6.25 mg/5mL & Codeine Phosphate 10 mg/5 mL) CV

(Contains alcohol 7.0% v/v)

Rx Only

**Promethazine with Codeine Oral Solution**  
 To deliver: **5 mL**  
 Promethazine HCL, USP 6.25mg  
 Codeine Phosphate, USP 10 mg  
 Alcohol 7.0% v/v  
 For Oral Use Only  
 Exp:01/31/2018 Lot#  
  
 7016655505  
**Rx Only** C-V  
 Store at Room Temperature  
 Rpkg. By: LOHXA Worcester MA 01608

NDC 70166-555-05

RxOnly

**Promethazine with Codeine**  
 Oral Solution

For Oral Use only

6.25mg-10mg/5mL



30 x 5mL Oral Syringes

**Promethazine with Codeine**

Oral Solution

6.25mg-10mg/5mL



LOT: EXP:

Each 5mL contains:

Promethazine Hydrochloride, USP ..... 6.25 mg  
 Codeine Phosphate, USP.....10 mg  
 Alcohol.....7% v/v

Store at Controlled Room Temperature 20 to 25 °C (68 to 77°F)  
 [See USP Controlled Room Temperature].

For Institutional Use  
 See insert for dosage and administration

Repackaged by:  
**Lohxa**

Worcester, MA 01608 U.S.A.



**PROMETHAZINE WITH CODEINE**

promethazine hydrochloride and codeine phosphate solution

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70166-555(NDC:60432-606)
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CV

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>PROMETHAZINE HYDROCHLORIDE</b> (UNII: R61ZEH711I) (PROMETHAZINE - UNII:FF28EJQ494)	PROMETHAZINE HYDROCHLORIDE	6.25 mg in 5 mL
<b>CODEINE PHOSPHATE</b> (UNII: GSL05Y1MN6) (CODEINE ANHYDROUS - UNII:UX6OWY2V7J)	CODEINE PHOSPHATE	10 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM PROPIONATE ANHYDROUS</b> (UNII: 39100PO49R)	
<b>ASCORBIC ACID</b> (UNII: PQ6CK8PD0R)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	

GLYCERIN (UNII: PDC6A3C0OX)	
ALCOHOL (UNII: 3K9958V90M)	
METHYLPARABEN (UNII: A2I8C7H9T)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics			
Color	purple	Score	
Shape		Size	
Flavor	RASPBERRY	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70166-555-05	30 in 1 BOX	02/22/2018	
1		1 in 1 BAG		
1		5 mL in 1 SYRINGE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA088875	12/17/1984	

**Labeler -** Lohxa (079872715)

**Registrant -** Lohxa (079872715)

Establishment			
Name	Address	ID/FEI	Business Operations
Lohxa		079872715	relabel(70166-555) , repack(70166-555)