SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 110 ALABASTER-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 120 IVORY-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 130 OPAL-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 140 PORCELAIN-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 150 LACE-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 160 SHELL-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 210 BIRCH-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 220 LINEN-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 230 ALDER-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 240 QUARTZ-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 250 SAND-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 260 CASHMERE-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 310 SILK-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 320 PINE-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 330 BAMBOO-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 340 OAK-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 350 MAPLE-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 360 CITRINE-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 410 SUNSTONE-
octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 420 BRONZE-
octinoxate and titanium dioxide emulsion
SHISEIDO AMERICAS CORPORATION

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION

Drug Facts

Active ingredients	***Purpose***
Octinoxate 4.9%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: ask a doctor
- **Sun Protection Measures**. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. – 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses

Inactive Ingredients

DIMETHICONE, WATER, GLYCERIN, TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE, PEG-10 DIMETHICONE, BUTYLENE GLYCOL, PHENYL TRIMETHICONE, DIMETHICONE CROSSPOLYMER, BIS-BUTYLDIMETHICONE POLYGLYCERYL-3, SORBITAN SESQUIISOSTEARATE, ISOPROPYL MYRISTATE, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, NACRE POWDER, MAGNESIUM ALUMINOMETASILICATE, POLYQUATERNIUM-51, THYMUS SERPYLLUM EXTRACT, SODIUM ACETYLATED HYALURONATE, ALUMINUM HYDROXIDE, STEARIC ACID, CYCLOPENTASILOXANE, DISTEARDIMONIUM HECTORITE, POLYMETHYLSILSESQUIOXANE, POLYSILICONE-2, TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN, LITHIUM MAGNESIUM SODIUM SILICATE, PEG/PPG-19/19 DIMETHICONE, CITRIC ACID, SODIUM CITRATE, BHT, SILICA, TOCOPHEROL, PHENOXYETHANOL, TITANIUM DIOXIDE, IRON OXIDES, MICA,

Other information

Protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 110 Alabaster

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 120 Ivory

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 130 Opal

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 140 Porcelain

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 150 Lace

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 160 Shell

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients | Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO 東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 210 Birch

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients | Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 220 Linen

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients | Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO 東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 230 Alder

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 240 Quartz

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 250 Sand

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO 東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 260 Cashmere

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients | Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO 東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 310 Silk

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients | Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 320 Pine

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 330 Bamboo

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 340 Oak

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 350 Maple

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 360 Citrine

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients / Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503



PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 410 Sunstone

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients — Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA
See our website for more information.

SELF-REFRESHING
Foundation

SHISEIDO GINZA TOKYO 東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 420 Bronze

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
SELF-REFRESHING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL 1 FL. OZ.

A SHISEIDO first in foundation that refreshes non-stop. ActiveForce technology synchronizes with skin and helps to resist heat, oil, humidity and motion for a fresh, natural finish that lasts all day. 24-hour wear. Flawless. Weightless comfort. Medium, buildable coverage.

DERMATOLOGIST-TESTED. NON-COMEDOGENIC.

Drug Facts

Active ingredients Purpose

OCTINOXATE 4.9% ..... Sunscreen

TITANIUM DIOXIDE 5.8% ..... Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove. ▶

SHISEIDO AMERICAS CORPORATION DIST. NEW YORK, NY 10017 MADE IN U.S.A.

www.shiseido.com

RECOMMENDED SKIN CANCER FOUNDATION DAILY USE

Drug Facts (continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses ▶

http://s1872.com/16072AA

See our website for more information.

SELF-REFRESHING Foundation

SHISEIDO GINZA TOKYO

東京 銀座

SYNCHRO SKIN SELF-REFRESHING

Foundation

Oil-Free

BROAD SPECTRUM SPF 30

Sunscreen

30mL 1 FL. OZ.

Drug Facts (continued)

Inactive ingredients

DIMETHICONE · WATER · GLYCERIN · TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE · PEG-10 DIMETHICONE · BUTYLENE GLYCOL · PHENYL TRIMETHICONE · DIMETHICONE CROSSPOLYMER · BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 · SORBITAN SESQUIISOSTEARATE · ISOPROPYL MYRISTATE · DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER · NACRE POWDER · MAGNESIUM ALUMINOMETASILICATE · POLYQUATERNIUM-51 · THYMUS SERPYLLUM EXTRACT · SODIUM ACETYLATED HYALURONATE · ALUMINUM HYDROXIDE · STEARIC ACID · CYCLOPENTASILOXANE · DISTEARDIMONIUM HECTORITE · POLYMETHYLSILSESQUIOXANE · POLYSILICONE-2 · TRIMETHYLSILOXYSILYLCARBAMOYL PULLULAN · LITHIUM MAGNESIUM SODIUM SILICATE · PEG/PPG-19/19 DIMETHICONE · CITRIC ACID · SODIUM CITRATE · BHT · SILICA · TOCOPHEROL · PHENOXYETHANOL · TITANIUM DIOXIDE · IRON OXIDES · MICA ·

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 110 ALABASTER

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-484
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	

DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-484-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 120 IVORY

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-485
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-485-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 130 OPAL

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-486
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-486-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 140 PORCELAIN

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-487
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	

DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-487-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 150 LACE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-488
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-488-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination		

1		Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 160 SHELL

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-489
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	

LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-489-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 210 BIRCH

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-490
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	

WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-490-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 220 LINEN

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-491
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	

MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-491-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 230 ALDER

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-492
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	

POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-492-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 240 QUARTZ

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-493
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-493-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 250 SAND

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-494
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	

SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-494-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 260 CASHMERE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-495
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	

BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-495-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 310 SILK

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-496
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-496-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 320 PINE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-497
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	

ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-497-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 330 BAMBOO

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-498
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-498-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 340 OAK

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-499
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	

PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-499-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 350 MAPLE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-500
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	

SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-500-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 360 CITRINE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-501

Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:58411-501			

1	NDC:58411-501-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 410 SUNSTONE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-502
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	

DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-502-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

SHISEIDO SYNCHRO SKIN SELF-REFRESHING FOUNDATION 420 BRONZE

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:58411-503
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.69 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	2.001 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
PHENYL TRIMETHICONE (UNII: DR0K5NOJ4R)	
DIMETHICONE CROSSPOLYMER (450000 MPA.S AT 12% IN CYCLOPENTASILOXANE) (UNII: UF7620L1W6)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
SILODRATE (UNII: 9T3UU8T0QK)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
SODIUM ACETYLATED HYALURONATE (UNII: WN66R7GL93)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
LAPONITE (UNII: D703131383)	
PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-503-10	1 in 1 CARTON	09/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/01/2019	

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Establishment

Name	Address	ID/FEI	Business Operations
Shiseido America Inc.		782677132	ANALYSIS(58411-484, 58411-485, 58411-486, 58411-487, 58411-488, 58411-489, 58411-490, 58411-491, 58411-492, 58411-493, 58411-494, 58411-495, 58411-496, 58411-497, 58411-498, 58411-499, 58411-500, 58411-501, 58411-502, 58411-503) , MANUFACTURE(58411-484, 58411-485, 58411-486, 58411-487, 58411-488, 58411-489, 58411-490, 58411-491, 58411-492, 58411-493, 58411-494, 58411-495, 58411-496, 58411-497, 58411-498, 58411-499, 58411-500, 58411-501, 58411-502, 58411-503)

Revised: 12/2024 SHISEIDO AMERICAS CORPORATION