

**PUBLIX FAST RELIEF EFFERVESCENT - ORIGINAL FLAVOR- aspirin tablet,  
effervescent  
PUBLIX SUPER MARKETS INC.**

-----  
**Publix Fast Relief Effervescent - Original Flavor**

**Active ingredient**

**(in each effervescent tablet)**

Buffered aspirin 325 mg **(NSAID)\***

\*Nonsteroidal anti-inflammatory drug

**Purpose**

**(in each effervescent tablet)**

Buffered aspirin 325 mg **(NSAID)\***. . . . . Pain reliever/fever reducer

\*Nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains due to:
- headache ■ muscle pain ■ backache
- toothache ■ menstrual pain ■ colds
- minor arthritis pain
- temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

## **Do not use**

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

## **Ask a doctor before use if**

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

## **Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for ■ diabetes ■ gout ■ arthritis

## **When using this product**

**When using this product** do not exceed recommended dosage

## **Stop use and ask a doctor if**

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

## **If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

## **In case of overdose**

In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking

adults and children 12 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 4 tablets in 24 hours

children under 12 years: consult a doctor

Take this product any time - morning, noon or night - when you need relief from headaches, backaches, body aches or muscle pain.

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.


## **Other information**

■ **each tablet contains:** sodium 524 mg

■ store at room temperature (59°-86°F). Avoid excessive heat.

## **Inactive ingredients**

anhydrous citric acid, sodium bicarbonate



# PUBLIX FAST RELIEF EFFERVESCENT - ORIGINAL FLAVOR

aspirin tablet, effervescent

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:56062-358
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

Ingredient Name	Strength

<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>		<b>Imprint Code</b>	L421
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56062-358-36	18 in 1 CARTON	12/18/2023	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/18/2023	

**Labeler** - PUBLIX SUPER MARKETS INC. (006922009)

Revised: 2/2024

PUBLIX SUPER MARKETS INC.