

HISTEX PD DROPS- triprolidine hydrochloride syrup
Allegis Pharmaceuticals, LLC

HISTEX™ PD Drops

Drug Facts

Active ingredient (in each 1 mL dropperful)

Triprolidine HCl 0.938 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if the child has

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor before use if the child is taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- sedatives and tranquilizers may increase the drowsiness effect

Stop use and ask a doctor if

- new symptoms occur

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed recommended dosage.

- For dosing, use only the enclosed dropper and not with any other drug product.

AGE	DOSE
Adults & Children 12 years of age or older:	2.67 mL (2.5 milligrams) every 4 to 6 hours, not to exceed 10.67 mL (10 milligrams in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	1.33 mL (1.25 milligrams) every 4 to 6 hours, not to exceed 5.33 mL (5 milligrams in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Other Information

Store at 15°-30° C (59°-86° F).

Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.

Inactive ingredients

bubble gum flavor, citric acid, glycerin, methylparaben, monoammonium glycyrrhizinate, potassium citrate, potassium sorbate, propylene glycol, propylparaben, purified water, sucralose.

Questions? Comments?

Call 1-866-633-9033.

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Label

NDC 28595-801-30

HISTEX™ PD

Drops

Antihistamine

Each dropperful (1 mL)

contains:

Triprolidine HCl 0.938 mg

Sugar-Free • Dye Free

Alcohol Free

Bubble Gum Flavor

1 fl oz (30 mL)

Drug Facts	
Active ingredient (in each 1 mL dropperful) Triprolidine HCl 0.938 mg.....	Purpose Antihistamine
Uses Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies: <input type="checkbox"/> runny nose, sneezing, <input type="checkbox"/> itching of the nose or throat, <input type="checkbox"/> itchy, watery eyes	
Warnings Do not exceed recommended dosage.	
Ask a doctor before use if the child has <input type="checkbox"/> a breathing problem such as emphysema or chronic bronchitis <input type="checkbox"/> glaucoma	
Ask a doctor before use if the child is taking sedatives or tranquilizers	
When using this product <input type="checkbox"/> excitability may occur, especially in children <input type="checkbox"/> may cause drowsiness <input type="checkbox"/> sedatives and tranquilizers may increase the drowsiness effect	
Stop use and ask a doctor if <input type="checkbox"/> new symptoms occur	
Keep out of the reach of children.	
In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions Do not exceed recommended dosage. <input type="checkbox"/> use only with enclosed dropper <input type="checkbox"/> do not use enclosed dropper for any other drug product	
AGE	DOSE
Adults & Children 12 years of age or older:	2.67 mL (2.5 milligrams) every 4 to 6 hours, not to exceed 10.67 mL (10 milligrams) in 24 hours, or as directed by a doctor.
Children 6 to under 12 years of age:	1.33 mL (1.25 milligrams) every 4 to 6 hours, not to exceed 5.33 mL (5 milligrams) in 24 hours, or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Drug Facts (continued)
Other Information Store at 15°-30° C (59°-86° F). Tamper evident by foil seal under cap. Do not use if foil seal is missing or broken.
Inactive ingredients bubble gum flavor, citric acid, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose.
Questions? Comments? Call 1-866-633-9033.

NDC 28595-801-30

HISTEX™ PD Drops

Antihistamine

Each dropperful (1 mL) contains:
Triprolidine HCl 0.938 mg

Sugar-Free • Dye Free
Alcohol Free

Bubble Gum Flavor

Tamper Evident: Do not use if seal is broken or missing

1 fl oz (30 mL)

KEEP OUT OF REACH OF CHILDREN.

Manufactured for:
Allegis Pharmaceuticals, LLC
Canton, MS 39046
www.allegispharma.com

Rev. Date: 04 /22

CALIBRATED
DROPPER
ENCLOSED



HISTEX PD DROPS

triprolidine hydrochloride syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:28595-801
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII: 2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	0.938 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM BENZOATE (UNII: OJ245FE5EU)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

SUCRALOSE (UNII: 96K6UQ3ZD4)

SORBITOL (UNII: 506T60A25R)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:28595-801-30	1 in 1 CARTON	03/06/2014	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/06/2014	

Labeler - Allegis Pharmaceuticals, LLC (792272861)

Revised: 1/2024

Allegis Pharmaceuticals, LLC