

COPPERTONE KIDS SUNSCREEN SPF 50- (to deliver) avobenzone 3%, homosalate 5%, octisalate 10%, octocrylene 5% spray
Beiersdorf Inc

Coppertone Kids Spray SPF 50

Drug Facts

Active ingredients

(To Deliver) Avobenzone 3%, Homosalate 10%, Octisalate 5%, Octocrylene 5%

Purpose

Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Flammable: Do not use near heat, flame, or while smoking

Do not use on damaged or broken skin

When using this product

- keep away from face to avoid breathing it
- keep out of eyes. Rinse with water to remove.
- use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.
- do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120°F.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before each use

- spray liberally and spread evenly by hand 15 minutes before sun exposure
- hold container 4 to 6 inches from the skin to apply
- do not spray directly into face. Spray on hands then apply to face.
- do not apply in windy conditions
- use in a well-ventilated area
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
 - children under 6 months: Ask a doctor

Other information

- protect this product from excessive heat and direct sun
- store between 20° to 25°C (68° to 77°F)
- may stain or damage some fabrics or surfaces

Inactive ingredients

SD alcohol 40-B (42.8% v/v), isobutane, neopentyl glycol diheptanoate, polyester-27, acrylates/octylacrylamide copolymer, dimethicone, fragrance

Questions?

1-866-288-3330

Copperton Sunscreen Spray

Kids 50

Says on Strong Even in Water Free of Oxybenzone, Octinoxate, Dyes & PABA

Water Resistant (80 Minutes)

Broad Spectrum SPF 50

**25%
MORE**

BONUS 6.9 oz
vs 5.5 oz

Coppertone
SUNSCREEN SPRAY



KiDs

50

**STAYS ON STRONG
EVEN IN WATER**

Free of Oxybenzone,
Octinoxate, Dyes & PABA

Water Resistant (80 Minutes)
Broad Spectrum SPF 50

NET WT 6.9 OZ (196 g)

Broad Spectrum UVA/UVB protection in a durable formula that keeps up with active kids. #1 Pediatrician Recommended Sunscreen Brand.

Drug Facts

Active ingredients

(To Deliver) Avobenzone 3%, Homosalate 10%, Octisalate 5%, Octocrylene 5%Sunscreen

Purpose

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Flammable: Do not use near heat, flame, or while smoking

Do not use on damaged or broken skin

When using this product

- keep away from face to avoid breathing it
- keep out of eyes. Rinse with water to remove.
- use only as directed. Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.
- do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120°F.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions

- shake well before each use
- spray liberally and spread evenly by hand 15 minutes before sun exposure
- hold container 4 to 6 inches from the skin to apply
- do not spray directly into face. Spray on hands then apply to face.
- do not apply in windy conditions
- use in a well-ventilated area
- reapply:
 - after 80 minutes of swimming or sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
 - children under 6 months: Ask a doctor

Other information

- protect this product from excessive heat and direct sun
- store between 20° to 25°C (68° to 77°F)
- may stain or damage some fabrics or surfaces

Inactive ingredients

SD alcohol 40-B (42.8% v/v), isobutane, neopentyl glycol diheptanoate, polyester-27, acrylates/octylacrylamide copolymer, dimethicone, fragrance

Questions? 1-866-288-3330



COPPERTONE KIDS SUNSCREEN SPF 50

(to deliver) avobenzone 3%, homosalate 5%, octisalate 10%, octocrylene 5% spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66800-5353
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	5 g in 100 g
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 g
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	10 g in 100 g
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
POLYESTER-7 (UNII: 0841698D2F)	
ACRYLATES/OCTYLACRYLAMIDE COPOLYMER (40000 MW) (UNII: 7LL6SY9YFV)	
ALCOHOL (UNII: 3K9958V90M)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
NEOPENTYL GLYCOL DIHEPTANOATE (UNII: 5LKW3C543X)	
FRAGRANCE FLORAL ORC0902236 (UNII: R66Z4YW3X0)	
ISOBUTANE (UNII: BXR49TP611)	

Product Characteristics

Color	white (White to Clear)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66800-5353-5	156 g in 1 CAN; Type 0: Not a Combination Product	11/30/2023	
2	NDC:66800-5353-6	196 g in 1 CAN; Type 0: Not a Combination Product	11/30/2023	
3	NDC:66800-5353-7	235 g in 1 CAN; Type 0: Not a Combination Product	11/30/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	11/30/2023	

