

**ALLERGY RELIEF- acetaminophen, diphenhydramine hcl, phenylephrine hcl tablet, film coated**  
**Walgreen Company**

-----  
**Walgreens 44-543 AR**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Diphenhydramine HCl 25 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever  
Antihistamine  
Nasal decongestant

***Uses***

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - headache
  - nasal congestion
  - sneezing
  - runny nose
  - sinus congestion and pressure
  - itchy, watery eyes
  - itching of the nose or throat
  - minor aches and pains

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- glaucoma
- heart disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- liver disease
- thyroid disease
- diabetes

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- see end flap for expiration date and lot number
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients**

corn starch, croscarmellose sodium, crospovidone, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, triacetin

**Questions or comments?**

**1-800-426-9391**

**Principal display panel****Walgreens**

WALGREENS PHARMACIST RECOMMENDED†

NDC 0363-5430-09

**Allergy Relief****SEVERE ALLERGY & SINUS HEADACHE**

**ACETAMINOPHEN** / PAIN RELIEVER

DIPHENHYDRAMINE HCl / ANTIHISTAMINE

PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

- Relief of headache, sinus congestion, sinus pain & pressure, runny nose, sneezing, itchy throat & itchy, watery eyes

**20 CAPLETS**

ACTUAL SIZE

DISTRIBUTED BY: **WALGREEN CO.**

**200 WILMOT RD., DEERFIELD, IL 60015**

**100% SATISFACTION GUARANTEED**

**walgreens.com**

©2021 Walgreen Co.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING**

50844    ORG091854309

†Our pharmacists  
recommend the  
Walgreens brand.

Walgreens



# Allergy Relief

SEVERE ALLERGY & SINUS HEADACHE

ACETAMINOPHEN / PAIN RELIEVER  
DIPHENHYDRAMINE HCl / ANTIHISTAMINE  
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

NDC 0363-5430-09

Walgreens



# Allergy Relief

B-2201-543-09-HR  
ORG091854309



Our pharmacists recommend the Walgreens brand.

DISTRIBUTED BY: WALGREEN CO.  
200 WILLMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com  
©2021 Walgreen Co.

NO PRINT/VARNISH  
LOT & EXP DATE ONLY

50844 ORG091854309

### Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

### Active ingredients (in each caplet)

Acetaminophen 325 mg Pain Reliever  
Diphenhydramine HCl 25 mg Antihistamine  
Phenylephrine HCl 5 mg Nasal decongestant

### Uses

Temporarily relieves these symptoms of hay fever or other upper respiratory allergies: sneezing, runny nose, sinus congestion and pressure, itchy, watery eyes, itching of the nose or throat, minor aches and pains

### Warnings

Liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. 3 or more alcoholic drinks every day while using this product. With other drugs containing acetaminophen. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

### Do not use

With any other drug containing acetaminophen (prescription or nonprescription) if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. With any other product containing diphenhydramine, even one used on skin. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or mental conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients.

### Drug Facts (continued)

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN. Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F).

Inactive ingredients: corn starch, croscarmellose

### Drug Facts (continued)

sodium, croscarmellose, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, silicon dioxide, stearic acid, titanium dioxide, thacetin

(Questions or comments? 1-800-426-9391)

### Other information

See end flap for expiration date and lot number. Children under 12 years: ask a doctor. Do not take more than 10 caplets in 24 hours. Take 2 caplets every 4 hours. Adult and children 12 years and over: do not take more than directed. Do not take more than directed.

### Directions

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### When using this product

Do not exceed recommended dosage. Excitability may occur, especially in children. Marked drowsiness may occur. Alcohol, sedatives, and tranquilizers may increase drowsiness. Use caution when driving a motor vehicle or operating machinery. Avoid alcoholic beverages.

### Stop use and ask a doctor if

fever gets worse or lasts more than 3 days, pain or nasal congestion gets worse or lasts more than 7 days, nervousness, dizziness, or sleeplessness occur. If pregnant or breast-feeding, ask a health professional before use.

### Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Drug Facts (continued)

Ask a doctor before use if you have: glaucoma, heart disease, high blood pressure, difficulty in urination due to enlargement of the prostate gland, a breathing problem such as emphysema or chronic bronchitis, liver disease, thyroid disease, diabetes.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin, taking sedatives or tranquilizers, do not exceed recommended dosage, excitability may occur, especially in children, marked drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, use caution when driving a motor vehicle or operating machinery, avoid alcoholic beverages.

Stop use and ask a doctor if: nervousness, dizziness, or sleeplessness occur, pain or nasal congestion gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur. If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take more than directed. Do not take more than directed.

Adult and children 12 years and over: do not take more than directed. Do not take more than directed.


Children under 12 years: ask a doctor. Do not take more than 10 caplets in 24 hours. Take 2 caplets every 4 hours.

See end flap for expiration date and lot number.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

ITEM 555466 W00000-0000-0  
3 1 1917 108636 1

W3ORGO821-F2



Walgreens 44-543 ORG0918

## ALLERGY RELIEF

acetaminophen, diphenhydramine hcl, phenylephrine hcl tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-5430
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRACETIN</b> (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
--------------	------	--------------	----------

<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	44;543
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5430-09	2 in 1 CARTON	11/04/2021	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/04/2021	

**Labeler** - Walgreen Company (008965063)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-5430) , pack(0363-5430)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-5430)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-5430)

Revised: 11/2024

Walgreen Company