PHILLIPS MILK OF MAGNESIA CONCENTRATED FRESH STRAWBERRY- phillips milk of magnesia concentrated fresh strawberry liquid Bayer HealthCare LLC.

Phillips' Milk of Magnesia Concentrated - Fresh Strawberry

Drug Facts

Active ingredient (in each 15 mL)

Magnesium hydroxide 2400 mg

Saline laxative

Use

relieves occasional constipation (irregularity). This product usually produces bowel movement in $\frac{1}{2}$ to 6 hours.

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed

by a doctor. Drink a full glas (8 oz) of liquid with each dose.

- for accurate dosing, only use the dosing cup provided
- mL = milliliter
- Adults and children 12 years and older: 15 mL to 30 mL
- Children under 12 years: ask a doctor

Other information

- each 15 mL contains: magnesium 1000 mg
- store at room temperature. Avoid freezing. Close cap tightly after use.

Inactive ingredients

anhydrous citric acid, D&C red #28, flavors, glycerin, propylene glycol, purified water, sodium hypochlorite, sorbitol, sucrose

Questions or comments? 1-800-986-0369 (Mon - Fri 9AM - 5PM EST)



Drug Facts

Active ingredient (in each 15 mL) Magnesium hydroxide 2400 mg.....

Purpose Saline faxative

Use relieves occasional constipation (irregularity). This product usually produces bowel movement in 1/2 to 6 hours.

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
 a sudden change in bowel habits that lasts over 14 days.

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 oz) of liquid with each dose.
- for accurate dosing, only use the dosing cup provided
- mL= milliliter

Drug Facts (continued)

- Adults and children 12 years and older: 15 mL to 30 mL
- Children under 12 years: ask a doctor

Other information

- each 15 mL contains: magnesium 1000 mg
- store at room temperature. Avoid freezing. Close cap tightly after use.

Inactive ingredients anhydrous citric acid, D&C red #28, flavors, glycerin propylene glycol, purified water, sodium hypochlorite, sorbitol, sucrose

Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST)

Bayer, the Bayer Cross, Phillips' and Phillips' Milk of Magnesia design are registered trademarks of Bayer



PHILLIPS MILK OF MAGNESIA CONCENTRATED FRESH **STRAWBERRY**

phillips milk of magnesia concentrated fresh strawberry liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0280-0033

Route of Administration ORAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength MAGNESIUM HYDROXIDE (UNII: NBZ 3QY004S) (MAGNESIUM CATION - UNII: T6V3LHY838) MAGNESIUM HYDROXIDE (UNII: NBZ 3QY004S) (MAGNESIUM CATION - UNII: T6V3LHY838) MAGNESIUM HYDROXIDE (UNII: NBZ 3QY004S) (MAGNESIUM CATION - UNII: T6V3LHY838)

| Inactive Ingredients | | | | |
|--|----------|--|--|--|
| Ingredient Name | Strength | | | |
| SODIUM HYPOCHLORITE (UNII: DY38VHM5OD) | | | | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | | | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| SUCROSE (UNII: C151H8M554) | | | | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | | | | |
| D&C RED NO. 28 (UNII: 767IP0Y5NH) | | | | |
| GLYCERIN (UNII: PDC6A3C0OX) | | | | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | | | | |

| Product Characteristics | | | | |
|-------------------------|------------|--------------|--|--|
| Color | | Score | | |
| Shape | | Size | | |
| Flavor | STRAWBERRY | Imprint Code | | |
| Contains | | | | |

Item Code Package Description Marketing Start Date Marketing End Date 1 NDC:02800033-09 Combination Product 06/20/2016

| Marketing Information | | | | | |
|-----------------------|---|-------------------------|-----------------------|--|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | | |
| OTC Monograph Drug | M007 | 12/18/2014 | | | |
| | | | | | |

Revised: 11/2023 Bayer HealthCare LLC.