

# **MIGRAINE RELIEF- acetaminophen, aspirin and caffeine tablet**

## **DISCOUNT DRUG MART**

-----  
**1205-DDM-2024-0613**

### ***Drug Facts***

#### **Active ingredient(s)**

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

#### **Purpose**

Pain reliever

Pain reliever

Pain reliver aid

#### **Use**

■ treats migraine

#### **Warnings**

##### **Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

##### **Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

##### **Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)

- shock

### **Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Medication overuse headache warning**

Headaches may worsen if this product is used for 10 or more days per month.

### **Caffeine warning**

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches

- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
  - diabetes
  - gout
  - arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**■ do not use more than directed**

- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

**Other information**

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- close cap tightly after use
- read all product information before using.
- keep this box for important information

**Inactive ingredients**

carnauba wax, hydroxypropyl cellulose, hypromellose, microcrystalline cellulose, polyethylene glycol, sodium stearyl fumarate, titanium dioxide

**Questions or comments?**

1-844-705-4384

**PRINCIPAL DISPLAY PANEL**

DISCOUNT drug mart FOOD FAIR

NDC 53943-305-03

†Compare to the active ingredients in Excedrin® Migraine  
Migraine Relief

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets

Pain Reliever / Pain Reliever Aid

100 COATED CAPLETS\*\*

\*\*Capsule-Shaped Tablets

INK AND COATING FREE  
FOR LOT AND  
EXPIRATION STAMPING



06230153660

11 The product is not manufactured or distributed in the USA.  
Consumer Health Care, a division of Cardinal Health



**DO NOT USE IF IMPRINTED SEAL  
UNDER CAP IS BROKEN OR MISSING**

KEEP BOX FOR REFERENCE

**Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg  
Aspirin 250 mg (NSAID)  
Caffeine 65 mg  
Hydrocodone bitartrate (opioid) and acetaminophen (pain reliever/fever reducer)

**Use**

Relieves pain

**Warnings**

Do not use if you have had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (paracetamol) or aspirin. Do not use if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if:**

- you have never had migraines diagnosed by a health professional
- you have a medical condition that is different from your usual migraines
- you have the worst headache of your life
- you have heart and still neck
- you have been diagnosed with liver or kidney disease
- you have been diagnosed with stomach or intestinal problems, such as heartburn
- you have a history of stomach problems, heart disease, liver disease, or kidney disease
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking any of the following medicines:
  - you have asthma
  - you are taking any of the following medicines:
    - you have a history of stomach problems, such as heartburn
    - you have a history of stomach problems, heart disease, liver disease, or kidney disease
    - you have high blood pressure, heart disease, liver disease, or kidney disease

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (paracetamol) or aspirin. Do not use if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if:**

- you have never had migraines diagnosed by a health professional
- you have a medical condition that is different from your usual migraines
- you have the worst headache of your life
- you have heart and still neck
- you have been diagnosed with liver or kidney disease
- you have been diagnosed with stomach or intestinal problems, such as heartburn
- you have a history of stomach problems, heart disease, liver disease, or kidney disease
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking any of the following medicines:
  - you have asthma
  - you are taking any of the following medicines:
    - you have a history of stomach problems, such as heartburn
    - you have a history of stomach problems, heart disease, liver disease, or kidney disease
    - you have high blood pressure, heart disease, liver disease, or kidney disease

Do not use if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (paracetamol) or aspirin. Do not use if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if:**

- you have never had migraines diagnosed by a health professional
- you have a medical condition that is different from your usual migraines
- you have the worst headache of your life
- you have heart and still neck
- you have been diagnosed with liver or kidney disease
- you have been diagnosed with stomach or intestinal problems, such as heartburn
- you have a history of stomach problems, heart disease, liver disease, or kidney disease
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking any of the following medicines:
  - you have asthma
  - you are taking any of the following medicines:
    - you have a history of stomach problems, such as heartburn
    - you have a history of stomach problems, heart disease, liver disease, or kidney disease
    - you have high blood pressure, heart disease, liver disease, or kidney disease

**Drug Facts (continued)**

Medication overdose headache warning: Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of the product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking the product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (paracetamol) or aspirin. Do not use if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if:**

- you have never had migraines diagnosed by a health professional
- you have a medical condition that is different from your usual migraines
- you have the worst headache of your life
- you have heart and still neck
- you have been diagnosed with liver or kidney disease
- you have been diagnosed with stomach or intestinal problems, such as heartburn
- you have a history of stomach problems, heart disease, liver disease, or kidney disease
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking any of the following medicines:
  - you have asthma
  - you are taking any of the following medicines:
    - you have a history of stomach problems, such as heartburn
    - you have a history of stomach problems, heart disease, liver disease, or kidney disease
    - you have high blood pressure, heart disease, liver disease, or kidney disease

**Drug Facts (continued)**

**Step use and ask a doctor if:**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - heart burn
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- your migraines are not relieved or worsen after first dose
- your or unexpected symptoms occur
- if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
- Other medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- swallow table 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age, ask a doctor

**Other information**

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- store cap tightly after use
- read all product information before using
- keep this box for important information

**Inactive ingredients** carboxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide

**Questions or comments?** 1-844-705-4394

**Drug Facts (continued)**

Medication overdose headache warning: Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of the product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking the product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.

**Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer with any other drug containing acetaminophen (paracetamol) or aspirin. Do not use if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if:**

- you have never had migraines diagnosed by a health professional
- you have a medical condition that is different from your usual migraines
- you have the worst headache of your life
- you have heart and still neck
- you have been diagnosed with liver or kidney disease
- you have been diagnosed with stomach or intestinal problems, such as heartburn
- you have a history of stomach problems, heart disease, liver disease, or kidney disease
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking any of the following medicines:
  - you have asthma
  - you are taking any of the following medicines:
    - you have a history of stomach problems, such as heartburn
    - you have a history of stomach problems, heart disease, liver disease, or kidney disease
    - you have high blood pressure, heart disease, liver disease, or kidney disease

**Drug Facts (continued)**

**Step use and ask a doctor if:**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - heart burn
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- your migraines are not relieved or worsen after first dose
- your or unexpected symptoms occur
- if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
- Other medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- swallow table 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age, ask a doctor

**Other information**

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- store cap tightly after use
- read all product information before using
- keep this box for important information

**Inactive ingredients** carboxymethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyethylene glycol, sodium lauryl sulfate, titanium dioxide

**Questions or comments?** 1-844-705-4394

NDC 53943-305-03

†Compare to the active ingredients in Excedrin® Migraine

**MIGRAINE RELIEF**

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets  
Pain Reliever / Pain Reliever Aid

Actual Size

**100 COATED CAPLETS\*\***  
\*\*Capsule-Shaped Tablets

NC

NC

NC

**MIGRAINE RELIEF**

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets

**drug mart**

FOOD MARKET

SATISFACTION GUARANTEED

IF DISASSEMBLED, RETURN UNSEALED PORTION AND PACKAGING TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISASSEMBLY, NAME, ADDRESS AND EASY TO FOLLOW RETURN TO STORE DISCOUNT DRUG MART, 211 COMMERCIAL DRIVE, MEDINA, OHIO 44026

<b>MIGRAINE RELIEF</b> acetaminophen, aspirin and caffeine tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53943-305
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>SODIUM STEARYL FUMARATE</b> (UNII: 7CV7WJK4UI)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>HYDROXYPROPYL CELLULOSE (110000 WAMW)</b> (UNII: 5Y0974F5PW)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Capsule Shaped Tablet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	l1
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-305-03	1 in 1 CARTON	01/11/2024	
1		100 in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214039	01/11/2024	

**Labeler** - DISCOUNT DRUG MART (047741335)