

PAIN RELIEF PM- acetaminophen and diphenhydramine hydrochloride tablet, coated
HARRIS TEETER

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HTE-1095-2019-1001

Drug Facts

| <i>Active ingredients (in each caplet)</i> | <i>Purpose</i> |
|---|------------------------|
| Acetaminophen 500 mg | Pain reliever |
| Diphenhydramine HCl 25 mg | Nighttime sleep aid |

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of serious underlying medical illness.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

| | |
|--|--|
| adults and children 12 years and over | <ul style="list-style-type: none"> ▪ take 2 caplets at bedtime ▪ do not take more than 2 caplets of this product in 24 hours |
| children under 12 years | <ul style="list-style-type: none"> ▪ do not use |

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Acetaminophen PM

†Compare to the Active Ingredients in Tylenol® PM Extra Strength Caplets

Extra Strength

Acetaminophen & Diphenhydramine

PAIN RELIEVER / NIGHTTIME SLEEP AID

Non-Habit Forming

For Adults

50 CAPLETS

Actual Size

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Merck Consumer Healthcare, distributor of Tylenol® PM Extra Strength Caplets. F.109502ATE_R1

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 500 mg.....Pain reliever
Diphenhydramine HCl 25 mg.....Nighttime sleep aid

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ with any other product containing diphenhydramine, even one used on skin
 ■ in children under 12 years of age
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

Ask a doctor before use if you have
 ■ liver disease ■ glaucoma
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin
 ■ taking sedatives or tranquilizers

When using this product
 ■ avoid alcoholic drinks
 ■ drowsiness will occur
 ■ do not drive a motor vehicle or operate machinery

Stop use and ask a doctor if
 ■ sleeplessness persists continuously for more than 2 weeks; insomnia may be a symptom of serious underlying medical illness.
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not take more than directed (see overdose warning)
 adults and ■ take 2 caplets at bedtime
 children 12 ■ do not take more than 2 caplets
 years and ■ of this product in 24 hours
 over
 children ■ do not use
 under 12 years

Other information
 ■ store between 20-25°C (68-77°F) in a dry place
 ■ retain carton for complete product information

Inactive ingredients colloidal silicon dioxide, copovidone, croscarmellose sodium, FD&C blue #1, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
 1-844-705-4384

PROUDLY DISTRIBUTED BY:
 HARRIS TEETER, LLC, MATTHEWS, NC 28105
yourplusguarantee.
 1-800-432-6111 or harristeeter.com

NC

Acetaminophen PM



**Extra Strength
Acetaminophen & Diphenhydramine HCl**

PAIN RELIEVER / NIGHTTIME SLEEP AID

- Non-Habit Forming
- For Adults

50 CAPLETS



COATING FREE AREA

NC

NC

PAIN RELIEF PM

acetaminophen and diphenhydramine hydrochloride tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:72036-095 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 500 mg |
| DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M) | DIPHENHYDRAMINE HYDROCHLORIDE | 25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| COPOVIDONE K25-31 (UNII: D9C330MD8B) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| LACTOSE (UNII: J2B2A4N98G) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ) | |
| SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|----------|------|--------------|----------|
| Color | blue | Score | no score |
| Shape | OVAL | Size | 17mm |
| Flavor | | Imprint Code | AAA;1031 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:72036-095-02 | 1 in 1 CARTON | 10/01/2011 | |
| 1 | | 50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| 2 | NDC:72036-095-03 | 1 in 1 CARTON | 10/01/2011 | |
| 2 | | 100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 10/01/2011 | |

Labeler - HARRIS TEETER (047279351)

Revised: 10/2019

HARRIS TEETER