

AZELAIC ACID- azelaic acid gel
Glenmark Pharmaceuticals Inc., USA

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use AZELAIC ACID GEL safely and effectively. See full prescribing information for AZELAIC ACID GEL.

AZELAIC ACID Gel, for topical use

Initial U.S. Approval: 1995

INDICATIONS AND USAGE

Azelaic acid gel, 15% is indicated for topical treatment of the inflammatory papules and pustules of mild to moderate rosacea (1).

Limitations of Use

Efficacy for treatment of erythema in rosacea in the absence of papules and pustules has not been evaluated. (1)

DOSAGE AND ADMINISTRATION

- Apply a thin layer twice daily to affected area(s). (2)
- Use only very mild soaps or soapless cleansing lotion and pat dry with a soft towel before applying azelaic acid gel. (2)
- Wash hands immediately following application. (2)
- Cosmetics may be applied after the application of azelaic acid gel has dried. (2)
- Avoid use of alcoholic cleansers, tinctures and astringents, abrasives and peeling agents. (2)
- For topical use. (2)
- Not for oral, ophthalmic or intravaginal use. (2)

DOSAGE FORMS AND STRENGTHS

Gel, 15% (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

- Hypersensitivity: Hypersensitivity reactions, including cases of angioedema, eye swelling, facial swelling, dyspnea, urticaria, and adverse skin reactions, have been reported. In case of known hypersensitivity to any component of the gel, avoid the use of azelaic acid gel. If hypersensitivity develops, discontinue treatment and institute appropriate therapy. (5.1)
- Skin Reactions: Skin irritation (i.e. pruritus, burning or stinging) may occur, usually during the first few weeks of treatment. If sensitivity or severe irritation develops and persists, discontinue treatment and institute appropriate therapy. (5.2)
- Hypopigmentation: Isolated cases of hypopigmentation occurred after azelaic acid use. Monitor patients with dark complexion for early signs of hypopigmentation.(5.2)
- Eye and Mucous Membrane Irritation: Azelaic acid gel has been reported to cause irritation of the eyes. Avoid contact with the eyes and mucous membranes. (5.3)
- Exacerbation of Asthma: Consult a physician if asthma is exacerbated with azelaic acid gel use. (5.4)

ADVERSE REACTIONS

The most common adverse reactions are burning/stinging/tingling (29%), pruritus (11%), scaling/dry skin/xerosis (8%) and erythema/irritation (4%). (6)

To report SUSPECTED ADVERSE REACTIONS, contact Glenmark Pharmaceuticals Inc., USA at 1 (888) 721-7115 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 12/2021

1 INDICATIONS AND USAGE**2 DOSAGE AND ADMINISTRATION****3 DOSAGE FORMS AND STRENGTHS****4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

5.1 Hypersensitivity

5.2 Skin Reactions

5.3 Eye and Mucous Membranes Irritation

5.4 Exacerbation of Asthma

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES**16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION**1 INDICATIONS AND USAGE**

Azelaic acid gel, 15% is indicated for topical treatment of the inflammatory papules and pustules of mild to moderate rosacea.

Limitations of Use

Although some reduction of erythema which was present in patients with papules and pustules of rosacea occurred in clinical studies, efficacy for treatment of erythema in rosacea in the absence of papules and pustules has not been evaluated.

2 DOSAGE AND ADMINISTRATION

- Cleanse affected area(s) using only very mild soaps or soapless cleansing lotion and pat dry with a soft towel before application of azelaic acid gel.

- Apply and gently massage a thin layer of azelaic acid gel into the affected areas on the face twice daily (morning and evening).
- Wash hands immediately following application of azelaic acid gel.
- Cosmetics may be applied after the application of azelaic acid gel has dried.
- Reassess the diagnosis if no improvement is observed upon completing 12 weeks of therapy.
- Avoid the use of occlusive dressings or wrappings.
- Instruct patients to avoid use of alcoholic cleansers, tinctures and astringents, abrasives and peeling agents.
- For topical use.
- Not for oral, ophthalmic or intravaginal use.

3 DOSAGE FORMS AND STRENGTHS

Azelaic Acid Gel, 15% is a white to yellowish white opaque gel. Each gram of azelaic acid gel contains 0.15 gm of azelaic acid (15% w/w).

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity

Hypersensitivity reactions, including cases of angioedema, eye swelling, facial swelling, dyspnea, urticaria, and adverse skin reactions, have been reported during post marketing surveillance.

Avoid the use of azelaic acid gel in patients with known hypersensitivity to any component of the gel. If hypersensitivity develops during treatment, discontinue azelaic acid gel and institute appropriate therapy.

5.2 Skin Reactions

Skin irritation (i.e. pruritus, burning or stinging) may occur during use of azelaic acid gel, usually during the first few weeks of treatment. If sensitivity or severe irritation develops and persists, discontinue treatment and institute appropriate therapy.

There have been isolated reports of hypopigmentation after use of azelaic acid. Since azelaic acid has not been well studied in patients with dark complexion, monitor these patients for early signs of hypopigmentation

5.3 Eye and Mucous Membranes Irritation

Azelaic acid gel has been reported to cause irritation of the eyes. Avoid contact with the eyes, mouth and other mucous membranes. If azelaic acid gel comes in contact with the eyes, wash the eyes with large amounts of water and consult a physician if eye irritation persists [*see Adverse Reactions (6.2)*].

5.4 Exacerbation of Asthma

Worsening of asthma has been reported in patients using azelaic acid formulations including azelaic acid gel. Consult a physician if asthma is exacerbated with use of azelaic acid gel.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In two vehicle-controlled and one active-controlled U.S. clinical trials, treatment safety was monitored in 788 subjects who used twice-daily azelaic acid gel for 12 weeks (N=333) or 15 weeks (N=124), or the gel vehicle (N=331) for 12 weeks. In all three trials, the most common treatment-related adverse events were: burning/stinging/tingling (29%), pruritus (11%), scaling/dry skin/xerosis (8%) and erythema/irritation (4%). In the active-controlled trial, overall adverse reactions (including burning, stinging/tingling, dryness/tightness/scaling, itching, and erythema/irritation/redness) were 19.4% (24/124) for azelaic acid gel compared to 7.1% (9/127) for the active comparator gel at 15 weeks.

Table 1: Adverse Events Occurring in ≥1% of Subjects in the Rosacea Trials by Treatment Group and Maximum Intensity*

	Azelaic acid gel, 15% N=457 (100%)			Vehicle N=331 (100%)		
	Mild N=99 (22%)	Moderate N=61 (13%)	Severe N=27 (6%)	Mild N=46 (14%)	Moderate N=30 (9%)	Severe N=5 (2%)
Burning/stinging/tingling	71 (16%)	42 (9%)	17 (4%)	8 (2%)	6 (2%)	2 (1%)
Pruritus	29 (6%)	18 (4%)	5 (1%)	9 (3%)	6 (2%)	0 (0%)
Scaling/dry skin/ xerosis	21 (5%)	10 (2%)	5 (1%)	31 (9%)	14 (4%)	1 (<1%)
Erythema/ irritation	6 (1%)	7 (2%)	2 (<1%)	8 (2%)	4 (1%)	2 (1%)
Contact dermatitis	2 (<1%)	3 (1%)	0 (0%)	1 (<1%)	0 (0%)	0 (0%)
Edema	3 (1%)	2 (<1%)	0 (0%)	3 (1%)	0 (0%)	0 (0%)
Acne	3 (1%)	1 (<1%)	0 (0%)	1 (<1%)	0 (0%)	0 (0%)

*Subjects may have >1 cutaneous adverse event; thus, the sum of the frequencies of preferred terms may exceed the number of subjects with at least 1 cutaneous adverse event.

In patients using azelaic acid formulations, the following adverse events have been reported: worsening of asthma, vitiligo, depigmentation, small depigmented spots, hypertrichosis, reddening (signs of keratosis pilaris) and exacerbation of recurrent herpes labialis.

Local Tolerability Studies

Azelaic acid gel and its vehicle caused irritant reactions at the application site in human dermal safety studies. Azelaic acid gel caused significantly more irritation than its vehicle in a cumulative irritation study. Some improvement in irritation was demonstrated over the course of the clinical trials, but this improvement might be attributed to subject dropouts. No phototoxicity or photoallergenicity were reported in human dermal safety studies.

6.2 Postmarketing Experience

The following adverse reactions have been identified post approval of azelaic acid gel. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate the frequency or establish a causal relationship to drug exposure:

Eyes: iridocyclitis upon accidental exposure of the eyes to azelaic acid gel.

Hypersensitivity: angioedema, eye swelling, facial swelling, urticaria.

Respiratory: worsening of asthma, dyspnea, wheezing.

Skin reactions: application site rash.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Azelaic acid is minimally absorbed systemically following topical route of administration, and maternal use is not expected to result in fetal exposure to the drug [*see Clinical Pharmacology (12.3)*].

In animal reproduction studies, embryofetal toxicity was noted when azelaic acid was administered orally during the period of organogenesis at doses 162, 19, and 65 times the maximum recommended human dose (MRHD) in rats, rabbits, and monkeys, respectively. Maternal toxicity was noted at these doses but no malformations were observed in these embryofetal developmental studies (*see Data*).

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Dermal embryofetal developmental toxicology studies have not been performed with azelaic acid, 15% gel. Oral embryofetal developmental studies were conducted with azelaic acid in rats, rabbits, and cynomolgus monkeys. Azelaic acid was administered during the period of organogenesis in all three animal species. Embryotoxicity was observed in rats, rabbits, and monkeys at oral doses of azelaic acid that generated some maternal toxicity. Embryotoxicity was observed in rats given 2500 mg/kg/day [162 times the MRHD based on body surface area (BSA) comparison], rabbits given 150 or

500 mg/kg/day (19 or 65 times the MRHD based on BSA comparison) and cynomolgus monkeys given 500 mg/kg/day (65 times the MRHD based on BSA comparison) azelaic acid. No malformations were observed in the oral embryofetal developmental studies conducted in rats, rabbits and cynomolgus monkeys.

An oral peri- and post-natal developmental study was conducted in rats. Azelaic acid was administered from gestational day 15 through day 21 postpartum up to a dose level of 2500 mg/kg/day. Embryotoxicity was observed in rats at an oral dose of 2500 mg/kg/day (162 times the MRHD based on BSA comparison) that generated some maternal toxicity. In addition, slight disturbances in the post-natal development of fetuses was noted in rats at oral doses that generated some maternal toxicity (500 and 2500 mg/kg/day; 32 and 162 times the MRHD based on BSA comparison). No effects on sexual maturation of the fetuses were noted in this study.

8.2 Lactation

Risk Summary

Azelaic acid is naturally present in human milk. When used as prescribed, azelaic acid is unlikely to be absorbed through the skin in clinically relevant amounts to cause a change in azelaic acid concentration in milk or milk production; therefore, breastfeeding is not expected to result in exposure of the infant to azelaic acid gel. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for azelaic acid gel and any potential adverse effects on the breastfed child from azelaic acid gel or from the underlying maternal condition.

8.4 Pediatric Use


The safety and effectiveness of azelaic acid gel have not been established in pediatric patients.

8.5 Geriatric Use

Clinical studies of azelaic acid gel did not include sufficient numbers of subjects aged 65 and over to determine if they respond differently than younger subjects.

11 DESCRIPTION

Azelaic Acid Gel, 15%, is a white to yellowish white opaque gel which contains azelaic acid, a naturally-occurring saturated dicarboxylic acid. It is for topical use. Chemically, azelaic acid is 1,7-heptanedicarboxylic acid. The molecular formula for azelaic acid is $C_9H_{16}O_4$. It has the following structure:



Azelaic acid has a molecular weight of 188.22 g/mol. It is a white to off-white solid that is insoluble in water, toluene and heptane, but soluble in acetone, methanol, isopropyl alcohol and isopropyl acetate.

Azelaic Acid Gel, 15% is a white to yellowish white opaque gel for topical use; each gram contains 0.15 gm azelaic acid (15% w/w) as the active ingredient in an aqueous gel base containing benzoic acid (as a preservative), carbomer homopolymer, disodium edetate, isopropyl myristate, polysorbate 80, propylene glycol, purified water and sodium hydroxide to adjust pH.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism(s) by which azelaic acid interferes with the pathogenic events in rosacea are unknown.

12.2 Pharmacodynamics

The pharmacodynamics of azelaic acid in association with the treatment of rosacea are unknown.

12.3 Pharmacokinetics

The percutaneous absorption of azelaic acid after topical application of azelaic acid gel could not be reliably determined. Mean plasma azelaic acid concentrations in rosacea subjects treated with azelaic acid gel twice daily for at least 8 weeks are in the range of 42 to 63.1 ng/mL. These values are within the maximum concentration range of 24 to 90.5 ng/mL observed in rosacea subjects treated with vehicle only. This indicates that azelaic acid gel does not increase plasma azelaic acid concentration beyond the range derived from nutrition and endogenous metabolism.

In vitro and human data suggest negligible cutaneous metabolism of ³H-azelaic acid after topical application of 20% azelaic acid cream. Azelaic acid is mainly excreted unchanged in the urine, but undergoes some β -oxidation to shorter chain dicarboxylic acids.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 2-year dermal mouse carcinogenicity study, azelaic acid pre-foam emulsion was administered twice daily to CD-1 mice at topical doses of 5%, 15%, and 30% (500, 1500, and 3000 mg/kg/day azelaic acid). No drug-related tumors were noted at concentrations up to 30% azelaic acid (396 times the MRHD based on AUC comparison).

Azelaic acid was not mutagenic or clastogenic in a battery of *in vitro* [Ames assay, HGPRT in V79 cells (Chinese hamster lung cells), and chromosomal aberration assay in human lymphocytes] and *in vivo* (dominant lethal assay in mice and mouse micronucleus assay) genotoxicity tests.

Oral administration of azelaic acid at dose levels up to 2500 mg/kg/day (162 times the MRHD based on BSA comparison) did not affect fertility or reproductive performance in male or female rats.

14 CLINICAL STUDIES

Azelaic acid gel was evaluated for the treatment of mild to moderate papulopustular rosacea in two multicenter, randomized, double-blind, vehicle-controlled, 12-week clinical trials having identical protocols and involving a total of 664 (active: 333; vehicle: 331) subjects aged 21 to 86 years (mean age = 49). Overall, 92.5% of subjects were Caucasian and 73% of subjects were female. Enrolled subjects had mild to moderate rosacea with a mean lesion count of 18 (range 8 to 60) inflammatory papules and pustules. The following subjects were excluded: a) those without papules and pustules; b) those with nodules, rhinophyma, or ocular involvement and c) those with a history of hypersensitivity to propylene glycol or to any other ingredients of the study drug. Azelaic acid gel or its vehicle were to be applied twice daily for 12 weeks; no other topical or systemic medication affecting the course of rosacea and/or evaluability was to be used during the studies. Subjects were instructed to avoid spicy foods, thermally hot food/drink and alcoholic beverages during the study. Subjects were also instructed to use only very mild soaps or soapless cleansing lotion for facial cleansing.

The primary efficacy endpoints included both 1) change from baseline in inflammatory lesion counts as well as 2) success defined as a score of “clear” or “minimal” with at least a 2-step reduction from baseline on the Investigator’s Global Assessment (IGA), defined as follows below:

CLEAR:

No papules and/or pustules; no or residual erythema; no or mild to moderate telangiectasia

MINIMAL:

Rare papules and/or pustules; residual to mild erythema; mild to moderate telangiectasia

MILD:

Few papules and/or pustules; mild erythema; mild to moderate telangiectasia

MILD TO MODERATE:

Distinct number of papules and/or pustules; mild to moderate erythema; mild to moderate telangiectasia

MODERATE:

Pronounced number of papules and/or pustules; moderate erythema; mild to moderate telangiectasia

MODERATE TO SEVERE:

Many papules and/or pustules, occasionally with large inflamed lesions; moderate erythema; moderate degree of telangiectasia

SEVERE:

Numerous papules and/or pustules, occasionally with confluent areas of inflamed lesions; moderate or severe erythema; moderate or severe telangiectasia

Primary efficacy assessment was based on the “intent-to-treat” (ITT) population with the “last observation carried forward” (LOCF).

Both trials demonstrated a statistically significant difference in favor of azelaic acid gel

over its vehicle in both reducing the number of inflammatory papules and pustules associated with rosacea (Table 2) as well as demonstrating success on the IGA in the ITT-LOCF population at the end of treatment.

Table 2: Inflammatory Papules and Pustules (ITT population)*

	Study One AZELAIC ACID GEL, 15% N=164	Study One VEHICLE N=165	Study Two Azelaic acid gel, 15% N=167	Study Two VEHICLE N=166
Mean Lesion Count Baseline	17.5	17.6	17.9	18.5
End of Treatment ¹	6.8	10.5	9	12.1
Mean Percent Reduction End of Treatment ¹	57.9%	39.9%	50.0%	38.2%

*ITT population with last observation carried forward (LOCF)

Although some reduction of erythema which was present in subjects with papules and pustules of rosacea occurred in clinical trials, efficacy for treatment of erythema in rosacea in the absence of papules and pustules has not been evaluated.

Azelaic acid gel was superior to the vehicle with regard to success based on the IGA of rosacea on a 7-point static score at the end of treatment (ITT population; Table 3).

Table 3: Investigator's Global Assessment at the End of Treatment*

	Study One Azelaic acid gel, 15% N=164	Study One VEHICLE N=165	Study Two Azelaic acid gel, 15% N=167	Study Two VEHICLE N=166
Clear, Minimal or Mild at End of Treatment (% of Subjects)	61%	40%	61%	48%

*ITT population with last observation carried forward (LOCF)

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

Azelaic Acid Gel, 15%, a white to yellowish white opaque gel, is supplied in 50 g tubes (NDC 68462-626-52).

Storage and Handling

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Inform patients using azelaic acid gel of the following

Administration instructions

- For topical use only.
- Before applying azelaic acid gel, cleanse affected area(s) with a very mild soap or a soapless cleansing lotion and pat dry with a soft towel.
- Wash hands immediately following application of azelaic acid gel.
- Cosmetics may be applied after the application of azelaic acid gel has dried.
- Avoid the use of occlusive dressings or wrappings.
- Avoid use of alcoholic cleansers, tinctures and astringents, abrasives and peeling agents [see *Dosage and Administration (2)*].

Hypersensitivity

- If allergic reactions occur, discontinue use and consult their healthcare provider [see *Warnings and Precautions (5.1)*].

Skin Irritation

- Skin irritation (e.g., pruritus, burning, or stinging) may occur during use of azelaic acid gel, usually during the first few weeks of treatment. If irritation is excessive or persists, or allergic reactions occur, discontinue use and consult your physician [see *Warnings and Precautions (5.2)*].

Hypopigmentation

- Advise patients to report abnormal changes in skin color to their healthcare provider [see *Warnings and Precautions (5.2)*].

Eye and Mucous Membranes Irritation

- Avoid contact with the eyes, mouth and other mucous membranes. If azelaic acid gel comes in contact with the eyes, wash the eyes with large amounts of water and consult their healthcare provider if eye irritation persists [see *Warnings and Precautions (5.3)*].

Exacerbation of Asthma

- Advise patients to report any worsening of asthma to their healthcare provider [see *Warnings and Precautions (5.4)*].

Manufactured by:

Glenmark Pharmaceuticals Limited
Colvale-Bardez, Goa 403513, India

Manufactured for:



Glenmark Pharmaceuticals Inc., USA

Mahwah, NJ 07430

Questions? 1 (888) 721-7115
www.glenmarkpharma-us.com

December 2021

Package/Label Display Panel

NDC 68462-626-52

Azelaic Acid Gel, 15 %

For Topical Use Only

NDC 68462-626-52

Azelaic Acid Gel, 15%

For Topical Use Only

Not for oral, ophthalmic or intravaginal use



Rx only

50 grams

Each gram of Azelaic Acid Gel, 15% contains: 0.15 gm azelaic acid (15% w/w) as the active ingredient in an aqueous gel base containing benzoic acid (as a preservative), carbomer homopolymer, disodium edetate, isopropyl myristate, polysorbate 80, propylene glycol, purified water and sodium hydroxide to adjust pH.

Dosage: Apply and gently massage a thin layer of Azelaic Acid Gel, 15% into the affected areas on the face twice daily (morning and evening).

See package insert for full prescribing information.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

To Open: Use pointed end on cap to puncture seal.

For lot number and expiry date see crimp.

Manufactured by:
**Glenmark
Pharmaceuticals Limited**
Colvale-Bardez, Goa 403513, India
GO/DRUGS/648

Manufactured for: **Glenmark
Pharmaceuticals
Inc., USA**
Mahwah, NJ 07430
01/21 PE580420121-1



Questions? 1 (888) 721-7115
www.glenmarkpharma-us.com

AZELAIC ACID

azelaic acid gel

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68462-626
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AZELAIC ACID (UNII: F2VW3D43YT) (AZELAIC ACID - UNII:F2VW3D43YT)	AZELAIC ACID	0.15 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68462-626-52	1 in 1 CARTON	11/19/2018	
1		50 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204637	11/19/2018	

Labeler - Glenmark Pharmaceuticals Inc., USA (130597813)

Establishment

Name	Address	ID/FEI	Business Operations
Glenmark Pharmaceuticals Limited		677318665	ANALYSIS(68462-626) , MANUFACTURE(68462-626)

