

CURANEX DM- dextromethorphan hydrobromide 20 mg guaifenesin 400 mg tablet
Wittman Pharma, Inc.

CuraNEX DM

- take with a full glass of water
- adults and children 12 years and over: 1 tablet every 4 hours.
- Do not take more than 6 tablets in 24 hours.
- children under 12 years: do not use

Other information

Do Not Use ~ you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use If you have

- cough accompanied by too much phlegm (mucus)
- persistent or chronic cough such as occurs~~ with smoking, asthma, chronic bronchitis, or emphysema

■ When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache.

These could be signs of a serious condition.

■ if pregnant or breast-feeding, ask a health professional before use.

Magnesium Stearate, Microcrystalline Cellulose,

Sodium Starch Glycolate

Cough Suppressant and Expectorant

- helps: loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves: ■ cough due to minor throat and bronchial irritation associated with the common cold
- the intensity of coughing ■ the impulse to cough to help you get to sleep

Keep out of reach of children.

- In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Purpose

(In each Immediate-re/ease tablet)

Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

Drug Facts

Active ingredients Purpose

(In each Immediate-re/ease tablet)

Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant

Use

- helps: loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves: ■ cough due to minor throat and bronchial irritation associated with the common cold
- the intensity of coughing ■ the impulse to cough to help you get to sleep

Warnings

Do Not Use ~ you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use If you have

- cough accompanied by too much phlegm (mucus)
- persistent or chronic cough such as occurs~~ with smoking, asthma, chronic bronchitis, or emphysema

(Panel 2)

Warnings (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache.

These could be signs of a serious condition.

- if pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

- In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- take with a full glass of water
- adults and children 12 years and over: 1 tablet every 4 hours.
- Do not take more than 6 tablets in 24 hours.

■ children under 12 years: do not use

Other information

■ store between 15°-30°C (59°-86°F)

Inactive ingredients

Magnesium Stearate, Microcrystalline Cellulose,

Sodium Starch Glycolate

Questions or Comments?

call 1-352-549-9917, Mon-Fri: 9:00AM - 5:00 PM



TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	Purpose
Active ingredients (in each immediate-release tablet) Dextromethorphan HBr 20 mg Cough suppressant Guaifenesin 400 mg Expectorant	
Use	
<ul style="list-style-type: none"> ■ helps: loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves: ■ cough due to minor throat and bronchial irritation associated with the common cold ■ the intensity of coughing ■ the impulse to cough to help you get to sleep 	
Warnings	
<p>Do Not Use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ cough accompanied by too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema 	Peel Here

Drug Facts (continued)
Warnings (continued)
<p>When using this product do not exceed recommended dosage.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. <p>These could be signs of a serious condition.</p> <ul style="list-style-type: none"> ■ If pregnant or breast-feeding, ask a health professional before use. <p>Keep out of reach of children.</p> <ul style="list-style-type: none"> ■ In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
Directions
<ul style="list-style-type: none"> ■ take with a full glass of water ■ adults and children 12 years and over: 1 tablet every 4 hours. ■ Do not take more than 6 tablets in 24 hours. ■ children under 12 years: do not use
Other information
<ul style="list-style-type: none"> ■ store between 15°-30°C (59°-86°F)
Inactive ingredients
Magnesium Stearate, Microcrystalline Cellulose, Sodium Starch Glycolate
Questions or Comments?
<p>Call 1-352-549-9917, Mon-Fri: 9:00 AM - 5:00 PM</p>

CURANEX DM

dextromethorphan hydrobromide 20 mg guaifenesin 400 mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83335-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg
DEXTROMETHORPHAN (UNII: 7355X3ROTS) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	13mm
Flavor		Imprint Code	DG
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83335-004-01	1 in 1 BOTTLE; Type 0: Not a Combination Product	11/01/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/01/2023	

Labeler - Wittman Pharma, Inc. (830980947)

Establishment

Name	Address	ID/FEI	Business Operations
Wittman Pharma, Inc.		830980947	manufacture(83335-004)

Revised: 4/2024

Wittman Pharma, Inc.