

**NAPROXEN SODIUM- naproxen sodium tablet**  
**TYA Pharmaceuticals**

-----  
**Perrigo Naproxen Sodium Tablets 220 mg Drug Facts**

**Active ingredient (in each tablet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - it feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older:

- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period

Children under 12 years:

- ask a doctor

**Other information**

- each tablet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

1-800-719-9260

**NAPROXEN SODIUM TABLET**

# NAPROXEN SODIUM 220MG TABLET(100CTBTL)

45802-490-78

LOT: XXXX

EXP: XX-XX-XX

MFR: PERRIGO

REPACKAGED BY: T.Y.A. PHARMACEUTICALS  
2930 CRESCENT DR.  
TALLAHASSEE, FL 32301  
(850) 385-0228



DOSE: SEE PACKAGE INSERT

## NAPROXEN SODIUM

naproxen sodium tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:64725-0490(NDC:45802-490)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	BLUE (Light Blue)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L490
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:64725-0490-1	100 in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	04/11/2014	

**Labeler** - TYA Pharmaceuticals (938389038)

**Registrant** - TYA Pharmaceuticals (938389038)

## Establishment

Name	Address	ID/FEI	Business Operations
TYA Pharmaceuticals		938389038	RELABEL(64725-0490) , REPACK(64725-0490)

Revised: 4/2014

TYA Pharmaceuticals