HALODINE ORAL ANTISEPTIC- povidone-iodine spray Halodine Llc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Halodine Oral Antiseptic Spray

Drug Facts

Active Ingredient

Povidone-Iodine USP, 1.25% (0.125% Available Iodine)

Purpose

Antiseptic

Uses • Oral Antiseptic • First aid to help prevent infection in minor oral irritation caused by dental procedures

Warnings For external use only.

Do not use • If allergic to iodine • In the eyes • On children less than 3 years old • For more than 7 days unless directed by a dentist or doctor **Stop use and promptly ask your dentist or doctor** if sore mouth symptoms do not improve within 7 days, if irritation, pain, or redness persists or worsens, or if swelling, rash, or fever develops.

Keep out of reach of children. If large amounts swallowed, get medical help or contact a Poison Control Center right away.

□

Directions For oral antisepsis in adults and children 3 years of age and older:

1. Press down pump completely and spray twice into mouth. 2. Swish vigorously in mouth for 15 seconds. 3. Slightly lift chin and tilt head back to gargle. Spit out excess. Children under 3 years of age: Consult a dentist or doctor

Other Information • Not made with natural rubber latex • Store at room temperature; avoid excessive heat

Inactive Ingredients hydroxyethylcellulose, purified water

Questions or comments? contact@halodine.com

Povidone-Iodine USP, 1.25%

[Helps Reduce Risk of Infection
4-Hour Formula

halodine.com

©PATENT PENDING

Manufactured in the U.S.A.

for Halodine LLC 111 NE 1st St,

Ste 806, Miami, FL 33132

Packaging

Halōdine

Oral Antiseptic Spray Povidone-Iodine USP, 1.25%

2fl oz (60ml)

Helps Reduce Risk of Infection 4-Hour Formula

NDC: 78371-401-01

Oral Antiseptic Spray Povidone-Iodine USP, 1.25%

2fl oz (60ml)

Helps Reduce Risk of Infection 4-Hour Formula

halodine.com PATENT PENDING

> LOT EXP

Oral Antiseptic Spray Povidone-lodine USP, 1.25%

2fl oz (60ml)

Helps Reduce Risk of Infection 4-Hour Formula

Manufactured in the U.S.A. for Halodine LLC 111 NE 1st St, Ste 806, Miami, FL 33132

Drug Facts

Active Ingredient

Purpose

Povidone-lodine USP, 1.25% (0.125% Available lodine)

Uses ■ Oral Antiseptic ■ First aid

to help prevent infection in minor oral irritation caused by dental procedures

Warnings For external use only.

Do not use ■ If allergic to iodine ■ In the eyes ■ On children less than 3 years old ■ For more than 7 days unless directed by a dentist or doctor

Stop use and promptly ask your dentist or doctor if sore mouth symptoms do not improve within 7 days.

Keep out of reach of children. If large amounts swallowed, get medical help or contact a Poison Control Center right away.

Directions

For oral antisepsis in adults and children 3 years of age and older:

- 1. Press down pump completely and spray twice into mouth. 2. Swish vigorously in mouth for 15 seconds.
- 3. Slightly lift chin and tilt head back to gargle. Spit out excess.

Children under 3 years of age: Consult a dentist or doctor

Other Information

Not made

with natural rubber latex

Store at room temperature; avoid excessive heat

Inactive Ingredients

hydraxyethylcellulose, purified water

Questions or comments? contact@halodine.com



Oral Antiseptic Spray Povidone-Iodine USP, 1.25%

2fl oz (60ml)

Helps Reduce Risk of Infection 4-Hour Formula

halodine.com

PATENT PENDING

Manufactured in the U.S.A. for Halodine LLC 111 NE 1st St. Ste 806, Miami, FL 33132

NDC: 78371-401-01

Active Ingredient

Purpose

Povidone-Iodine USP, 1.25%Antiseptic

(0.125% Available Iodine)

Uses ■ Oral Antiseptic ■ First aid to help prevent infection in minor oral irritation caused by dental procedures Warnings For external use only.

Do not use ■ If allergic to iodine ■ In the eyes ■ On children less than 3 years old ■ For more than 7 days unless directed by a dentist or doctor Stop use and promptly ask your dentist or doctor if sore mouth symptoms do not improve within 7 days, if irritation, pain, or redness persists or worsens, or if swelling, rash, or fever develops. Keep out of reach of children. If large amounts swallowed, get medical help or contact a Poison Control Center right away.

Directions For oral antisepsis in adults and children 3 years of age and older:

- 1. Press down pump completely and spray twice into mouth.
- 2. Swish vigorously in mouth for 15 seconds. 3. Slightly lift chin and tilt head back to gargle. Spit out excess. Children under 3 years of age: Consult a dentist or doctor

Other Information ■ Not made with natural rubber latex Store at room temperature; avoid excessive heat

Inactive Ingredients hydroxyethylcellulose, purified water Questions or comments? contact@halodine.com

HALODINE ORAL ANTISEPTIC

povidone-iodine spray

Product Information

NDC:78371-401 Product Type HUMAN OTC DRUG Item Code (Source)

ORAL Route of Administration

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PO VIDO NE-IO DINE (UNII: 85H0 HZU99 M) (IO DINE - UNII:9679 TC07X4)	IODINE	1.25 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
HYDRO XYETHYL CELLULO SE, UNSPECIFIED (UNII: T4V6TWG28D)	

WATER (UNII: 059QF0KO0R)

Packaging

Marketing Start Marketing End Item Code **Package Description** Date Date

	60 mL in 1 BOTTLE, SPRAY; Type 0: Not a Com Product	10/0 1/20 20			
Marketing Information					
Marketing Catego	ory Application Number or Monograph	Citation Marketing Start Date	Marketing End Date		
OTC monograph not f	final part356	10/01/2020			

Labeler - Halodine Llc (117526113)

Revised: 10/2020 Halodine Llc