

VIMIZIM- elosulfase alfa injection, solution, concentrate
BioMarin Pharmaceutical Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VIMIZIM safely and effectively. See full prescribing information for VIMIZIM.

VIMIZIM (elosulfase alfa) injection, for intravenous use

Initial U.S. Approval: 2014

WARNING: RISK OF ANAPHYLAXIS

See full prescribing information for complete boxed warning.

Life-threatening anaphylactic reactions have occurred in some patients during Vimizim infusions. Anaphylaxis, presenting as cough, erythema, throat tightness, urticaria, flushing, cyanosis, hypotension, rash, dyspnea, chest discomfort, and gastrointestinal symptoms in conjunction with urticaria, have been reported to occur during infusions, regardless of duration of the course of treatment. Closely observe patients during and after Vimizim administration and be prepared to manage anaphylaxis. Inform patients of the signs and symptoms of anaphylaxis and have them seek immediate medical care should symptoms occur. Patients with acute respiratory illness may be at risk of serious acute exacerbation of their respiratory compromise due to hypersensitivity reactions, and require additional monitoring (5.1, 5.2, 6).

----- **INDICATIONS AND USAGE** -----

Vimizim is a hydrolytic lysosomal glycosaminoglycan (GAG)-specific enzyme indicated for patients with Mucopolysaccharidosis type IVA (MPS IVA; Morquio A syndrome) (1).

----- **DOSAGE AND ADMINISTRATION** -----

2 mg per kg body weight administered once every week as an intravenous infusion over a minimum of 3.5 to 4.5 hours, based on infusion volume (2.1, 2.3).

----- **DOSAGE FORMS AND STRENGTHS** -----

Injection: 5 mg/5 mL (1 mg/mL) in single-use vials (3).

----- **CONTRAINDICATIONS** -----

None (4)

----- **WARNINGS AND PRECAUTIONS** -----

· Anaphylaxis and Hypersensitivity Reactions: Life-threatening anaphylaxis and hypersensitivity reactions have been observed in some patients during treatment with Vimizim. If anaphylaxis or severe hypersensitivity reactions occur, immediately stop the infusion and initiate appropriate medical treatment. Pre-treatment with antihistamines with or without antipyretics is recommended prior to the start of infusion (5.1).

· Risk of Acute Respiratory Complications: Patients with acute febrile or respiratory illness may be at higher risk of life-threatening complications from hypersensitivity reactions. Careful consideration should be given to the patient's clinical status prior to administration of Vimizim and consider delaying the Vimizim infusion (5.2).

----- **ADVERSE REACTIONS** -----

The most common adverse reactions ($\geq 10\%$ in Vimizim patients and occurring at a higher incidence than placebo-treated patients) were pyrexia, vomiting, headache, nausea, abdominal pain, chills, and fatigue (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact BioMarin at 1-866-906-6100 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **USE IN SPECIFIC POPULATIONS** -----

Pediatric Use: The safety and effectiveness of Vimizim have not been established in pediatric patients less than 5 years of age (8.4).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 2/2014

1 INDICATIONS AND USAGE**2 DOSAGE AND ADMINISTRATION**

- 2.1 Recommended Dose
- 2.2 Preparation Instructions
- 2.3 Administration Instructions
- 2.4 Storage and Stability

3 DOSAGE FORMS AND STRENGTHS**4 CONTRAINDICATIONS****5 WARNINGS AND PRECAUTIONS**

- 5.1 Anaphylaxis and Hypersensitivity Reactions
- 5.2 Risk of Acute Respiratory Complications
- 5.3 Spinal or Cervical Cord Compression

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Immunogenicity

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE**11 DESCRIPTION****12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES**16 HOW SUPPLIED/STORAGE AND HANDLING****17 PATIENT COUNSELING INFORMATION**

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION**WARNING: RISK OF ANAPHYLAXIS**

Life-threatening anaphylactic reactions have occurred in some patients during Vimizim infusions. Anaphylaxis, presenting as cough, erythema, throat tightness, urticaria, flushing, cyanosis, hypotension, rash, dyspnea, chest discomfort, and gastrointestinal symptoms (e.g., nausea, abdominal pain, retching, and vomiting) in conjunction with urticaria, have been reported to occur during Vimizim infusions, regardless of duration of the course of treatment. Closely observe patients during and after Vimizim administration and be prepared to manage anaphylaxis. Inform patients of the signs and symptoms of anaphylaxis and have them seek immediate medical care should symptoms occur. Patients with acute respiratory illness may be at risk of serious acute exacerbation of their respiratory compromise due to hypersensitivity reactions, and require additional monitoring [see *Warnings and Precautions (5.1, 5.2) and Adverse Reactions (6)*].

1 INDICATIONS AND USAGE

Vimizim (elosulfase alfa) is indicated for patients with Mucopolysaccharidosis type IVA (MPS IVA; Morquio A syndrome).

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dose

The recommended dose is 2 mg per kg given intravenously over a minimum range of 3.5 to 4.5 hours, based on infusion volume, once every week. Pre-treatment with antihistamines with or without antipyretics is recommended 30 to 60 minutes prior to the start of the infusion [see *Warnings and Precautions (5.1)*].

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

2.2 Preparation Instructions

Important Information: This product should be prepared and administered under the supervision of a healthcare professional with the ability to manage medical emergencies.

Determine the number of vials to be diluted based on the individual patient's weight and the recommended dose of 2 mg/kg.

Dilute the calculated dose to a final volume of 100 mL or 250 mL using 0.9% Sodium Chloride Injection, USP.

The final volume is based on the patient's weight as follows:

- For patients who weigh less than 25 kg, the final volume should be 100 mL;
- For patients who weigh 25 kg or more, the final volume should be 250 mL.

The solution should be clear to slightly opalescent and colorless to pale yellow when diluted. Do not use if the solution is discolored or if there is particulate matter in the solution. Note that a diluted solution with slight flocculation (e.g., thin translucent fibers) is acceptable for administration.

Avoid agitation during preparation. Gently rotate the bag to ensure proper distribution. Do not shake the solution.

2.3 Administration Instructions

Administer the diluted solution to patients using a low-protein binding infusion set equipped with a low-protein binding 0.2 micrometer (μm) in-line filter.

Note: The safety and effectiveness of Vimizim have not been established in pediatric patients less than 5 years of age [see Use in Specific Populations (8.4)].

For patients who weigh less than 25 kg: initial infusion rate should be 3 mL per hour for the first 15 minutes and, if tolerated, increased to 6 mL per hour for the next 15 minutes. If this rate is tolerated, then the rate may be increased every 15 minutes in 6 mL per hour increments, not to exceed 36 mL per hour. The total volume of the infusion should be delivered over a minimum of 3.5 hours.

For patients who weigh 25 kg or more: initial infusion rate should be 6 mL per hour for the first 15 minutes and, if tolerated, the infusion rate may be increased to 12 mL per hour for the next 15 minutes. If this rate is tolerated, then the rate may be increased every 15 minutes in 12 mL per hour increments, not to exceed 72 mL per hour. The total volume of the infusion should be delivered over a minimum of 4.5 hours.

The infusion rate may be slowed, temporarily stopped, or discontinued for that visit in the event of hypersensitivity reactions [see *Warnings and Precautions (5.1)*]. Do not infuse with other products in the

infusion tubing. Compatibility with other products has not been evaluated.

2.4 Storage and Stability

Vimizim does not contain preservatives; therefore the product should be used immediately after dilution. If immediate use is not possible, the diluted product may be stored for up to 24 hours at 2°C to 8°C (36°F to 46°F) followed by up to 24 hours at 23°C to 27°C (73°F to 81°F). Administration of Vimizim should be completed within 48 hours from the time of dilution. Vials are for single-use only. Discard any unused product. Do not freeze or shake. Protect from light.

3 DOSAGE FORMS AND STRENGTHS

Injection: 5 mg/5 mL (1 mg/mL) in single-use vials.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Anaphylaxis and Hypersensitivity Reactions

Anaphylaxis and hypersensitivity reactions have been reported in patients treated with Vimizim. In premarketing clinical trials, 18 of 235 (7.7%) patients treated with Vimizim experienced signs and symptoms consistent with anaphylaxis. These 18 patients experienced 26 anaphylactic reactions during infusion with signs and symptoms including cough, erythema, throat tightness, urticaria, flushing, cyanosis, hypotension, rash, dyspnea, chest discomfort, and gastrointestinal symptoms (e.g., nausea, abdominal pain, retching, and vomiting) in conjunction with urticaria. These cases of anaphylaxis occurred as early as 30 minutes from the start of infusion and up to three hours after infusion. Anaphylaxis occurred as late into treatment as the 47th infusion.

In clinical trials with Vimizim, 44 of 235 (18.7%) patients experienced hypersensitivity reactions, including anaphylaxis. Hypersensitivity reactions have occurred as early as 30 minutes from the start of infusion but as late as six days after infusion. Frequent symptoms of hypersensitivity reactions (occurring in more than 2 patients) included anaphylactic reactions, urticaria, peripheral edema, cough, dyspnea, and flushing.

Due to the potential for anaphylaxis, appropriate medical support should be readily available when Vimizim is administered. Observe patients closely for an appropriate period of time after administration of Vimizim, taking into account the time to onset of anaphylaxis seen in premarketing clinical trials. Inform patients of the signs and symptoms of anaphylaxis, and instruct them to seek immediate medical care should signs and symptoms occur.

Because of the potential for hypersensitivity reactions, administer antihistamines with or without antipyretics prior to infusion. Management of hypersensitivity reactions should be based on the severity of the reaction and include slowing or temporary interruption of the infusion and/or administration of additional antihistamines, antipyretics, and/or corticosteroids for mild reactions. However, if severe hypersensitivity reactions occur, immediately stop the infusion of Vimizim and initiate appropriate treatment.

Consider the risks and benefits of re-administering Vimizim following a severe reaction.

5.2 Risk of Acute Respiratory Complications

Patients with acute febrile or respiratory illness at the time of Vimizim infusion may be at higher risk of life-threatening complications from hypersensitivity reactions. Careful consideration should be given to the patient's clinical status prior to administration of Vimizim and consider delaying the Vimizim

infusion.

Sleep apnea is common in MPS IVA patients. Evaluation of airway patency should be considered prior to initiation of treatment with Vimizim. Patients using supplemental oxygen or continuous positive airway pressure (CPAP) during sleep should have these treatments readily available during infusion in the event of an acute reaction, or extreme drowsiness/sleep induced by antihistamine use.

5.3 Spinal or Cervical Cord Compression

Spinal or cervical cord compression (SCC) is a known and serious complication of MPS IVA and may occur as part of the natural history of the disease. In clinical trials, SCC was observed both in patients receiving Vimizim and patients receiving placebo. Patients with MPS IVA should be monitored for signs and symptoms of SCC (including back pain, paralysis of limbs below the level of compression, urinary and fecal incontinence) and given appropriate clinical care.

6 ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The following serious adverse reactions are described below and elsewhere in the labeling:

- Anaphylaxis and hypersensitivity reactions [see *Warnings and Precautions (5.1)*].

The most common adverse reactions ($\geq 10\%$) observed across pre-marketing clinical trials were similar in type and frequency as those observed in the placebo-controlled trial (see Table 1). The acute reactions requiring intervention were managed by either temporarily interrupting or discontinuing infusion, and administering additional antihistamine, antipyretics, or corticosteroids.

6.1 Clinical Trials Experience

A 24-week, randomized, double-blind, placebo-controlled clinical trial of Vimizim was conducted in 176 patients with MPS IVA, ages 5 to 57 years old. Approximately half of the patients (49%) were male. Of the 176 patients, 65% were White, 23% Asian, 3% Black, and 10% Other race. The majority of patients (78%) were non-Hispanic. Patients were randomized to three treatment groups: Vimizim 2 mg/kg once per week (n=58), Vimizim 2 mg/kg once every other week (n=59), or placebo (n=59). All patients were treated with antihistamines prior to each infusion.

Table 1 summarizes the most common adverse reactions that occurred in the placebo-controlled trial with an incidence of $\geq 10\%$ in patients treated with Vimizim 2 mg/kg once per week and with a higher incidence than in the placebo-treated patients.

Table 1: Adverse Reactions That Occurred in the Placebo-Controlled Trial in At Least 10% of Patients in the Vimizim 2 mg/kg Once Per Week Group and with a Higher Incidence than in the Placebo Group

Adverse Reaction	Vimizim 2 mg/kg once per week	Placebo
	N= 58 n (%)	N= 59 n (%)
Pyrexia	19 (33%)	8 (14%)
Vomiting	18 (31%)	4 (7%)
Headache	15 (26%)	9 (15%)
Nausea	14 (24%)	4 (7%)
Abdominal pain	12 (21%)	1 (1.7%)
Chills	6 (10.3%)	1 (1.7%)

Fatigue	6 (10.3%)	2 (3.4%)
---------	-----------	----------

Extension Trial

An open-label extension trial was conducted in 173 patients who completed the placebo-controlled trial [see *Clinical Studies (14)*]. No new adverse reactions were reported.

6.2 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. All patients treated with Vimizim 2 mg/kg once per week in the placebo-controlled trial developed anti-drug antibodies by Week 4. Anti-drug antibody titers were sustained or increased for the duration of Vimizim treatment. Because all patients developed anti-drug antibodies, associations between antibody titers and reductions in treatment effect or the occurrence of anaphylaxis or other hypersensitivity reactions could not be determined.

All patients treated with Vimizim 2 mg/kg once per week tested positive for neutralizing antibodies capable of inhibiting the drug from binding to the mannose-6-phosphate receptor at least once during the trial. Binding to this receptor is required for Vimizim to be taken into cells where it is active.

Neutralizing antibody titers were not determined in the patients. Therefore, the possibility of an association between neutralizing antibody titer and treatment effect cannot be assessed.

Assessment of the incidence of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to Vimizim with the incidence of antibodies to other products may be misleading.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

There is a Morquio A Registry that collects data on pregnant women with MPS IVA who are treated with Vimizim. Contact MARS@bmrn.com or call 1-800-983-4587 for information and enrollment [see *Patient Counseling Information (17)*].

Risk Summary

There are no adequate and well-controlled studies with Vimizim in pregnant women. However, animal reproduction studies have been conducted for elosulfase alfa. In these studies, no effects on embryo-fetal development were observed in rats given daily administration of elosulfase alfa up to 33 times the human steady-state AUC (area under the curve) at the recommended human weekly dose pre-mating and through the period of organogenesis. No effects on embryo-fetal development were observed in rabbits given daily administration of elosulfase alfa at doses up to 8 times the human steady-state AUC at the recommended weekly dose during organogenesis, which produced maternal toxicity. A dose-dependent increase in stillbirths was observed when elosulfase alfa was administered daily in rats during organogenesis through lactation at doses 5 times the human steady-state AUC at the recommended human weekly dose. An increase in pup mortality was observed at doses producing maternal toxicity. Vimizim should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Clinical Considerations

Disease-associated maternal and embryo/fetal risk

Pregnancy can adversely affect the health of females affected with MPS IVA and lead to adverse

pregnancy outcomes for both mother and fetus.

Animal Data

All reproductive studies with rats included pre-treatment with diphenhydramine to prevent or minimize hypersensitivity reactions. The effects of elosulfase alfa were evaluated based on comparison to a control group treated with diphenhydramine alone. Daily intravenous (IV) administration of up to 20 mg/kg elosulfase alfa in rats (33 times the human steady-state AUC at the recommended weekly dose of 2 mg/kg) during a 15-day pre-mating period, mating, and the period of organogenesis, produced no maternal toxicity or effects on embryo-fetal development. Daily intravenous administration of up to 10 mg/kg in rabbits (8 times the human steady-state AUC at the recommended weekly dose) during the period of organogenesis had no effects on embryo-fetal development. However, maternal toxicity (gross changes in liver) was observed in rabbits given doses of 1 mg/kg/day and higher (0.1 times the human steady-state AUC at the recommended weekly dose). Elosulfase alfa produced an increase in the percentage of stillbirths when administered daily to rats at doses of 6 mg/kg IV and higher (5 times the human steady-state AUC at the recommended weekly dose) during the period of organogenesis through lactation. Daily administration of 20 mg/kg IV (33 times the human steady-state AUC at the recommended weekly dose) produced maternal toxicity and an increase in mortality of offspring during the lactation period. This study lacked a full evaluation of neurodevelopmental milestones; however, no effects of elosulfase alfa were noted in tests for learning and memory.

8.3 Nursing Mothers

It is not known if Vimizim is present in human milk. Elosulfase alfa is present in milk from treated rats [see *Use in Specific Populations (8.1)*]. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Vimizim and any potential adverse effects on the breastfed child from the drug or from MPS IVA. Exercise caution when administering Vimizim to a nursing mother. There is a Morquio A Registry that also collects data on breastfeeding women with MPS IVA who are treated with Vimizim. Contact MARS@bmrn.com or call 1-800-983-4587 for information and enrollment [see *Patient Counseling Information (17)*].

8.4 Pediatric Use

Safety and effectiveness of Vimizim have been established in pediatric patients 5 years of age and older. Use of Vimizim in patients 5 years of age and older is supported by an adequate and well-controlled trial in pediatric and adult patients. Clinical trials with Vimizim were conducted in 176 patients (median age 12 years, range 5 to 57 years old) with the majority of patients in the pediatric age group (53% aged 5 to 11 years, 27% aged 12 to 17 years) [see *Clinical Studies (14)*]. Safety and effectiveness in pediatric patients below 5 years of age have not been established.

8.5 Geriatric Use

Clinical studies of Vimizim did not include any patients aged 65 and over. It is not known whether they respond differently from younger patients.

10 OVERDOSAGE

There is no experience with overdose with Vimizim.

11 DESCRIPTION

Vimizim is a formulation of elosulfase alfa, which is a purified human enzyme produced by recombinant DNA technology in a Chinese hamster ovary cell line. Human N-acetylgalactosamine-6-sulfatase (EC 3.1.6.4) is a hydrolytic lysosomal glycosaminoglycan-specific enzyme that hydrolyzes sulfate from either galactose-6-sulfate or N-acetyl-galactosamine-6-sulfate on the non-reducing ends of the glycosaminoglycans keratan sulfate (KS) and chondroitin-6-sulfate (C6S).

Elosulfase alfa is a soluble glycosylated dimeric protein with two oligosaccharide chains per monomer. Each monomeric peptide chain contains 496 amino acids and has an approximate molecular mass of 55 kDa (59 kDa including the oligosaccharides). One of the oligosaccharide chains contains bis-mannose-6-phosphate (bisM6P). bisM6P binds a receptor at the cell surface and the binding mediates cellular uptake of the protein to the lysosome. Elosulfase alfa has a specific activity of 2.6 to 6.0 units/mg. One activity unit is defined as the amount of the enzyme required to convert 1 micromole of sulfated monosaccharide substrate D-galactopyranoside-6-sulfate (Gal-6S) to de-sulfated-galactose (Gal) and free sulfate per minute at 37°C.

Vimizim is intended for intravenous infusion and is supplied as a sterile, nonpyrogenic, colorless to pale yellow, clear to slightly opalescent solution that must be diluted with 0.9% Sodium Chloride for Injection, USP prior to administration. Vimizim is supplied in clear Type 1 glass 5 mL vials. Each vial provides 5 mg elosulfase alfa, 31.6 mg L-arginine hydrochloride, 0.5 mg polysorbate 20, 13.6 mg sodium acetate trihydrate, 34.5 mg sodium phosphate monobasic monohydrate, and 100 mg sorbitol in a 5 mL extractable solution with a pH between 5.0 to 5.8. Vimizim does not contain preservatives. Each vial is for single use only.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Mucopolysaccharidoses comprise a group of lysosomal storage disorders caused by the deficiency of specific lysosomal enzymes required for the catabolism of glycosaminoglycans (GAG).

Mucopolysaccharidosis IVA (MPS IVA, Morquio A Syndrome) is characterized by the absence or marked reduction in N-acetylgalactosamine-6-sulfatase activity. The sulfatase activity deficiency results in the accumulation of the GAG substrates, KS and C6S, in the lysosomal compartment of cells throughout the body. The accumulation leads to widespread cellular, tissue, and organ dysfunction. Vimizim is intended to provide the exogenous enzyme N-acetylgalactosamine-6-sulfatase that will be taken up into the lysosomes and increase the catabolism of the GAGs KS and C6S. Elosulfase alfa uptake by cells into lysosomes is mediated by the binding of mannose-6-phosphate-terminated oligosaccharide chains of elosulfase alfa to mannose-6-phosphate receptors.

In the absence of an animal disease model that recapitulates the human disease phenotype, elosulfase alfa pharmacological activity was evaluated using human primary chondrocytes from two MPS IVA patients. Treatment of MPS IVA chondrocytes with elosulfase alfa induced clearance of KS lysosomal storage from the chondrocytes.

12.2 Pharmacodynamics

The pharmacodynamic effect of Vimizim was assessed by reductions in urinary KS levels. The relationship of urinary KS to other measures of clinical response has not been established [see *Clinical Studies (14)*]. No association was observed between antibody development and urinary KS levels.

12.3 Pharmacokinetics

The pharmacokinetics of elosulfase alfa were evaluated in 23 patients with MPS IVA who received intravenous infusions of Vimizim 2 mg/kg once weekly, over approximately 4 hours, for 22 weeks. Eleven patients were aged 5 to 11 years, six were aged 12 to 17 years, and six were aged 18 to 41 years. Table 2 summarizes the pharmacokinetic parameters at Week 0 and Week 22. Mean AUC_{0-t} and C_{max} increased to 2.8- and 2.9-fold, respectively, at Week 22 compared to Week 0. Mean t_{1/2} increased from 7.5 min at Week 0 to 35.9 min at Week 22. These changes are likely related to the development of neutralizing antibodies in all patients.

Table 2: Pharmacokinetic Parameters

	Week 0 (N = 22)*	Week 22 (N = 22)*
--	------------------	-------------------

Pharmacokinetic Parameter	Mean (SD)	Mean (SD)
AUC _{0-t} , min x µg/mL [†]	238 (100)	577 (416)
C _{max} , µg/mL [‡]	1.49 (0.534)	4.04 (3.24)
T _{max} , min [§]	172 (75.3)	202 (90.8)
CL, mL/min/kg [¶]	10.0 (3.73) [#]	7.08 (13.0) [♣]
V _{dss} , mL/kg [♥]	396 (316) [♦]	650 (1842) [♣]
t _{1/2} , min [♣]	7.52 (5.48) [#]	35.9 (21.5) [♣]

* The pharmacokinetics of elosulfase alfa was evaluated in 23 individual patients. However, 1 patient was not tested at Week 0 and another patient was not tested at Week 22.

[†]AUC_{0-t}, area under the plasma concentration-time curve from time zero to the time of last measurable concentration;

[‡]C_{max}, observed maximum plasma concentration;

[§]T_{max}, time from zero to maximum plasma concentration;

[¶]CL, total clearance of drug after intravenous administration;

[#]N = 15;

[♣]N = 20

[♥]V_{dss}, apparent volume of distribution at steady-state;

[♦]N = 14;

[♣]t_{1/2}, elimination half-life

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals to evaluate carcinogenic potential or studies to evaluate mutagenic potential have not been performed with elosulfase alfa. Based on the mechanism of action, elosulfase alfa is not expected to be tumorigenic. Daily intravenous administration of elosulfase alfa in rats at doses up to 20 mg/kg (55 times the human steady-state AUC in male rats and 33 times the human steady-state AUC in female rats at the recommended human weekly dose) had no effects on fertility or reproductive performance.

14 CLINICAL STUDIES

The safety and efficacy of Vimizim were assessed in a 24-week, randomized, double-blind, placebo-controlled clinical trial of 176 patients with MPS IVA. The age of patients ranged from 5 to 57 years. The majority of the patients (82%) presented with a medical history of musculoskeletal conditions, which includes knee deformity (52%), kyphosis (31%), hip dysplasia (22%), prior spinal fusion surgery (22%) and arthralgia (20%). At baseline, all enrolled patients could walk more than 30 meters (m) but less than 325 m in six minutes.

Patients received Vimizim 2 mg/kg once per week (n=58), Vimizim 2 mg/kg once every other week (n=59), or placebo (n=59).

The primary endpoint was the change from baseline in the distance walked in six minutes (six minute walk test, 6-MWT) at Week 24. The other endpoints included changes from baseline in the rate of stair climbing in three minutes (three-minute stair climb test, 3-MSCT) and changes from baseline in urine KS levels at Week 24. The treatment effect in the distance walked in 6 minutes, compared to placebo, was 22.5 m (CI₉₅, 4.0, 40.9; p=0.0174) in patients who received Vimizim 2 mg/kg once per week. There was no difference in the rate of stair climbing between patients who received Vimizim 2 mg/kg once per

week and those who received placebo. Patients who received Vimizim 2 mg/kg once every other week performed similarly in the 6-MWT and 3-MSCT as those who received placebo. The reduction in urinary KS levels from baseline, a measure of pharmacodynamic effect, was greater in the Vimizim treatment groups compared to placebo. The relationship between urinary KS and other measures of clinical response has not been established.

Table 3: Results from Placebo-Controlled Clinical Trial

	Vimizim 2 mg/kg once per week			Placebo			Vimizim vs. Placebo
	Baseline	Week 24	Change	Baseline	Week 24	Change	Mean Difference in Changes
N	58	57*	57	59	59	59	
Six-Minute Walk Test (Meters)							
Mean	203.9	243.3	36.5	211.9	225.4	13.5	23.0 [†]
± SD	± 76.32	± 83.53	± 58.49	± 69.88	± 83.22	± 50.63	(CI ₉₅ , 2.9, 43.1)
Median	216.5	251.0	20.0	228.9	229.4	9.9	22.5 [‡]
Min,	42.4,	52.0,	-57.8,	36.2,	50.6,	-99.2,	(CI ₉₅ , 4.0, 40.9)
Max	321.5	399.9	228.7	312.2	501.0	220.5	(p = 0.0174) ^{‡,§}

* One patient in the Vimizim group dropped out after 1 infusion

[†] Observed Vimizim mean change – Placebo mean change

[‡] ANCOVA Model-based Vimizim mean change – Placebo mean change, adjusted for baseline 6MWT categories (less than or equal to 200 meters, greater than 200 meters) and age groups (5-11, 12-18, 19 or older)

[§] p-value based on the model-based difference in means

Extension Trial

Patients who participated in the placebo-controlled trial were eligible to continue treatment in an open-label extension trial. One hundred seventy-three of 176 patients enrolled in the extension trial in which patients received Vimizim 2 mg/kg once per week (n=86) or Vimizim 2 mg/kg once every other week (n=87). In patients who continued to receive Vimizim 2 mg/kg once per week for another 48 weeks (for a total of 72-week exposure), walking ability showed no further improvement beyond the first 24 weeks of treatment in the placebo-controlled trial.

16 HOW SUPPLIED/STORAGE AND HANDLING

Vimizim is supplied as a concentrated solution for infusion (1 mg per mL) requiring dilution. One vial of 5 mL contains 5 mg Vimizim.

NDC 68135-100-01, 5 mL vial

Store Vimizim under refrigeration at 2°C to 8°C (36°F to 46°F). Do not freeze or shake. Protect from light.

Diluted Vimizim should be used immediately. If immediate use is not possible, diluted Vimizim may be stored for up to 24 hours at 2°C to 8°C (36°F to 46°F) followed by up to 24 hours at 23°C to 27°C (73°F to 81°F) during administration.

17 PATIENT COUNSELING INFORMATION

Anaphylactic Reactions

Advise the patients and caregivers that reactions related to administration and infusion may occur during

Vimizim treatment, including life-threatening anaphylaxis. Patients who have experienced anaphylactic reactions may require observation during and after Vimizim administration. Inform patients of the signs and symptoms of anaphylaxis and have them seek immediate medical care should symptoms occur. The risks and benefits of re-administering Vimizim following a severe reaction should be considered. Patients with acute respiratory illness may be at risk of serious acute exacerbation of their respiratory compromise due to hypersensitivity reactions. Pre-medication and reduction of infusion rate may alleviate those reactions associated with the infusion [see Warnings and Precautions (5.1, 5.2)].

Morquio A Registry

Inform patients of the Morquio A Registry established in order to better understand the variability and progression of the disease in the population as a whole, and to monitor and evaluate long-term treatment effects of Vimizim. The Morquio A Registry will also monitor the effect of Vimizim on pregnant women, nursing mothers and their offspring, and determine if Vimizim is excreted in breast milk. Patients should be encouraged to participate and advised that their participation is voluntary and may involve long-term follow-up. For more information, contact MARS@bmrn.com or call 1-800-983-4587.

Manufactured by:

BioMarin Pharmaceutical Inc. Novato, CA 94949

US License Number 1649

1-866-906-6100 (phone)

V1/2014

PACKAGE LABEL

NDC 68135-100-01

VIMIZIMTM
(elosulfase alfa)

Injection

5 mg/5 mL (1 mg/mL)

For intravenous infusion

Must be diluted prior to use

Rx Only

For single use only.
Discard unused portions.
Usual dosage: see package insert.
Store at 2°C to 8°C (36°F to 46°F)
Do not freeze
Manufactured by:
BioMarin Pharmaceutical Inc.
Novato, CA 94949
U.S. License No. 1649

NDC 68135-100-01
Vimizim[™]
(elosulfase alfa)
Injection
5 mg/5 mL (1 mg/mL)
For intravenous infusion
Must be diluted prior to use
Rx Only
BIOMARIN[®]

Varnish free area for printing of LOT and EXP

Vimizim US Vial

NDC 68135-100-01


VIMIZIM™
(elosulfase alfa)

5 mg/5 mL
(1 mg/mL)

For intravenous infusion

Must be diluted prior to use

Rx Only



VIMIZIM

elosulfase alfa injection, solution, concentrate

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:68 135-100
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ELOSULFASE ALFA (UNII: ODJ69JZG85) (ELOSULFASE ALFA - UNII:ODJ69JZG85)	ELOSULFASE ALFA	5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ARGININE HYDRO CHLORIDE (UNII: F7LTHIE20Y)	31.6 mg in 5 mL
SORBITOL (UNII: 506T60A25R)	100 mg in 5 mL
POLYSORBATE 20 (UNII: 7T1F30V5YH)	0.5 mg in 5 mL
WATER (UNII: 059QF0KO0R)	
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	34.5 mg in 5 mL
SODIUM ACETATE (UNII: 4550K0SC9B)	13.6 mg in 5 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68135-100-01	1 in 1 CARTON	02/14/2014	
1		5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA125460	02/14/2014	

Labeler - BioMarin Pharmaceutical Inc. (079722386)**Establishment**

Name	Address	ID/FEI	Business Operations
AndersonBrecon (UK) Ltd.		762771269	LABEL(68135-100) , PACK(68135-100)

Establishment

Name	Address	ID/FEI	Business Operations
AndersonBrecon Inc.		053217022	LABEL(68135-100) , PACK(68135-100)

Establishment

Name	Address	ID/FEI	Business Operations
BioMarin International Limited		985066780	ANALYSIS(68135-100) , API MANUFACTURE(68135-100) , LABEL(68135-100) , PACK(68135-100)

Establishment

Name	Address	ID/FEI	Business Operations
BioMarin Pharmaceutical Inc.		010004135	ANALYSIS(68135-100) , API MANUFACTURE(68135-100)

Establishment			
Name	Address	ID/FEI	Business Operations
Vetter Pharma Fertigung GmbH & Co. KG		316 126 754	ANALYSIS(68 135-100)

Establishment			
Name	Address	ID/FEI	Business Operations
Vetter Pharma Fertigung GmbH & Co. KG		344217323	ANALYSIS(68 135-100) , MANUFACTURE(68 135-100)

Establishment			
Name	Address	ID/FEI	Business Operations
Vetter Pharma-Fertigung GmbH & Co. KG		312670654	ANALYSIS(68 135-100) , MANUFACTURE(68 135-100)

Revised: 2/2014

BioMarin Pharmaceutical Inc.