

**POTASSIUM CHLORIDE- potassium chloride tablet, film coated, extended release**  
**Slate Run Pharmaceuticals, LLC**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use POTASSIUM CHLORIDE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for POTASSIUM CHLORIDE EXTENDED-RELEASE TABLETS.**

**POTASSIUM CHLORIDE extended-release tablets, for oral use**

**Initial U.S. Approval: 1948**

----- **INDICATIONS AND USAGE** -----

Potassium Chloride Extended-Release Tablets are a potassium salt, indicated for the treatment and prophylaxis of hypokalemia with or without metabolic alkalosis in patients for whom dietary management with potassium-rich foods or diuretic dose reduction is insufficient. (1)

----- **DOSAGE AND ADMINISTRATION** -----

- Monitor serum potassium and adjust dosages accordingly (2.1)
- If serum potassium is less than 2.5 mEq/L, use intravenous potassium instead of oral supplementation (2.1)
- Take with meals and with a glass of water or other liquid. Swallow tablets whole without crushing, chewing or sucking. (2.1)
- Treatment of hypokalemia: Doses range from 40 to 100 mEq/day in divided doses. Limit doses to 40 mEq per dose. (2.2)
- Prevention of hypokalemia: Typical dose is 20 mEq per day. (2.2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Tablets: 600 mg (8 mEq) and 750 mg (10 mEq) (3)

----- **CONTRAINDICATIONS** -----

- Concomitant use with triamterene and amiloride (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Gastrointestinal Irritation: Take with meals (5.1)

----- **ADVERSE REACTIONS** -----

- The most common adverse reactions are nausea, vomiting, flatulence, abdominal pain/discomfort and diarrhea. (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Slate Run Pharmaceuticals, LLC at 1-888-341-9214 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

----- **DRUG INTERACTIONS** -----

- Triamterene and amiloride: Concomitant use is contraindicated (7.1)
- Renin-angiotensin-aldosterone inhibitors: Monitor for hyperkalemia (7.2)
- Nonsteroidal anti-inflammatory drugs: Monitor for hyperkalemia (7.3)

----- **USE IN SPECIFIC POPULATIONS** -----

- Cirrhosis: Initiate therapy at the low end of the dosing range (8.6)
- Renal Impairment: Initiate therapy at the low end of the dosing range (8.7)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 11/2019**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

2.1 Administration and Monitoring

2.2 Dosing

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Gastrointestinal Adverse Reactions

### **6 ADVERSE REACTIONS**

### **7 DRUG INTERACTIONS**

7.1 Triamterene or amiloride

7.2 Renin-angiotensin-aldosterone Inhibitors

7.3 Nonsteroidal Anti-inflammatory Drugs (NSAIDs)

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Cirrhotics

8.7 Renal Impairment

### **10 OVERDOSAGE**

10.1 Symptoms

10.2 Treatment

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Potassium Chloride Extended-Release Tablets are indicated for the treatment and prophylaxis of hypokalemia with or without metabolic alkalosis, in patients for whom dietary management with potassium-rich foods or diuretic dose reduction is insufficient.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Administration and Monitoring**

If serum potassium concentration is less than 2.5 mEq/L, use intravenous potassium instead of oral supplementation.

## *Monitoring*

Monitor serum potassium and adjust dosages accordingly. Monitor serum potassium periodically during maintenance therapy to ensure potassium remains in desired range.

The treatment of potassium depletion, particularly in the presence of cardiac disease, renal disease, or acidosis, requires careful attention to acid-base balance, volume status, electrolytes, including magnesium, sodium, chloride, phosphate, and calcium, electrocardiograms, and the clinical status of the patient. Correct volume status, acid-base balance, and electrolyte deficits as appropriate.

## *Administration*

Take potassium chloride extended-release tablets with meals and with a glass of water or other liquid. Do not take potassium chloride extended-release tablets on an empty stomach because of its potential for gastric irritation [see *Warnings and Precautions (5.1)*].

Swallow tablets whole without crushing, chewing or sucking.

## **2.2 Dosing**

Dosage must be adjusted to the individual needs of each patient. Dosages greater than 40 mEq per day should be divided such that no more than 40 mEq is given in a single dose.

*Treatment of Hypokalemia:* Typical dose range is 40 to 100 mEq per day.

*Maintenance or Prophylaxis:* Typical dose range is 20 mEq per day.

## **3 DOSAGE FORMS AND STRENGTHS**

Potassium Chloride Extended-Release Tablets, USP are supplied as:

The 8 mEq (600 mg) tablets are white to off-white, round, biconvex, coated tablets, debossed with “YH” and “106” on one side, and “8” on the other side.

The 10 mEq (750 mg) tablets are white to off-white, round, biconvex, coated tablets, debossed with “YH” and “105” on one side, and “10” on the other side.

## **4 CONTRAINDICATIONS**

Potassium chloride is contraindicated in patients on triamterene and amiloride.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Gastrointestinal Adverse Reactions**

Solid oral dosage forms of potassium chloride can produce ulcerative and/or stenotic lesions of the gastrointestinal tract, particularly if the drug maintains contact with the gastrointestinal mucosa for prolonged periods. Consider the use of liquid potassium in patients with dysphagia, swallowing disorders, or severe gastrointestinal motility disorders.

If severe vomiting, abdominal pain, distention, or gastrointestinal bleeding occurs, discontinue potassium chloride extended-release tablets and consider possibility of ulceration, obstruction or perforation.

Potassium chloride extended-release tablets should not be taken on an empty stomach because of its potential for gastric irritation [see *Dosage and Administration (2.1)*].

## **6 ADVERSE REACTIONS**

The following adverse reactions have been identified with use of oral potassium salts. Because these

reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The most common adverse reactions to oral potassium salts are nausea, vomiting, flatulence, abdominal pain/discomfort, and diarrhea.

There have been reports hyperkalemia and of upper and lower gastrointestinal condition including obstruction, bleeding, ulceration, perforation.

Skin rash has been reported rarely.

## **7 DRUG INTERACTIONS**

### **7.1 Triamterene or amiloride**

Use with triamterene or amiloride can produce severe hyperkalemia. Concomitant use is contraindicated [*see Contraindications (4)*].

### **7.2 Renin-angiotensin-aldosterone Inhibitors**

Drugs that inhibit the renin-angiotensin-aldosterone system (RAAS) including angiotensin converting enzyme (ACE) inhibitors, angiotensin receptor blockers (ARBs), spironolactone, eplerenone, or aliskiren produce potassium retention by inhibiting aldosterone production. Closely monitor potassium in patients on concomitant RAAS inhibitors.

### **7.3 Nonsteroidal Anti-inflammatory Drugs (NSAIDs)**

NSAIDs may produce potassium retention by reducing renal synthesis of prostaglandin E and impairing the renin-angiotensin system. Closely monitor potassium in patients on concomitant NSAIDs.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no human data related to use of potassium chloride extended-release tablets during pregnancy, and animal reproduction studies have not been conducted. Potassium supplementation that does not lead to hyperkalemia is not expected to cause fetal harm.

The background risk for major birth defects and miscarriage in the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

### **8.2 Lactation**

#### Risk Summary

The normal potassium ion content of human milk is about 13 mEq per liter. Since oral potassium becomes part of the body potassium pool, so long as body potassium is not excessive, the contribution of potassium chloride supplementation should have little or no effect on the level in human milk.

### **8.4 Pediatric Use**

Safety and effectiveness in the pediatric population have not been established.

### **8.5 Geriatric Use**

Clinical studies of potassium chloride extended-release tablets did not include sufficient numbers of

subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **8.6 Cirrhotics**

Based on published literature, the baseline corrected serum concentrations of potassium measured over 3 hours after administration in cirrhotic subjects who received an oral potassium load rose to approximately twice that of normal subjects who received the same load. Patients with cirrhosis should usually be started at the low end of the dosing range, and the serum potassium level should be monitored frequently [see *Clinical Pharmacology* (12.3)] .

## **8.7 Renal Impairment**

Patients with renal impairment have reduced urinary excretion of potassium and are at substantially increased risk of hyperkalemia. Patients with impaired renal function, particularly if the patient is on RAAS inhibitors or NSAIDs, should usually be started at the low end of the dosing range because of the potential for development of hyperkalemia [see *Drug Interactions* ( 7.2, 7.3)] . The serum potassium level should be monitored frequently. Renal function should be assessed periodically.

# **10 OVERDOSAGE**

## **10.1 Symptoms**

The administration of oral potassium salts to persons with normal excretory mechanisms for potassium rarely causes serious hyperkalemia. However, if excretory mechanisms are impaired, potentially fatal hyperkalemia can result [see *Contraindications and Warnings*].

It is important to recognize that hyperkalemia is usually asymptomatic and may be manifested only by an increased serum potassium concentration (6.5 to 8.0 mEq/L) and characteristic electrocardiographic changes (peaking of T-waves, loss of P-wave, depression of S-T segment and prolongation of the QT interval). Late manifestations include muscle paralysis and cardiovascular collapse from cardiac arrest (9 to 12 mEq/L).

## **10.2 Treatment**

Treatment measures for hyperkalemia include the following:

1. Elimination of foods and medications containing potassium and of any agents with potassium-sparing properties.
2. Intravenous administration of 300 to 500 mL/hr of 10% dextrose solution containing 10 to 20 units of crystalline insulin per 1,000 mL.
3. Correction of acidosis, if present, with intravenous sodium bicarbonate.
4. Use of exchange resins, hemodialysis or peritoneal dialysis.

In treating hyperkalemia, it should be recalled that in patients who have been stabilized on digitalis, too rapid a lowering of the serum potassium concentration can produce digitalis toxicity.

The extended release feature means that absorption and toxic effects may be delayed for hours. Consider standard measures to remove any unabsorbed drug.

## 11 DESCRIPTION

Potassium Chloride Extended-Release Tablets, USP are a solid oral dosage form of potassium chloride. Each contains 600 mg or 750 mg of potassium chloride equivalent to 8 mEq or 10 mEq of potassium in a wax matrix tablet.

Potassium Chloride Extended-Release Tablets, USP are an electrolyte replenisher. The chemical name is potassium chloride, and the structural formula is KCl. Potassium chloride, USP is a white, granular powder or colorless crystals. It is odorless and has a saline taste. Its solutions are neutral to litmus. It is freely soluble in water and insoluble in alcohol.

*Inactive Ingredients:* colloidal silicon dioxide, hydroxypropyl cellulose and hydrogenated vegetable oil. The film coating contains lecithin, polyethylene glycol, polyvinyl alcohol, talc and titanium dioxide.

FDA approved dissolution test specifications differ from USP.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The potassium ion is the principal intracellular cation of most body tissues. Potassium ions participate in a number of essential physiological processes including the maintenance of intracellular tonicity, the transmission of nerve impulses, the contraction of cardiac, skeletal and smooth muscle and the maintenance of normal renal function.

The intracellular concentration of potassium is approximately 150 to 160 mEq per liter. The normal adult plasma concentration is 3.5 to 5 mEq per liter. An active ion transport system maintains this gradient across the plasma membrane.

Potassium is a normal dietary constituent and under steady state conditions the amount of potassium absorbed from the gastrointestinal tract is equal to the amount excreted in the urine. The usual dietary intake of potassium is 50 to 100 mEq per day.

### 12.3 Pharmacokinetics

The potassium chloride in potassium chloride extended-release tablets is completely absorbed before it leaves the small intestine. The wax matrix is not absorbed and is excreted in the feces; in some instances the empty matrices may be noticeable in the stool. When the bioavailability of the potassium ion from the potassium chloride extended-release tablets is compared to that of a true solution the extent of absorption is similar.

The extended-release properties of potassium chloride extended-release tablets are demonstrated by the finding that a significant increase in time is required for renal excretion of the first 50% of the potassium chloride extended-release tablets dose as compared to the solution.

Increased urinary potassium excretion is first observed 1 hour after administration of potassium chloride extended-release tablets, reaches a peak at approximately 4 hours, and extends up to 8 hours. Mean daily steady-state plasma levels of potassium following daily administration of potassium chloride extended-release tablets cannot be distinguished from those following administration of potassium chloride solution or from control plasma levels of potassium ion.

#### Specific Populations

##### *Cirrhotics*

Based on published literature, the baseline corrected serum concentrations of potassium measured over 3 hours after administration in cirrhotic subjects who received an oral potassium load rose to approximately twice that of normal subjects who received the same load.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

Carcinogenicity, mutagenicity and fertility studies in animals have not been performed. Potassium is a normal dietary constituent.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Potassium Chloride Extended-Release Tablets, USP contain 600 mg or 750 mg of Potassium Chloride, USP (equivalent to 8 mEq and 10 mEq respectively), and are provided as extended release tablets.

**Table 1: How Supplied**

Dose	Shape	Color	Debossment	NDC #: 70436-xxx-xx		
				Bottle of 100 tablets	Bottle of 500 tablets	Bottle of 1000 tablets
600 mg (8 mEq)	Round	White to off-white	“YH” and “106” on one side, and “8” on the other side	001-01	001-02	001-03
750 mg (10 mEq)	Round	White to off-white	“YH” and “105” on one side, and “10” on the other side	002-01	002-02	002-03

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature.] Protect from light and moisture.

Dispense in a tight container as defined in the USP with a child-resistant closure.

## 17 PATIENT COUNSELING INFORMATION

- Inform patients to take each dose with meals and with a full glass of water or other liquid, and to not crush, chew, or suck the tablets. Inform patients that the wax matrix is not absorbed and is excreted in the feces; in some instances the empty matrices may be noticeable in the stool.
- Advise patients seek medical attention if tarry stools or other evidence of gastrointestinal bleeding is noticed.

Manufactured by:  
Yichang Humanwell Oral Solid Dosage Plant  
Yichang, Hubei, China 443112

Distributed by:  
Slate Run Pharmaceuticals, LLC  
Columbus, Ohio 43215

Revised 08/2019

## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

### NDC 70436-001-01

Potassium Chloride Extended-release Tablets, USP

8 mEq (600 mg)

100 Tablets

Rx only

NDC 70436-001-01 Rx only

## Potassium Chloride Extended-Release Tablets USP

**8 mEq (600 mg)**

**Usual Dose:** See accompanying package insert for full prescribing information. Dosage must be adjusted to the individual needs of each patient.

**100 Tablets**


**Each extended-release tablet contains:**  
Potassium Chloride, USP 600 mg.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight container as defined in the USP with a child-resistant closure.

**For Patient's Information:** Be aware that the expended matrix is not absorbed and may be excreted intact in the stool.

**Keep out of reach of children.**

Manufactured by: Yichang Humanwell Oral Solid Dosage Plant, Yichang, Hubei, China 443112  
Distributed by: Slate Run Pharmaceuticals, LLC, Columbus, Ohio 43215  
1000001/02 Rev. 08/2019



**NDC 70436-001-02**

Potassium Chloride Extended-release Tablets, USP

8 mEq (600 mg)

500 Tablets

Rx only

NDC 70436-001-02 Rx only

## Potassium Chloride Extended-Release Tablets USP

**8 mEq (600 mg)**

**Usual Dose:** See accompanying package insert for full prescribing information. Dosage must be adjusted to the individual needs of each patient.

**500 Tablets**


**Each extended-release tablet contains:**  
Potassium Chloride, USP 600 mg.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight container as defined in the USP with a child-resistant closure.

**For Patient's Information:** Be aware that the expended matrix is not absorbed and may be excreted intact in the stool.

**Keep out of reach of children.**

Manufactured by: Yichang Humanwell Oral Solid Dosage Plant, Yichang, Hubei, China 443112  
Distributed by: Slate Run Pharmaceuticals, LLC, Columbus, Ohio 43215  
1000002/02 Rev. 08/2019



**NDC 70436-001-03**

Potassium Chloride Extended-release Tablets, USP

8 mEq (600 mg)

1000 Tablets

Rx only



NDC 70436-001-03

Rx only

# Potassium Chloride Extended-Release Tablets USP

**8 mEq (600 mg)**

**Usual Dose:** See accompanying package insert for full prescribing information. Dosage must be adjusted to the individual needs of each patient.

SLATE RUN  
PHARMACEUTICALS

1000 Tablets

**Each extended-release tablet contains:**  
Potassium Chloride, USP 600 mg.


Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight container as defined in the USP with a child-resistant closure.

**For Patient's Information:** Be aware that the expended matrix is not absorbed and may be excreted intact in the stool.

**Keep out of reach of children.**

Manufactured by: Yichang Humanwell Oral Solid Dosage Plant, Yichang, Hubei, China 443112  
Distributed by: Slate Run Pharmaceuticals, LLC, Columbus, Ohio 43215  
10000003/02

Rev. 08/2019



## NDC 70436-002-01

Potassium Chloride Extended-release Tablets, USP

10 mEq (750 mg)

100 Tablets

Rx only

NDC 70436-002-01

Rx only

# Potassium Chloride Extended-Release Tablets USP

**10 mEq (750 mg)**

**Usual Dose:** See accompanying package insert for full prescribing information. Dosage must be adjusted to the individual needs of each patient.

SLATE RUN  
PHARMACEUTICALS

100 Tablets

**Each extended-release tablet contains:**  
Potassium Chloride, USP 750 mg.


Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight container as defined in the USP with a child-resistant closure.

**For Patient's Information:** Be aware that the expended matrix is not absorbed and may be excreted intact in the stool.

**Keep out of reach of children.**

Manufactured by: Yichang Humanwell Oral Solid Dosage Plant, Yichang, Hubei, China 443112  
Distributed by: Slate Run Pharmaceuticals, LLC, Columbus, Ohio 43215  
10000004/02

Rev. 08/2019



## NDC 70436-002-02

Potassium Chloride Extended-release Tablets, USP

10 mEq (750 mg)

500 Tablets

Rx only

NDC 70436-002-02 Rx only

# Potassium Chloride Extended-Release Tablets USP

10 mEq (750 mg)

**Usual Dose:** See accompanying package insert for full prescribing information. Dosage must be adjusted to the individual needs of each patient.

**Each extended-release tablet contains:**  
Potassium Chloride, USP 750 mg.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight container as defined in the USP with a child-resistant closure.


**For Patient's Information:** Be aware that the expended matrix is not absorbed and may be excreted intact in the stool.

**Keep out of reach of children.**

Manufactured by: Yichang Humanwell Oral Solid Dosage Plant, Yichang, Hubei, China 443112  
Distributed by: Slate Run Pharmaceuticals, LLC, Columbus, Ohio 43215  
10000005/02

Rev. 08/2019





500 Tablets

**NDC 70436-002-03**

Potassium Chloride Extended-release Tablets, USP

10 mEq (750 mg)

1000 Tablets

Rx only

NDC 70436-002-03 Rx only

# Potassium Chloride Extended-Release Tablets USP

10 mEq (750 mg)

**Usual Dose:** See accompanying package insert for full prescribing information. Dosage must be adjusted to the individual needs of each patient.

**Each extended-release tablet contains:**  
Potassium Chloride, USP 750 mg.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light and moisture. Dispense in a tight container as defined in the USP with a child-resistant closure.

**For Patient's Information:** Be aware that the expended matrix is not absorbed and may be excreted intact in the stool.

**Keep out of reach of children.**

Manufactured by: Yichang Humanwell Oral Solid Dosage Plant, Yichang, Hubei, China 443112  
Distributed by: Slate Run Pharmaceuticals, LLC, Columbus, Ohio 43215  
10000006/02

Rev. 08/2019





1000 Tablets

**POTASSIUM CHLORIDE**

potassium chloride tablet, film coated, extended release

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:70436-002
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
POTASSIUM CHLORIDE (UNII: 660YQ981I0) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CATION	750 mg

## Inactive Ingredients

Ingredient Name	Strength
HYDROGENATED COTTONSEED OIL (UNII: Z82Y2C65EA)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
TALC (UNII: 7SEV7J4R1U)	

## Product Characteristics

Color	white (white to off-white)	Score	no score
Shape	ROUND (Biconvex)	Size	13mm
Flavor		Imprint Code	YH;105;10
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70436-002-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2018	
2	NDC:70436-002-02	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2018	
3	NDC:70436-002-03	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209314	11/15/2018	

## POTASSIUM CHLORIDE

potassium chloride tablet, film coated, extended release

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70436-001
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POTASSIUM CHLORIDE (UNII: 660YQ981I0) (POTASSIUM CATION - UNII:295O53K152)	POTASSIUM CATION	600 mg

## Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>HYDROGENATED COTTONSEED OIL</b> (UNII: Z82Y2C65EA)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>POLYVINYL ALCOHOL</b> (UNII: 532B59J990)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

### Product Characteristics

<b>Color</b>	white (white to off-white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Biconvex)	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	YH;106;8
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70436-001-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2018	
2	NDC:70436-001-02	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2018	
3	NDC:70436-001-03	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209314	11/15/2018	

**Labeler** - Slate Run Pharmaceuticals, LLC (039452765)

**Registrant** - Yichang Humanwell Pharmaceutical Co., Ltd. (527225336)

### Establishment

Name	Address	ID/FEI	Business Operations
Yichang Humanwell Oral Solid Dosage Plant		421371958	manufacture(70436-001, 70436-002)

Revised: 11/2019

Slate Run Pharmaceuticals, LLC