

OPHYTE TATTOO NUMBING CREAM- numbing cream cream
Changsha Beiliang E-commerce Co., Ltd

83887-002

Active Ingredient

Lidocaine 5%

Purpose

Local Anesthetic

Use

For temporary relief of local and anorectal itching, discomfort, and inflammation associated with anorectal disorders.

For temporary relief of pain, burning, and soreness.

Warnings

If symptoms worsen or do not improve within 7 days, seek medical advice.

Do not exceed the recommended daily dosage without doctor's instruction.

Consult a doctor immediately if bleeding occurs.

Do not insert this product into the rectum using fingers or any applicator.

Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.

Note: Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients

Do not use

Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.

When Using

When Using

If symptoms worsen or do not improve within 7 days, seek medical advice. Do not

exceed the recommended daily dosage without doctor's instruction. Consult a doctor immediately if bleeding occurs.

Do not insert this product into the rectum using fingers or any applicator.

Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.

Stop Use

Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients.

Ask Doctor

Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients

Keep Out Of Reach Of Children

This package is child-resistant. Keep out of reach of children

Directions

Adults: When practical, cleanse the affected area with mild soap and warm water, and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.

Other information

Keep away from direct sunlight or heat.

Store at room temperature between 15°C and 30°C (59°F-86°F).

This package is child-resistant. Keep out of reach of children. In case of accidental overdose or ingestion, call a doctor or poison control center immediately. Do not use this product if the seal is broken or missing.

Inactive ingredients

Vitamin E Allantoin, Mineral Oil, Dimethicone, Methylparaben, Propylparaben, Petrolatum, IPBC, Diazolidinyl Urea, Glycerin, Cetostearyl Alcohol, Isopropyl Myristate, Propylene Glycol, Ammonium Methacrylate, Purified Water, Laurocapram, Steareth-2, Steareth-21, Rose Oil

PRINCIPAL DISPLAY PANEL



USAGE METHOD

- Clean and dry the area of skin where the tattoo will be applied
- Apply a generous amount of cream onto the skin
- Cover the cream-coated area with plastic wrap for approximately 40-60 minutes
- Remove plastic wrap, gently pat to help cream absorb. Wipe off residue, wait 15-20 mins for maximum numbness before proceeding.

Distributed by:

Ophye care LLC
1242 Broadway St Ste 314C
Boulder, CO 80502
Tel:1-719-201-4680

Manufactured by:

Changsha Beilang E-commerce Co., Ltd
Changsha, Hunan, China-410000
www.pressycare.com

SHELF LIFE: 3 YEARS



Ophye

Tattoo
NUMBING CREAM

6 HOURS
MAXIMUM STRENGTH
Numbing Cream with
Vitamin E

EXTERNAL
NET: 30g/1.0oz

Warnings (For external use only)

- If symptoms worsen or do not improve within 7 days, seek medical advice.
- Do not exceed the recommended daily dosage without doctor's instruction.
- Consult a doctor immediately if bleeding occurs.
- Do not insert this product into the rectum using fingers or any applicator.
- Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain. **Note:** Discontinue use and consult a doctor if symptoms persist or worsen including redness, irritation, swelling, or pain due to possible allergies to product ingredients.

Inactive Ingredients

Vitamin E, Allantoin, Mineral Oil, Dimethicone, Methylparaben, Propylparaben, Pterocissum, IPAC, Decylolyl Urea, Glycerin, Cetylstearyl Alcohol, Isopropyl Myristate, Propylene Glycol, Ammonium Methacrylate, Purified Water, Laureth-2, Steareth-21, Rose Oil.

DRUG FACTS

Active Ingredient	Purpose
Lidocaine 5%	Local Anesthetic

Uses

- For temporary relief of local and anorectal itching, discomfort, and inflammation associated with anorectal disorders.
- For temporary relief of pain, burning, and soreness.

Directions

- Adults: When practical, cleanse the affected area with mild soap and warm water, and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.
- Children under 12 years of age: Consult a doctor. Keep out of reach of children.
- Apply externally to the affected area up to 6 times daily.

Other Information

- Keep away from direct sunlight or heat.
- Store at room temperature between 15°C and 30°C (59°F - 86°F).
- This package is child-resistant. Keep out of reach of children.
- In case of accidental overdose or ingestion, call a doctor or poison control center immediately.
- Do not use this product if the seal is broken or missing.

■ C:100 M:90 Y:0 K:0

■ C:0 M:50 Y:100 K:0

MFG: 12/11/2023
EXP: 12/10/2026
LOT: 20231211-01



C:100 M:90 Y:0 K:0

C:0 M:50 Y:100 K:0

DRUG FACTS

Active Ingredient
Lidocaine HCl

Purpose
Local Anesthetic

Uses
For temporary relief of local and mucosal itching, discomfort, and inflammation associated with insect bites/stings.

Warnings (See external use only)
 • For external use only.
 • If symptoms worsen or do not improve within 7 days, seek medical advice.
 • Do not exceed the recommended daily dosage, amount or duration.
 • Discontinue use immediately if bleeding occurs.
 • Do not exceed the labeled directions using fingers or any applicator.
 • Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including rash, irritation, swelling, or pain.
 • **Notes:** Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients.

DRUG FACTS (Continued)

Directions
 • Adults: When needed, massage the affected area with mild soap and warm water, and rinse thoroughly. Gently dry by patting or blotting with a clean tissue or cloth, and then reapplication of the product.
 • Children: Under 17 years of age: Consult a doctor. Keep out of reach of children.
 • Keep container tightly closed when not in use.

Other Information
 • Keep away from direct sunlight or heat.
 • Store at room temperature between 20°C and 25°C (68°F - 80°F).
 • The package is child-resistant. Keep out of reach of children.
 • In case of accidental overdose or ingestion, call a doctor or poison control center immediately.
 • Do not use the product if the seal is broken or missing.

Inactive Ingredients
 Valerol, E. Alanolol, Mineral Oil, Dimethylsiloxane, Methylparaben, Propylparaben, Propylene Glycol, Disodium EDTA, Stearic Acid, Citric Acid, Benzyl Alcohol, Methylparaben, Propylparaben, Ammonium Methacrylate, Purified Water, Hydroxypropyl Methylcellulose, Benzyl Alcohol, Benzyl Alcohol.

USAGE METHOD:

- Clean and dry the area of skin where the tattoo will be applied.
- Apply a generous amount of cream onto the skin.
- Cover the cream-treated area with plastic wrap for approximately 40-60 minutes.
- Remove plastic wrap, gently pat to help cream absorb. Wipe off residue, wait 15-20 min for treatment numbness before proceeding.

Distributed by:
 Opiyo Cosmetics
 1842 Broadway St. Ste 214C
 Beverly, CA 90705
 Tel: 310-778-2000
 www.opiyo.com

Manufactured by:
 Dingyha (Beijing) Cosmetics Co., Ltd.
 Dingyha, Hainan, China 570000
 www.opiyo.com

100% VEGAN
 100% CRUELTY FREE

EXTERNAL

Tattoo Opiyo

NUMBING CREAM

6 HOURS MAXIMUM STRENGTH

NET: 50g/1.75oz

MFG: 12/11/2023
 EXP: 12/10/2026
 LOT: 20231211-01

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
IODOPROPYNYL BUTYLCARBAMATE (UNII: 603P14DHEB)
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)
ALLANTOIN (UNII: 344S277G0Z)
GLYCERIN (UNII: PDC6A3C0OX)
ROSE OIL (UNII: WUB68Y35M7)
DIMETHICONE (UNII: 92RU3N3Y1O)
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)
METHYLPARABEN (UNII: A2I8C7HI9T)
AMMONIUM METHACRYLATE (UNII: J2243103QO)
STEARETH-2 (UNII: V56DFE46J5)
MINERAL OIL (UNII: T5L8T28FGP)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
PETROLATUM (UNII: 4T6H12BN9U)
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)
LAUROCAPRAM (UNII: 1F3X9DRV9X)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
WATER (UNII: 059QF0KO0R)
STEARETH-21 (UNII: 53J3F32P58)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83887-002-01	30 g in 1 TUBE; Type 0: Not a Combination Product	12/22/2023	
2	NDC:83887-002-02	50 g in 1 BOTTLE; Type 0: Not a Combination Product	12/22/2023	
3	NDC:83887-002-03	60 g in 1 BOTTLE; Type 0: Not a Combination Product	12/22/2023	
4	NDC:83887-002-04	100 g in 1 BOTTLE; Type 0: Not a Combination Product	12/22/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	12/22/2023	

Labeler - Changsha Beiliang E-commerce Co., Ltd (632138945)

Establishment

Name	Address	ID/FEI	Business Operations
Changsha Beiliang E-commerce Co., Ltd		632138945	label(83887-002) , manufacture(83887-002)