

HURRICAINA TOPICAL ANESTHETIC- benzocaine gel
Beutlich Pharmaceuticals, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HurriCaine Gel LV Syringe - Mint

Active ingredient

Benzocaine 20%

Purpose

Oral Anesthetic

Uses

for the temporary relief of occasional minor irritation and pain, associated with:

- canker sores
- sore mouth and throat
- minor injury of the mouth and gums
- minor dental procedures
- minor irritation of the mouth and gums caused by dentures or orthodontic appliances

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or irritation, pain, or redness persists or worsens, see your dentist or doctor promptly.

Do not use

Do not use:

- if the tray seal is broken or missing
- if cap is missing or damaged
- for teething

- in children under 2 years of age

When using this product

- avoid contact with eyes

Keep out of reach of children.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

Directions**Directions:**

Do not exceed recommended dosage.

- adults and children 2 years of age and older: apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.
- children under 12 years of age: should be supervised in the use of the product
- children under 2 years of age: do not use

Other information

- store at 20°-25° C (68°-77° F)

Inactive ingredients

- flavor, polyethylene glycol, sodium saccharin

Questions or comments?

1-800-238-8542

M-F: 8:00 a.m. - 4:30 p.m. ET

Principal Display Panel

Manufactured for:
Beutlich Pharmaceuticals, LLC
Bunnell, FL 32110

NDC 0283-0920-34



FRESH MINT

HURRICAIN[®]
Topical Anesthetic GEL
20% Benzocaine Oral Anesthetic

HURRICAIN[®]
Topical Anesthetic GEL
20% Benzocaine Oral Anesthetic

Drug Facts	
Active ingredient Benzocaine 20%	Purpose Oral Anesthetic
Uses for the temporary relief of occasional minor irritation and pain, associated with <ul style="list-style-type: none"> • canker sores • sore mouth and throat • minor injury of the mouth and gums • minor dental procedures • minor irritation of the mouth and gums caused by dentures or orthodontic appliances 	
Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: <ul style="list-style-type: none"> • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy Allergy alert: Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.	

Drug Facts (continued)							
Warnings (continued) Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. If sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persist or worsens, see your dentist or doctor promptly. Do not use <ul style="list-style-type: none"> • if tray seal is broken or missing • for teething • in children under 2 years of age When using this product avoid contact with eyes Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.							
Directions do not exceed recommended dosage <table border="0"> <tr> <td>adults and children 2 years of age and older</td> <td>apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.</td> </tr> <tr> <td>children under 12 years of age</td> <td>should be supervised in the use of the product</td> </tr> <tr> <td>children under 2 years of age</td> <td>do not use</td> </tr> </table>		adults and children 2 years of age and older	apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.	children under 12 years of age	should be supervised in the use of the product	children under 2 years of age	do not use
adults and children 2 years of age and older	apply to the affected area. Allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a dentist or doctor.						
children under 12 years of age	should be supervised in the use of the product						
children under 2 years of age	do not use						

Drug Facts (continued)			
Other information store at 20-25 °C (68-77 °F)			
Inactive ingredients flavor, polyethylene glycol, sodium saccharin			
Questions or comments? 1-800-238-8542 M-F: 8:00 a.m. - 4:30 p.m. ET			
Please retain Drug Facts for future reference. HurriCaine [®] is a registered trademark of Beutlich Pharmaceuticals, LLC, 7775 S. US Hwy 1, Suite H, Bunnell, FL 32110. www.beutlich.com			
<table border="1"> <tr> <td>LOT</td> <td>No UV Coating in this Area</td> </tr> </table>		LOT	No UV Coating in this Area
LOT	No UV Coating in this Area		

HSPM 889 1120

Gel Syringe Unit: 0283-1820-43 1.5g in 1 syringe
 Gel Syringe Pack of 4: 0283-1820-34 4 x 1.5g in 1 box.



HURRICAIN[®] TOPICAL ANESTHETIC

benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0283-0920
Route of Administration	ORAL, DENTAL, PERIODONTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0283-0920-34	4 in 1 BOX	01/31/2021	
1	NDC:0283-0920-43	1.5 g in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	01/31/2021	

Labeler - Beutlich Pharmaceuticals, LLC (005209325)

Registrant - Beutlich Pharmaceuticals, LLC (005209325)

Establishment

Name	Address	ID/FEI	Business Operations
Dental Technologies, Inc.		148312838	manufacture(0283-0920)