

ANTACID ANTIGAS MAXIMUM STRENGTH- aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Chain Drug Consortium, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients

Aluminum hydroxide (equivalent to Dried Gel, USP) 400 mg

Magnesium hydroxide 400 mg

Simethicone 40 mg

Purpose

Antacid

Antigas

Uses

relieves: • heartburn • sour stomach • acid indigestion
• upset stomach due to these symptoms • symptoms of gas

Warnings

Ask a doctor before use if you have
• kidney disease • a magnesium restricted diet

Ask a doctor or pharmacist before use if you are

presently taking a prescription drug. Antacids may interact with certain prescription drugs

Stop use and ask a doctor

if symptoms last for more
than 2 weeks

If pregnant or breast-feeding

ask a health professional before use.

Keep this and all drugs out of the reach of children.

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Other information

store at room temperature tightly closed

do not freeze

each teaspoon contains:

potassium 2 mg, magnesium 167 mg

Inactive ingredients

benzyl alcohol,
flavors, glycerin, hydroxyethyl cellulose,
potassium citrate, propylene glycol,
purified water, sorbitol

Directions

- shake well before use
- do not exceed 12 teaspoons (60 mL) in a 24 hour period or use the maximum dosage for more than 2 weeks
- adults and children 12 years of age and older: take 2 to 4 teaspoons (10-20 mL) between meals and at bedtime or as directed by a doctor
- children under 12 years of age: ask a doctor

Package Label

Drug Facts

Active ingredients

(in each 5 mL = 1 teaspoon)

		Purpose
Aluminum hydroxide (equivalent to Dried Gel, USP)	400 mg.....	Antacid
Magnesium hydroxide	400 mg.....	Antacid
Simethicone	40 mg.....	Antigas

Uses relieves: • heartburn • sour stomach
• acid indigestion • upset stomach due to these symptoms
• symptoms of gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last for more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep this and all drugs out of the reach of children.

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Directions • shake well before use

- do not exceed 12 teaspoons (60 mL) in a 24 hour period or use the maximum dosage for more than 2 weeks
- adults & children 12 years of age and older: take 2 to 4 teaspoons (10-20 mL) between meals and at bedtime or as directed by a doctor
- children under 12 years of age: ask a doctor

Other information • do not freeze

- store at room temperature tightly closed

• **each teaspoon contains:**

potassium 2 mg, magnesium 167 mg

Inactive ingredients benzyl alcohol,

flavors, glycerin, hydroxyethyl cellulose,

potassium citrate, propylene glycol,

purified water, sorbitol

TAMPER EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Johnson & Johnson Merck, distributor of Mylanta®

Distributed by: Chain Drug Consortium, LLC.,
2300 NW Corporate Blvd., Suite 115
Boca Raton, FL 33431

Questions or Comments? 1-866-534-4631

68-01374B



Antacid Antigas Maximum Strength

ANTACID ANTIGAS MAXIMUM STRENGTH

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:68016-050

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE (2000 MPAS AT 1%) (UNII: S38J6RZN16)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	LEMON (Lemon)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-050-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part331	06/03/2009	

Labeler - Chain Drug Consortium, LLC (101668460)

Revised: 2/2016

Chain Drug Consortium, LLC