

# HEPARIN SODIUM IN SODIUM CHLORIDE- heparin sodium injection, solution

## Baxter Healthcare Corporation

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use HEPARIN SODIUM IN SODIUM CHLORIDE INJECTION safely and effectively. See full prescribing information for HEPARIN SODIUM IN SODIUM CHLORIDE INJECTION.

HEPARIN SODIUM IN SODIUM CHLORIDE injection, for intravenous use

Initial U.S. Approval: 1939

### RECENT MAJOR CHANGES

Dosage and Administration Preparation of Administration (2.1) 01/2022

Warnings and Precautions (5.2, 5.6, 5.7) 01/2022

### INDICATIONS AND USAGE

Heparin Sodium in Sodium Chloride Injection at a concentration of 2 units/mL is indicated as an anticoagulant to maintain catheter patency. (1)

### DOSAGE AND ADMINISTRATION

Infuse through intravenous catheter at a rate of 6 units per hour. (2.2)

### DOSAGE FORMS AND STRENGTHS

- Injection: 1,000 USP units in Sodium Chloride in 500 mL single-dose infusion bag (2 USP units per mL) (3)
- Injection: 2,000 USP units in Sodium Chloride per 1,000 mL single-dose infusion bag (2 USP units per mL) (3)

### CONTRAINDICATIONS

Heparin Sodium in Sodium Chloride Injection is contraindicated in patients with the following conditions: (4)

- With an uncontrollable active bleeding state, except when this is due to disseminated intravascular coagulation (5.1)
- With a history of heparin-induced thrombocytopenia (HIT) or heparin-induced thrombocytopenia with thrombosis (HITT) (5.2)
- With severe thrombocytopenia (5.2, 5.3)
- Known hypersensitivity to heparin or pork products (5.5)

### WARNINGS AND PRECAUTIONS

- Hemorrhage: Fatal cases have occurred. Monitor for signs of bleeding and manage promptly. (5.1)
- HIT and HITT: Monitor for signs and symptoms and discontinue if indicative of HIT or HITT. (5.2)
- Hyperkalemia: Measure plasma potassium in patients at risk of hyperkalemia before starting heparin therapy and periodically in all patients. (5.6)
- Elevations of serum aminotransferases: Interpret elevation of these enzymes with caution. (5.7)

### ADVERSE REACTIONS

Most common adverse reactions are: hemorrhage, thrombocytopenia, HIT and HITT, hypersensitivity, and elevations of aminotransferase levels. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare at 1-866-888-2472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

Drugs that interfere with platelet aggregation or drugs that counteract coagulation may induce bleeding (7.2)

### USE IN SPECIFIC POPULATIONS

Geriatric Use: A higher incidence of bleeding has been reported in patients over 60 years of age, especially women (5.7, 8.5)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 1/2022**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Preparation for Administration
- 2.2 Maintenance of Catheter Patency

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Hemorrhage
- 5.2 Heparin-Induced Thrombocytopenia and Heparin-Induced Thrombocytopenia with Thrombosis
- 5.3 Thrombocytopenia
- 5.4 Heparin Resistance
- 5.5 Hypersensitivity
- 5.6 Hyperkalemia
- 5.7 Elevations of Serum Aminotransferases
- 5.8 Laboratory Tests

### **6 ADVERSE REACTIONS**

- 6.1 Postmarketing Experience

### **7 DRUG INTERACTIONS**

- 7.1 Oral Anticoagulants
- 7.2 Platelet Inhibitors
- 7.3 Other Medications that May Interfere with Heparin

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Heparin Sodium in Sodium Chloride Injection at a concentration of 2 units/mL is indicated as an anticoagulant to maintain catheter patency.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Preparation for Administration

Do not administer unless solution is clear, and seal is intact. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit. Do not infuse this product under pressure.

To Open:

Tear overwrap down side at slit and remove solution container. Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Check for minute leaks by squeezing inner bag firmly. If leaks are found discard solution as sterility may be impaired. Do not add supplementary medication.

Preparation for Administration:

1. Suspend container from eyelet support.
2. Remove plastic protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

All injections in VIAFLEX PLUS plastic containers are intended for administration using sterile equipment.

Single-dose container.

Discard any unused portion.

Do not reconnect partially used bags.

Because dosages of this drug are titrated to response, **no additives should be made to Heparin Sodium in Sodium Chloride Injection.**

Heparin sodium is not effective by oral administration and Heparin Sodium in Sodium Chloride Injection should not be given orally.

### 2.2 Maintenance of Catheter Patency

The recommended starting dose is 6 units per hour by intravenous infusion through an intravenous catheter to maintain catheter patency.

## 3 DOSAGE FORMS AND STRENGTHS

- Injection: 1,000 USP units per 500 mL (2 units per mL) clear solution in a single-dose infusion bag
- Injection: 2,000 USP units per 1,000 mL (2 units per mL) clear solution in a single-

dose infusion bag

## 4 CONTRAINDICATIONS

The use of Heparin Sodium in Sodium Chloride Injection is contraindicated in patients with the following conditions:

- Uncontrollable active bleeding state, except when this is due to disseminated intravascular coagulation [see *Warnings and Precautions (5.1)*]
- History of heparin-induced thrombocytopenia (HIT) and heparin-induced thrombocytopenia with thrombosis (HITT) [see *Warnings and Precautions (5.2)*]
- Severe thrombocytopenia [see *Warnings and Precautions (5.2, 5.3)*]
- Known hypersensitivity to heparin or pork products (e.g., anaphylactoid reactions) [see *Adverse Reactions (6.1)*]

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Hemorrhage

Avoid using heparin in the presence of major bleeding, except when the benefits of heparin therapy outweigh the potential risks.

Hemorrhage can occur at virtually any site in patients receiving heparin. Fatal hemorrhages have occurred. A higher incidence of bleeding has been reported in patients, particularly women, over 60 years of age [see *Use in Specific Populations (8.5)*]. An unexplained fall in hematocrit, fall in blood pressure, or any other unexplained symptom should lead to serious consideration of hemorrhagic event.

Use heparin sodium with caution in disease states in which there is increased risk of hemorrhage, including:

- Cardiovascular - Subacute bacterial endocarditis, severe hypertension.
- Surgical - During and immediately following (a) spinal tap or spinal anesthesia or (b) major surgery, especially involving the brain, spinal cord, or eye.
- Hematologic - Conditions associated with increased bleeding tendencies, such as hemophilia, thrombocytopenia, and some vascular purpuras.
- Gastrointestinal - Ulcerative lesions and continuous tube drainage of the stomach or small intestine.
- Patients with hereditary antithrombin III deficiency receiving concurrent antithrombin III therapy - The anticoagulant effect of heparin is enhanced by concurrent treatment with antithrombin III (human) in patients with hereditary antithrombin III deficiency. To reduce the risk of bleeding, reduce the heparin dose during concomitant treatment with antithrombin III (human).
- Other - Menstruation, liver disease with impaired hemostasis.

### 5.2 Heparin-Induced Thrombocytopenia and Heparin-Induced Thrombocytopenia with Thrombosis

Heparin-induced thrombocytopenia (HIT) is a serious immune-mediated reaction. HIT occurs in patients treated with heparin and is due to the development of antibodies to a platelet Factor 4-heparin complex that induce *in vivo* platelet aggregation. HIT may

progress to the development of venous and arterial thromboses, a condition referred to as heparin-induced thrombocytopenia with thrombosis (HITT). Thrombotic events may also be the initial presentation for HITT. These serious thromboembolic events include deep vein thrombosis, pulmonary embolism, cerebral vein thrombosis, limb ischemia, stroke, myocardial infarction, mesenteric thrombosis, renal arterial thrombosis, skin necrosis, gangrene of the extremities that may lead to amputation, and possibly death.

Once HIT or HITT is diagnosed or strongly suspected, discontinue all heparin sources (including heparin flushes) and use an alternative anticoagulant.

Immune-mediated HIT is diagnosed based on clinical findings supplemented by laboratory tests confirming the presence of antibodies to heparin, or platelet activation induced by heparin. Obtain platelet counts at baseline and periodically during heparin administration. A drop in platelet count greater than 50% from baseline is considered indicative of HIT. Platelet counts begin to fall 5 to 10 days after exposure to heparin in heparin-naïve individuals and reach a threshold by days 7 to 14. In contrast, “rapid onset” HIT can occur very quickly (within 24 hours following heparin initiation), especially in patients with a recent exposure to heparin (i.e., previous 3 months). Thrombosis development shortly after documenting thrombocytopenia is a characteristic finding in almost half of all patients with HIT.

Monitor any degree of thrombocytopenia closely. If the platelet count falls below 100,000/mm<sup>3</sup> or if recurrent thrombosis develops, promptly discontinue heparin, evaluate for HIT and HITT, and, if necessary, administer an alternative anticoagulant. HIT or HITT can occur up to several weeks after the discontinuation of heparin therapy. Evaluate patients presenting with thrombocytopenia or thrombosis after discontinuation of heparin sodium for HIT or HITT.

### **5.3 Thrombocytopenia**

Thrombocytopenia in patients receiving heparin has been reported at frequencies up to 30%. It can occur 2 to 20 days (average 5 to 9) following the onset of heparin therapy. Obtain platelet counts before and periodically during heparin therapy. Mild thrombocytopenia (count greater than 100,000/mm<sup>3</sup>) may remain stable or reverse even if heparin is continued. However, monitor thrombocytopenia of any degree closely. If the count falls below 100,000/mm<sup>3</sup> or if recurrent thrombosis develops, promptly discontinue heparin, evaluate for HIT and HITT, and, if necessary, administer an alternative anticoagulant [*see Warnings and Precautions (5.2)*].

### **5.4 Heparin Resistance**

Increased resistance to heparin is frequently encountered in fever, thrombosis, thrombophlebitis, infections with thrombosing tendencies, myocardial infarction, cancer, in postsurgical patients, and patients with anti-thrombin deficiency. Consider measurement of anti-thrombin levels if heparin resistance is suspected. Monitor coagulation tests frequently in such patients. It may be necessary to adjust the dose of heparin based on coagulation test monitoring, such as anti-Factor Xa levels and/or partial thromboplastin time.

### **5.5 Hypersensitivity**

Hypersensitivity reactions with chills, fever and urticaria as the most usual manifestations and also asthma, rhinitis, lacrimation, and anaphylactoid reactions have

been reported. Patients with documented hypersensitivity to heparin should be given the drug only in clearly life-threatening situations. Because Heparin Sodium in Sodium Chloride Injection is derived from animal tissue, it should be used with caution in patients with a history of allergy to pork products.

## **5.6 Hyperkalemia**

Heparin can suppress adrenal secretion of aldosterone leading to hyperkalemia, particularly in patients with diabetes mellitus, chronic renal failure, pre-existing metabolic acidosis, a raised plasma potassium, or taking potassium sparing drugs. The risk of hyperkalemia appears to increase with duration of therapy but is usually reversible upon discontinuation of heparin.

Measure plasma potassium in patients at risk of hyperkalemia before starting heparin therapy and periodically in all patients treated for more than 5 days or earlier as deemed fit by the clinician.

## **5.7 Elevations of Serum Aminotransferases**

Significant elevations of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels have occurred patients who have received heparin. Elevation of these enzymes in patients receiving heparin should be interpreted with caution. These elevations typically resolve upon heparin discontinuation.

## **5.8 Laboratory Tests**

Periodic platelet counts, hematocrits, and tests for occult blood in stool are recommended during the entire course of heparin therapy, regardless of the route of administration.

## **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are described elsewhere in the labeling:

- Hemorrhage [*see Warnings and Precautions (5.1)*]
- Heparin-Induced Thrombocytopenia and Heparin-Induced Thrombocytopenia with Thrombosis [*see Warnings and Precautions (5.2)*]
- Thrombocytopenia [*see Warnings and Precautions (5.3)*]
- Heparin Resistance [*see Warnings and Precautions (5.4)*]
- Hypersensitivity [*see Warnings and Precautions (5.5)*]
- Hyperkalemia [*see Warnings and Precautions (5.6)*]
- Elevations of Serum Aminotransferases [*see Warnings and Precautions (5.7)*]

### **6.1 Postmarketing Experience**

The following adverse reactions have been identified during post-approval use of heparin sodium. Because the reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency.

#### Hemorrhage

Hemorrhage is the chief complication that may result from heparin therapy [*see*

*Warnings and Precautions (5.1)].* An overly prolonged clotting time or minor bleeding during therapy can usually be controlled by withdrawing the drug [see *Overdosage (10)*]. Gastrointestinal or urinary tract bleeding during anticoagulant therapy may indicate the presence of an underlying occult lesion. Bleeding can occur at any site but certain specific hemorrhagic complications may be difficult to detect:

- Adrenal hemorrhage, with resultant acute adrenal insufficiency, has occurred during anticoagulant therapy, including fatal cases.
- Ovarian (corpus luteum) hemorrhage developed in a number of women of reproductive age receiving short or long-term anticoagulant therapy.
- Retroperitoneal hemorrhage

**Vascular Disorders:** Contusion, Vasospastic reactions (including episodes of painful, ischemic, and cyanosed limbs).

*HIT and HITT, including delayed onset cases, Thrombocytopenia [see Warnings and Precautions (5.2, 5.3)].*

*Histamine-like reactions:* Such reactions have been observed at the site of injections. Necrosis of the skin has been reported at the site of subcutaneous injection of heparin, occasionally requiring skin grafting.

*Hypersensitivity:* General hypersensitivity reactions have been reported, with chills, fever, and urticaria as the most usual manifestations, and asthma, rhinitis, lacrimation, headache, nausea and vomiting, and anaphylactoid reactions, including shock, occurring more rarely. Itching and burning, especially on the plantar site of the feet, may occur [see *Warnings and Precautions (5.5)*].

*Musculoskeletal, Connective Tissue and Bone Disorders:* Osteoporosis with long-term administration of heparin.

*Metabolism and Nutrition Disorders:* Hyperkalemia.

*General Disorders and Administration Site Conditions:* Erythema, mild pain, ulceration.

*Elevations of serum aminotransferases:* Significant elevations of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels have occurred patients who have received heparin.

*Others:* Cutaneous necrosis after systemic administration, delayed transient alopecia, priapism, and rebound hyperlipemia on discontinuation of heparin sodium have also been reported.

## **7 DRUG INTERACTIONS**

### **7.1 Oral Anticoagulants**

Heparin sodium may prolong the one-stage prothrombin time. Therefore, when heparin sodium is given with dicumarol or warfarin sodium, a period of at least 5 hours after the last intravenous dose or 24 hours after the last subcutaneous dose should elapse before blood is drawn if a valid prothrombin time is to be obtained.

### **7.2 Platelet Inhibitors**

Drugs such as NSAIDs (including acetylsalicylic acid, ibuprofen, indomethacin, and celecoxib), dextran, phenylbutazone, thienopyridines, dipyridamole, hydroxychloroquine, glycoprotein IIb/IIIa antagonists (including abciximab, eptifibatide, and tirofiban), and others that interfere with platelet-aggregation reactions (the main hemostatic defense of heparinized patients) may induce bleeding and should be used with caution in patients receiving heparin sodium. To reduce the risk of bleeding, a reduction in the dose of antiplatelet agent or heparin is recommended.

### **7.3 Other Medications that May Interfere with Heparin**

Digitalis, tetracyclines, nicotine or antihistamines may partially counteract the anticoagulant action of heparin sodium. Intravenous nitroglycerin administered to heparinized patients may result in a decrease of the partial thromboplastin time with subsequent rebound effect upon discontinuation of nitroglycerin. Careful monitoring of partial thromboplastin time and adjustment of heparin dosage are recommended during coadministration of heparin and intravenous nitroglycerin.

Antithrombin III (human) – The anticoagulant effect of heparin is enhanced by concurrent treatment with antithrombin III (human) in patients with hereditary antithrombin III deficiency. To reduce the risk of bleeding, a reduced dosage of heparin is recommended during treatment with antithrombin III (human).

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no available data on Heparin Sodium in Sodium Chloride Injection use in pregnant women to inform a drug-associated risk of major birth defects and miscarriage. In published reports heparin exposure during pregnancy did not show evidence of an increased risk of adverse maternal or fetal outcomes in humans (see *Data*). Consider the benefits and risks of Heparin Sodium in Sodium Chloride Injection for the mother and possible risks to the fetus when prescribing Heparin Sodium in Sodium Chloride Injection to a pregnant woman.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. However, the background risk in the U.S. general population of major birth defects is 2 to 4% and of miscarriage is 15 to 20% of clinically recognized pregnancies.

#### Data

##### *Human Data*

The maternal and fetal outcomes associated with uses of heparin via various dosing methods and administration routes during pregnancy have been investigated in numerous studies. These studies generally reported normal deliveries with no maternal or fetal bleeding and no other complications.

##### *Animal Data*

In a published study conducted in rats and rabbits, pregnant animals received heparin



intravenously during organogenesis at a dose of 10,000 units/kg/day, more than 50 times the maximum human daily dose based on body weight. The number of early resorptions increased in both species. There was no evidence of teratogenic effects.

## 8.2 Lactation

### Risk Summary

There is no information regarding the presence of heparin in human milk, the effects on the breastfed child, or the effects on milk production. Due to its large molecular weight, heparin is not likely to be excreted in human milk.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for heparin and any potential adverse effects on the breastfed child from heparin or from the underlying maternal condition.

## 8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

## 8.5 Geriatric Use

A higher incidence of bleeding has been reported in patients over 60 years of age, especially women [see *Warnings and Precautions (5.1)*]. Clinical studies indicate that lower doses of heparin may be indicated in these patients [see *Clinical Pharmacology (12.3)*].

## 10 OVERDOSAGE

An overdose requires immediate medical attention and treatment. Bleeding is the chief sign of heparin overdose. Easy bruising, petechial formations, nosebleeds, blood in urine or tarry stools may be the first signs or symptoms of a heparin overdose. In the event of symptomatic heparin overdose, consider stopping heparin infusion.

Neutralization of heparin effect:

When clinical circumstances (bleeding) require reversal of heparinization, protamine sulfate (1% solution) by slow infusion will neutralize heparin sodium. **No more than 50 mg** should be administered, **very slowly** in any 10-minute period. Each mg of protamine sulfate neutralizes approximately 100 USP heparin units. The amount of protamine required decreases over time as heparin is metabolized. Although the metabolism of heparin is complex, it may, for the purpose of choosing a protamine dose, be assumed to have a half-life of about 30 minutes after intravenous injection.

Ideally, the dose required to neutralize the action of heparin should be guided by blood coagulation tests or calculated from a protamine neutralization test.


Because fatal reactions often resembling anaphylaxis have been reported with protamine, it should be given only when resuscitation techniques and treatment of anaphylactoid shock are readily available.

For additional information consult the labeling of Protamine Sulfate Injection.

Blood or plasma transfusions may be necessary; these dilute but do not neutralize heparin.

## 11 DESCRIPTION

Heparin is a heterogenous group of straight-chain anionic mucopolysaccharides, called glycosaminoglycans having anticoagulant properties. Although others may be present, the main sugars occurring in heparin are: (1)  $\alpha$ -L-iduronic acid 2-sulfate, (2) 2-deoxy-2-sulfamino- $\alpha$ -D-glucose 6-sulfate, (3)  $\beta$ -D-glucuronic acid, (4) 2-acetamido-2-deoxy- $\alpha$ -D-glucose, and (5)  $\alpha$ -L-iduronic acid. These sugars are present in decreasing amounts, usually in the order (2) > (1) > (4) > (3) > (5), and are joined by glycosidic linkages, forming polymers of varying sizes. Heparin is strongly acidic because of its content of covalently linked sulfate and carboxylic acid groups. In heparin sodium, the acidic protons of the sulfate units are partially replaced by sodium ions.



Heparin Sodium in Sodium Chloride Injection is a buffered, sterile, nonpyrogenic solution of Heparin Sodium, USP derived from porcine intestinal mucosa, standardized for anticoagulant activity supplied in single dose containers for vascular administration. It contains no antimicrobial agents. The potency is determined by a biological assay using a USP reference standard based on units of heparin activity per milligram. Composition, osmolarity, pH and ionic concentration are shown in Table 1.

**Table 1. Composition, Osmolarity, pH, and Ionic Concentration of Heparin in 0.9% Sodium Chloride Injection**

Product Description and	1,000 USP Heparin Units in 0.9% Sodium	1,000 USP Heparin Units in 0.9% Sodium Chloride	2,000 USP Heparin Units in 0.9% Sodium Chloride	2,000 USP Heparin Units in 0.9% Sodium

<b>Size</b>	<b>Chloride Injection, 500 mL</b>	<b>Chloride Injection, 500 mL</b>	<b>Chloride Injection, 1,000 mL</b>	<b>Chloride Injection, 1,000 mL</b>
NDC#	0338-0431-03	0338-0424-18	0338-0433-04	0338-0428-12
Manufacturing Location	Jayuya, PR	Toongabbie, AUS	Jayuya, PR	Toongabbie, AUS
Heparin Sodium, USP (units/mL)	2	2	2	2
Sodium Chloride, USP (NaCl) (g/L)	9	9	9	9
Dibasic Sodium Phosphate Heptahydrate, USP (Na <sub>2</sub> HPO <sub>4</sub> *7H <sub>2</sub> O) (g/L)	4.34	----	4.34	----
Dibasic Sodium Phosphate Dodecahydrate, USP (Na <sub>2</sub> HPO <sub>4</sub> *12H <sub>2</sub> O) (g/L)	----	5.80	----	5.80
Citric Acid Hydrous, USP (C <sub>6</sub> H <sub>8</sub> O <sub>7</sub> *H <sub>2</sub> O) (g/L)	0.4	0.4	0.4	0.4
Osmolarity* (mOsmol/L)	322	322	322	322
pH	7.0 (6.0 to 8.0)	7.0 (6.0 to 8.0)	7.0 (6.0 to 8.0)	7.0 (6.0 to 8.0)
Sodium (mEq/L)	186	186	186	186
Chloride (mEq/L)	154	154	154	154
Phosphate (as HPO <sub>4</sub> <sup>=</sup> ) (mEq/L)	32 (16 mmol/L)	32 (16 mmol/L)	32 (16 mmol/L)	32 (16 mmol/L)
Citrate (mEq/L)	6	6	6	6

\* Normal physiologic osmolarity range is approximately 280 to 310 mOsmol/L.

Administration of substantially hypertonic solutions ( $\geq$  600 mOsmol/L) may cause vein damage.

This VIAFLEX PLUS plastic container is fabricated from a specially formulated polyvinyl chloride (PL 146 Plastic). VIAFLEX PLUS on the container indicates the presence of a drug additive in a drug vehicle. The VIAFLEX PLUS plastic container system utilizes the same container as the VIAFLEX plastic container system. The amount of water that can permeate from inside the container into the overwrap is insufficient to affect the solution significantly. Solutions in contact with the plastic container can leach out certain of its chemical components in very small amounts within the expiration period, e.g., di-2-ethylhexyl phthalate (DEHP), up to 5 parts per million. However, the safety of the plastic

has been confirmed in tests in animals according to USP biological tests for plastic containers as well as by tissue culture toxicity studies.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Heparin interacts with the naturally occurring plasma protein, Antithrombin III, to induce a conformational change, which markedly enhances the serine protease activity of Antithrombin III, thereby inhibiting the activated coagulation factors involved in the closing sequence, particularly Xa and IIa. Small amounts of heparin inhibit Factor Xa, and larger amounts inhibit thrombin (Factor IIa). Heparin also prevents the formation of a stable fibrin clot by inhibiting the activation of the fibrin stabilizing factor. Heparin does not have fibrinolytic activity; therefore, it will not lyse existing clots.

### **12.2 Pharmacodynamics**

Various times (activated clotting time, activated partial thromboplastin time, prothrombin time, whole blood clotting time) are prolonged by full therapeutic doses of heparin; in most cases, they are not measurably affected by low doses of heparin. Bleeding time is usually unaffected by heparin.

### **12.3 Pharmacokinetics**

#### **Absorption**

Heparin is not absorbed through the gastrointestinal tract and therefore administered via parenteral route. Peak plasma concentration and the onset of action are achieved immediately after intravenous administration.

#### **Distribution**

Heparin is highly bound to antithrombin, fibrinogens, globulins, serum proteases and lipoproteins. The volume of distribution is 0.07 L/kg.

#### **Elimination**

##### *Metabolism*

Heparin does not undergo enzymatic degradation.

##### *Excretion*

Heparin is mainly cleared from the circulation by liver and reticuloendothelial cells mediated uptake into extravascular space. Heparin undergoes biphasic clearance, a) rapid saturable clearance (zero order process due to binding to proteins, endothelial cells and macrophages) and b) slower first order elimination. Low doses of heparin are cleared mostly by a saturable, rapid, zero-order process. Slower first order elimination usually occurs with very high doses of heparin and is dependent on renal function. The plasma half-life is dose-dependent, and it ranges from 0.5 to 2 h.

#### **Specific Populations**

##### *Geriatric Patients*

Patients over 60 years of age, following similar doses of heparin, may have higher

plasma levels of heparin and longer activated partial thromboplastin times (APTTs) compared with patients under 60 years of age [see *Use in Specific Populations (8.5)*].

## **Renal and Hepatic Impairment**

The rate of clearance of unfractionated heparin may be decreased in patients with renal or hepatic impairment. Patients with renal or hepatic impairment, following similar doses of heparin may have higher plasma levels of heparin compared with patient with normal renal and hepatic function [see *Warnings and Precautions (5.1)*].

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term studies in animals have been performed to evaluate carcinogenic potential of heparin. Also, no reproduction studies in animals have been performed concerning mutagenesis or impairment of fertility.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Heparin Sodium in Sodium Chloride Injection in VIAFLEX PLUS plastic container is supplied as follows:

<b>Product Description</b>	<b>Size</b>	<b>Code</b>	<b>NDC</b>
1,000 USP Heparin Units in Sodium Chloride Injection	500 mL	2B0953	0338-0431-03
1,000 USP Heparin Units in Sodium Chloride Injection	500 mL	AHB0953U	0338-0424-18
2,000 USP Heparin Units in Sodium Chloride Injection	1,000 mL	2B0944	0338-0433-04
2,000 USP Heparin Units in Sodium Chloride Injection	1,000 mL	AHB0944U	0338-0428-12

Store at 25°C (77°F). Brief exposure up to 40°C (104°F) does not adversely affect the product. Avoid excessive heat.

## **17 PATIENT COUNSELING INFORMATION**

### Hemorrhage

Inform patients that it may take them longer than usual to stop bleeding, that they may bruise and/or bleed more easily when they are treated with heparin, and that they should report any unusual bleeding or bruising to their physician. Hemorrhage can occur at virtually any site in patients receiving heparin. Fatal hemorrhages have occurred [see *Warnings and Precautions (5.1)*].

### Prior to Surgery

Advise patients to inform physicians and dentists that they are receiving heparin before any surgery is scheduled [see *Warnings and Precautions (5.1)*].

### Heparin-Induced Thrombocytopenia

Inform patients of the risk of heparin-induced thrombocytopenia (HIT). HIT may progress to the development of venous and arterial thromboses, a condition known as heparin-induced thrombocytopenia with thrombosis (HITT). HIT and HITT can occur up to several weeks after the discontinuation of heparin therapy [see *Warnings and Precautions (5.2)*].

### Hypersensitivity

Inform patients that generalized hypersensitivity reactions have been reported. Necrosis of the skin has been reported at the site of subcutaneous injection of heparin [see *Warnings and Precautions (5.5)*, *Adverse Reactions (6.1)*].

### Other Medications

Because of the risk of hemorrhage, advise patients to inform their physicians and dentists of all medications they are taking, including non-prescription medications, and before starting any new medication [see *Drug Interactions (7.3)*].

Manufactured by:

**Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

Baxter, Viaflex and Viaflex Plus are trademarks of Baxter International Inc.

07-19-00-3508

ANZ881901209C

**PACKAGE/LABEL PRINCIPAL DISPLAY PANEL**

LOT

EXP

NDC 0338-0431-03

# Heparin

Heparin Sodium in 0.9% Sodium Chloride Injection

**1,000 units per 500 mL**

(2 units / mL)

1

2

3

**500 mL STERILE SINGLE DOSE CONTAINER** EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 4.34 mg DIBASIC SODIUM PHOSPHATE HEPTAHYDRATE USP 400 mcg CITRIC ACID USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L PHOSPHATE (AS HPO<sub>4</sub><sup>=</sup>) 32 mEq/L CITRATE 6 mEq/L OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL DOSAGE *SEE INSERT* **Do NOT ADD SUPPLEMENTARY MEDICATION** STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE **Rx ONLY**

4

## **Baxter**

USA

2B0953

**Container Label**

Container Label

LOT EXP

NDC 0338-0431-03

## **Heparin**

Heparin Sodium in 0.9% Sodium Chloride Injection

**1,000 units per 500 mL**

(2 units/mL)

**500 mL STERILE SINGLE DOSE CONTAINER** EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 4.34 mg DIBASIC SODIUM PHOSPHATE HEPTAHYDRATE USP 400 mcg CITRIC ACID USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L PHOSPHATE (AS HPO<sub>4</sub><sup>=</sup>) 32 mEq/L CITRATE 6 mEq/L OSMOLARITY 322 mOsmol/L

(ACTUAL) USUAL DOSAGE SEE INSERT **DO NOT ADD SUPPLEMENTARY MEDICATION** STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE **RX ONLY**

**Baxter Logo**

USA 2B0953

1  
2

LOT

EXP

NDC 0338-0433-04

**Heparin** 1

Heparin Sodium in 0.9% Sodium Chloride Injection 2

**2,000 units per 1,000 mL**  
(2 units / mL) 3

4

5

**1,000 mL STERILE SINGLE DOSE CONTAINER**

EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 4.34 mg DIBASIC SODIUM PHOSPHATE HEPTAHYDRATE USP 400 mcg CITRIC ACID USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L PHOSPHATE (AS HPO<sub>4</sub><sup>=</sup>) 32 mEq/L CITRATE 6 mEq/L OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL DOSAGE *SEE INSERT* **DO NOT ADD SUPPLEMENTARY MEDICATION** STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE **Rx ONLY** 6  
7  
8

9

**Baxter**

USA

2B0944

**Container Label**

Container Label

LOT EXP



NDC 0338-0433-04

**Heparin**

Heparin Sodium in 0.9% Sodium  
Chloride Injection

2,000 units per 1,000 mL  
(2 units / mL)

**1,000 mL STERILE SINGLE DOSE CONTAINER**

EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN  
SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM  
CHLORIDE USP 4.34 mg DIBASIC SODIUM PHOSPHATE  
HEPTAHYDRATE USP 400 mcg CITRIC ACID USP  
pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE  
154 mEq/L PHOSPHATE (AS  $\text{HPO}_4=$ ) 32 mEq/L CITRATE  
6 mEq/L OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL  
DOSAGE SEE INSERT **DO NOT ADD SUPPLEMENTARY  
MEDICATION** STORE IN MOISTURE BARRIER OVERWRAP AT  
ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE  
**RX ONLY**

**Baxter Logo**

USA 2B0944

07-25-58-614

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9

NDC 0338-0424-18

# Heparin

Heparin Sodium in 0.9% Sodium Chloride Injection

1,000 units per 500 mL  
(2 units / mL)

**500 mL STERILE SINGLE DOSE CONTAINER**

EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 5.80 mg DIBASIC SODIUM PHOSPHATE DODECAHYDRATE USP 400 mcg CITRIC ACID MONOHYDRATE USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L PHOSPHATE (AS HPO<sub>4</sub><sup>=</sup>) 32 mEq/L CITRATE 6 mEq/L OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL DOSAGE SEE INSERT **DO NOT ADD SUPPLEMENTARY MEDICATION** STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE **Rx ONLY**



**Baxter**  
MADE IN AUSTRALIA

Ⓐ AHB0953U  
88-25-02-015A

LOT XXXXX

EXP DATE YYYY/MM

### Container Label

#### Container Label

NDC 0338-0424-18

### Heparin

Heparin Sodium in 0.9% Sodium Chloride Injection

1,000 units per 500 mL  
(2 units / mL)

**500 mL STERILE SINGLE DOSE CONTAINER**

EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 5.80 mg DIBASIC SODIUM PHOSPHATE DODECAHYDRATE USP 400 mcg CITRIC ACID MONOHYDRATE USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L PHOSPHATE (AS HPO<sub>4</sub><sup>=</sup>) 32 mEq/L CITRATE 6 mEq/L OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL DOSAGE SEE INSERT **DO NOT ADD SUPPLEMENTARY MEDICATION** STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE **Rx ONLY**

**Baxter Logo**  
**MADE IN AUSTRALIA**

AHB0953U

A

88-25-02-015A

LOT XXXXX

EXP

DATE

YYYY/MM

1

2

3

4

Barcode

(01)00303380424184

NDC 0338-0428-12

# Heparin

Heparin Sodium in 0.9% Sodium Chloride Injection

2,000 units per 1,000 mL  
(2 units / mL)

GS1 128 Barcode to be placed here NLT 12.7mm x 62.5mm (01)00303380428120

**1,000 mL STERILE SINGLE DOSE CONTAINER**

EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 5.80 mg DIBASIC SODIUM PHOSPHATE DODECAHYDRATE USP 400 mcg CITRIC ACID MONOHYDRATE USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L PHOSPHATE (As HPO<sub>4</sub><sup>-2</sup>) 32 mEq/L CITRATE 6 mEq/L OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL DOSAGE SEE INSERT DO NOT ADD SUPPLEMENTARY MEDICATION STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE (77°F OR 25°C) UNTIL READY TO USE Rx ONLY

**Baxter**  
MADE IN AUSTRALIA

(A) AHB0944U  
88-25-02-016

LOT XXXXX

EXP DATE YYYY/MM

**Container Label**

Container Label

NDC 0338-0428-12

## Heparin

Heparin Sodium in 0.9% Sodium Chloride Injection

2,000 units per 1,000 mL  
(2 units / mL)

**1,000 mL STERILE SINGLE DOSE CONTAINER**

EACH mL CONTAINS 2 USP HEPARIN UNITS HEPARIN SODIUM (PORCINE INTESTINAL MUCOSA) USP 9 mg SODIUM CHLORIDE USP 5.80 mg DIBASIC SODIUM PHOSPHATE DODECAHYDRATE USP 400 mcg CITRIC ACID MONOHYDRATE USP pH 7.0 (6.0 TO 8.0) SODIUM 186 mEq/L CHLORIDE 154 mEq/L

PHOSPHATE (AS HPO4=) 32 mEq/L CITRATE 6 mEq/L  
OSMOLARITY 322 mOsmol/L (ACTUAL) USUAL DOSAGE  
SEE INSERT **DO NOT ADD SUPPLEMENTARY MEDICATION**  
STORE IN MOISTURE BARRIER OVERWRAP AT ROOM TEMPERATURE  
(77°F OR 25°C) UNTIL READY TO USE **RX ONLY**

**Baxter Logo**  
**MADE IN AUSTRALIA**

AHB0944U  
A  
88-25-02-016

LOT XXXXX  
EXP  
DATE  
YYYY/MM

1  
2  
3  
4  
5  
6  
7  
8  
9

GS1 128 Barcode to be placed  
here NLT 12.7mm x 62.5mm  
(01)00303380428120

<b>HEPARIN SODIUM IN SODIUM CHLORIDE</b>			
heparin sodium injection, solution			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-0431
<b>Route of Administration</b>	INTRAVENOUS		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	200 [USP'U] in 100 mL
<b>Inactive Ingredients</b>			

Ingredient Name	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	900 mg in 100 mL
<b>SODIUM PHOSPHATE, DIBASIC, HEPTAHYDRATE</b> (UNII: 70WT22SF4B)	434 mg in 100 mL
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	40 mg in 100 mL
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0431-03	18 in 1 CARTON	04/28/1982	
1		500 mL in 1 BAG; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018609	04/28/1982	

## HEPARIN SODIUM IN SODIUM CHLORIDE

heparin sodium injection, solution

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-0433
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HEPARIN SODIUM</b> (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	200 [USP'U] in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	900 mg in 100 mL
<b>SODIUM PHOSPHATE, DIBASIC, HEPTAHYDRATE</b> (UNII: 70WT22SF4B)	434 mg in 100 mL
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	40 mg in 100 mL
<b>WATER</b> (UNII: 059QF0KO0R)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0433-	12 in 1 CARTON	04/28/1982	

1	04	12 in 1 CARTON	04/20/1982	
1		1000 mL in 1 BAG; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA018609		04/28/1982	

## HEPARIN SODIUM IN SODIUM CHLORIDE

heparin sodium injection, solution

<b>Product Information</b>			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-0424
<b>Route of Administration</b>	INTRAVENOUS		

<b>Active Ingredient/Active Moiety</b>		
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	200 [USP U] in 100 mL

<b>Inactive Ingredients</b>	
<b>Ingredient Name</b>	<b>Strength</b>
SODIUM CHLORIDE (UNII: 451W47IQ8X)	900 mg in 100 mL
SODIUM PHOSPHATE, DIBASIC, HEPTAHYDRATE (UNII: 70WT22SF4B)	580 mg in 100 mL
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	40 mg in 100 mL
WATER (UNII: 059QF0KO0R)	

<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0338-0424-18	1 in 1 CARTON	04/19/2019	
1		500 mL in 1 BAG; Type 0: Not a Combination Product		

<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA018609		04/28/1982	

# HEPARIN SODIUM IN SODIUM CHLORIDE

heparin sodium injection, solution

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-0428
<b>Route of Administration</b>	INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HEPARIN SODIUM (UNII: ZZ45AB24CA) (HEPARIN - UNII:T2410KM04A)	HEPARIN	200 [USP'U] in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	900 mg in 100 mL
SODIUM PHOSPHATE, DIBASIC, HEPTAHYDRATE (UNII: 70WT22SF4B)	580 mg in 100 mL
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	40 mg in 100 mL
WATER (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0428-12	12 in 1 CARTON	04/23/2020	
1		1000 mL in 1 BAG; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018609	04/28/1982	

**Labeler** - Baxter Healthcare Corporation (005083209)

## Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		189326168	ANALYSIS(0338-0431, 0338-0433, 0338-0424, 0338-0428) , LABEL(0338-0431, 0338-0433, 0338-0424, 0338-0428) , MANUFACTURE(0338-0431, 0338-0433, 0338-0424, 0338-0428) , PACK(0338-0431, 0338-0433, 0338-0424, 0338-0428) , STERILIZE(0338-0431, 0338-0433, 0338-0424, 0338-0428)

## Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------



Baxter Healthcare Corporation	059140764	ANALYSIS(0338-0431, 0338-0433, 0338-0424, 0338-0428) , MANUFACTURE(0338-0431, 0338-0433, 0338-0424, 0338-0428) , LABEL(0338-0431, 0338-0433, 0338-0424, 0338-0428) , PACK(0338-0431, 0338-0433, 0338-0424, 0338-0428) , STERILIZE(0338-0431, 0338-0433, 0338-0424, 0338-0428)
-------------------------------	-----------	---

## Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0338-0431, 0338-0433, 0338-0424, 0338-0428)

## Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Pty Ltd		750455891	ANALYSIS(0338-0431, 0338-0433, 0338-0424, 0338-0428) , MANUFACTURE(0338-0431, 0338-0433, 0338-0424, 0338-0428) , LABEL(0338-0431, 0338-0433, 0338-0424, 0338-0428) , PACK(0338-0431, 0338-0433, 0338-0424, 0338-0428) , STERILIZE(0338-0431, 0338-0433, 0338-0424, 0338-0428)

Revised: 1/2022

Baxter Healthcare Corporation