

**LOPERAMIDE HYDROCHLORIDE- loperamide hydrochloride tablet**  
**Family Dollar Services, Inc.**

-----  
**Drug Facts**

**ACTIVE INGREDIENT (IN EACH CAPLET)**

Loperamide HCl USP, 2 mg

**PURPOSE**

Anti-diarrheal

**USE**

Controls symptoms of diarrhea, including Travelers' Diarrhea

**WARNINGS**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Do not use**

If you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease

**Ask a doctor or pharmacist before use if you are**

Taking antibiotics

**When using this product**

- tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- diarrhea lasts for more than 2 days
- symptoms get worse
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding**

Ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**DIRECTIONS**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

#### OTHER INFORMATION

- store between 20° – 25° C (68° – 77° F)
- see side panel for lot number and expiration date
- **TAMPER EVIDENT: THIS PRODUCT PROTECTED WITH SEALED BLISTER UNITS. DO NOT USE IF ANY ARE TORN OR BROKEN.**

#### INACTIVE INGREDIENTS

Anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no.10, FD&C blue no.1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

#### QUESTIONS?

Call **1800-406-7984**

#### PRINCIPAL DISPLAY PANEL

**NDC 55319-554-53**

**FAMILY wellness™**

**COMPARE TO THE ACTIVE INGREDIENT OF IMODIUM® A-D\***

**Anti-Diarrheal**

**Loperamide HCL Tablets USP, 2 mg**

**Controls the Symptoms of Diarrhea**

**12 CAPLETS†**

**Each caplet (†capsule-shaped tablet) contains Loperamide HCl USP, 2 mg**

**DISTRIBUTED BY: FAMILY DOLLAR SERVICES, INC.,**

**5085259/R0411**

5085259



**Drug Facts (continued)**

**Ask a doctor before use if you have**  
 ■ fever ■ mucus in the stool ■ a history of liver disease

**Ask a doctor or pharmacist before use if you are taking antibiotics**

**When using this product**  
 ■ dizziness, drowsiness or lightheadedness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**  
 ■ diarrhea lasts for more than 2 days ■ symptoms get worse  
 ■ you get abdominal swelling or bloating. These may be signs of a serious condition.


**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
 ■ Drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ Find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

**Other information**  
 ■ store between 20° - 25° C (68° - 77° F)  
 ■ see side panel for batch number and expiration date  
 ■ TAMPER EVIDENT: THIS PRODUCT PROTECTED WITH SEALED BLISTER UNITS.  
 DO NOT USE IF ANY ARE TORN OR BROKEN.

5085259



**Drug Facts**

**Active ingredient (in each caplet)**  
 Loperamide HCl USP, 2 mg  
 Anti-diarrheal

**Use**  
 Controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**  
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
 Do not use if you have bloody or black stool

NDC 5319-554-53

**FAMILY wellness™**

COMPARE TO THE ACTIVE INGREDIENT OF IODIUM® A-D\*

**100% SATISFACTION OR YOUR MONEY BACK GUARANTEED**

**Anti-Diarrheal**  
 Loperamide HCl Tablets USP, 2 mg  
 Controls the Symptoms of Diarrhea

**12 CAPLETS\***  
 Each caplet (\*capsule-shaped tablet) contains Loperamide HCl USP, 2 mg

**ACTUAL SIZE**

\*This product is not manufactured or distributed by McNeil-PPC, Inc., distributor of Iodium® A-D. Iodium® is a registered trademark of Johnson & Johnson.

**Keep the carton. It contains important information.**

Expiry Date:

**NON VARNISH**

Batch No.



5085259

**Anti-Diarrheal**  
 Loperamide HCl Tablets USP, 2 mg  
 Controls the Symptoms of Diarrhea



R0411

DISTRIBUTED BY: FAMILY DOLLAR SERVICES, INC.  
 10401 MONROE RD, MATTHEWS, NC 28105 USA  
 Mfr 100% satisfied  
 Return package and unused product within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

SKU 0915632  
V-40153

Questions? call 1-800-406-7984

**Drug Facts (continued)**  
**Inactive Ingredients** anhydrous lactose, croscarmellose sodium, crospovidone, D&C yellow no. 10, FD&C blue no. 1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

## LOPERAMIDE HYDROCHLORIDE

loperamide hydrochloride tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55319-554
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
TRISTEARIN (UNII: P6OCJ2551R)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE	<b>Size</b>	9 mm
<b>Flavor</b>		<b>Imprint Code</b>	123
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-554-53	12 in 1 BLISTER PACK		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA074091	02/01/1993	
------	------------	------------	--

**Labeler** - Family Dollar Services, Inc. (024472631)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

**Establishment**

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		051565745	manufacture(55319-554)

Revised: 9/2012

Family Dollar Services, Inc.