

**GLYCERIN- glycerin suppository suppository**  
**MAJOR® PHARMACEUTICALS**

-----  
**MAJOR® Glycerin**

***Drug Facts***

***Active ingredient***

***(in each suppository)***

Glycerin, USP 2 g

***Purpose***

Hyperosmotic laxative

***Uses***

For relief of occasional constipation. This product usually produces a bowel movement in ¼ to 1 hour.

***Warnings***

**For rectal use only.**

**Do not use**

laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor

**Ask a doctor before use if**

- you have a sudden change in bowel habits lasting more than 2 weeks.
- already used a laxative for more than 1 week.

**When using this product**

rectal discomfort or burning sensation may occur

**Stop using this product and consult a doctor if you have**

- rectal bleeding
- no bowel movement within 1 hour of using this product

This may indicate a serious condition.

**If pregnant or breastfeeding,**

ask a doctor before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

Single Daily dosage: 1 suppository or as directed by a doctor

Adults and children 6 years and over

- tear wrapper prior to immediate use
- remove the secondary wrapper before inserting into the rectum as follows:
  - hold suppository with pointed end up
  - carefully separate tabs with fingernail
  - gently pull apart by pulling tabs down on both sides
    - remove suppository from wrapper
    - insert one suppository well up into the rectum
- Children 2 to 6 years use Children's Glycerin Suppositories
- Children under 2 years of age: ask a doctor

***Other information***

- Only remove overwrap prior to immediate use
- Keep from excessive heat
- Do not use if overwrap is torn

***Inactive ingredients***

polyethylene glycol 400, sodium stearate

***Questions?***

**(855) 879-4368**

**Principal Display Panel**

MAJOR®

Glycerin  
Suppositories

2 g

Laxative

For fast, gentle relief of occasional constipation

Unit Dose

25 Individually Wrapped Adult Suppositories

Retain carton for complete product information

**Drug Facts**

**Active ingredient (in each suppository):** Glycerin, USP 2 g **Purpose:** Hyperosmotic Laxative

**Use:** For relief of occasional constipation. This product usually produces a bowel movement in 1/4 to 1 hour.

**Warnings**

**For rectal use only.**

**Do not use** laxative products when abdominal pain, nausea or vomiting are present unless directed by a doctor

**Ask a doctor before use if:**

- you have a sudden change in bowel habits lasting for more than 2 weeks.
- already used a laxative for more than 1 week.

**When using this product** rectal discomfort or burning sensation may occur

**Stop use and ask a doctor if you have:**

- rectal bleeding
- no bowel movement within 1 hour of using this product

This may indicate a serious condition.

- if pregnant or breastfeeding, ask a doctor before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately (1-800-222-1222).

**Directions**

Single Daily dosage: 1 suppository or as directed by a doctor

- Adults and children 6 years and over:
  - tear wrapper prior to immediate use
  - remove the secondary wrapper before inserting into the rectum as follows:
    - hold suppository with pointed end up
    - carefully separate tabs with fingernail
    - gently pull apart by pulling tabs down on both sides
  - remove suppository from wrapper
  - insert one suppository well up into the rectum
- Children 2 to 6 years use Children's Glycerin Suppositories
- Children under 2 years of age: ask a doctor

**Drug Facts** (continued)

**Other information**

- Only remove overwrap prior to immediate use.
- Keep from excessive heat
- Do not use if overwrap is torn

**Inactive ingredients:** polyethylene glycol 400, sodium stearate

**Questions or comments?** (855) 878-4368

Distributed by:  
**MAJOR® PHARMACEUTICALS**  
Indianapolis, IN 46268  
(800) 616-2471  
www.majorpharmaceuticals.com

\*This product is not manufactured or distributed by C.B. Fleet Company, Inc., owner of the registered trademark Fleet® Glycerin Suppositories

Rev. 05/23 M-182 Re-order No. 701355 FG-46

3 09047 33317 2

**MAJOR®**

NDC 0904-7333-17

Compare to the active ingredient in Fleet® Glycerin Suppositories\*

**Glycerin Suppositories**

**2 g**

**Laxative**

For fast, gentle relief of occasional constipation

**Unit Dose**  
**25 Individually Wrapped Adult Size Suppositories**

Principal Display Panel

MAJOR®

Glycerin Suppositories

2 g

Laxative

For fast, gentle relief of occasional constipation

DO NOT OPEN UNTIL IMMEDIATE USE

1 Individually Wrapped Adult Suppository

**MAJOR<sup>®</sup>**

NDC 0904-7333-01

Compare to the active  
ingredient in Fleet<sup>®</sup>  
Glycerin Suppositories\*

# Glycerin Suppository

**2 g**

**Laxative**

For fast, gentle relief of occasional constipation

**DO NOT OPEN UNTIL IMMEDIATE USE**

**Unit Dose**

**1 Individually Wrapped Adult Size Suppository**

## GLYCERIN

glycerin suppository suppository

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:0904-7333

Route of Administration	RECTAL
-------------------------	--------

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	2 g

**Inactive Ingredients**

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7333-17	25 in 1 CARTON	07/03/2024	
1	NDC:0904-7333-01	1 in 1 POUCH; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	07/03/2024	

**Labeler -** MAJOR® PHARMACEUTICALS (191427277)

Revised: 7/2024

MAJOR® PHARMACEUTICALS