

SPECTOGARD SCOUR-CHEK- spectinomycin solution
Bimeda, Inc.

Liquid

SpectoGard®

Scour-Chek®

(spectinomycin)

ORAL SOLUTION FOR PIG SCOURS

Anti-infective

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only As Directed

Approved by FDA under ANADA # 200-364

INDICATIONS: For use in pigs under four (4) weeks of age for treatment and control of porcine enteric colibacillosis (scours) caused by *E. coli* susceptible to spectinomycin.

EACH mL CONTAINS: Spectinomycin (from spectinomycin dihydrochloride pentahydrate) 50 mg

RESIDUE WARNINGS: This product is only intended for use in pigs under four (4) weeks of age or weighing less than 15 lbs.

Do not administer within 21 days of slaughter.

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

240 mL:

DIRECTIONS FOR USE: The plastic doser supplied in each package with SpectoGard Scour-Chek makes treatment easy. After inserting the doser in the bottle, screw the top on tightly to avoid spilling. Push the piece of clear plastic tubing over the end of the pump. Then press the plunger a few times to fill the pump and the clear plastic tube with medication. To administer SpectoGard Scour-Chek, insert the plastic tube in the pig's mouth and press the plunger to obtain the desired dose.

When not in use remove the plunger from the bottle. Put the clear plastic tube in the neck of the bottle and press the plunger a few times to remove the medication from the pump. Replace the original cap on the bottle and always rinse out the doser with water to prevent sticking.

1000 mL/1 Gallon:

DIRECTIONS FOR USE: The 1000 mL/1 Gallon bottle of SpectoGard Scour-Chek should be used only to refill the 240 mL bottle in the following manner.

1. Remove the cap and doser from the 240 mL bottle.
2. Unscrew the cap from the 1000 mL/1 Gallon bottle.
3. Pour, from the 1000 mL/1 Gallon bottle, the volume desired.
4. Reinsert the doser in the 240 L bottle and screw the cap on tightly.
5. Screw the cap tightly on the 1000 mL/1 Gallon bottle.

INDICATIONS: For use in pigs under four (4) weeks of age for treatment and control of porcine enteric colibacillosis (scours) caused by *E. coli* susceptible to spectinomycin.

EACH mL CONTAINS: Spectinomycin (from spectinomycin dihydrochloride pentahydrate)..... 50 mg

RESIDUE WARNINGS: This product is only intended for use in pigs under four (4) weeks of age or weighing less than 15 lbs.

Do not administer within 21 days of slaughter.

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

SpectoGard® and Scour-Chek® are Registered Trademarks of Bimeda, Inc.

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

N.A. Corp Address: Bimeda, Inc., One Tower Lane,
Oakbrook Terrace, IL 60181



Liquid

SpectoGard®

Scour-Chek®

(spectinomycin)

ORAL SOLUTION FOR PIG SCOURS

Anti-Infective

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

Approved by FDA under ANADA # 200-364



Net Contents:
240 mL (8 fl oz)

DOSE: Pigs under 10 lbs. - 1 pump (1 mL) twice daily.
Pigs over 10 lbs. - 2 pumps (2 mL) twice daily.

Each pump of the plunger delivers 1 mL of solution containing 50 mg of spectinomycin. Treatment may be continued twice daily for 3 to 5 days. If pigs do not improve within 48 hours, re diagnosis is suggested.

DIRECTIONS FOR USE: The plastic doser supplied in each package with SpectoGard Scour-Chek makes treatment easy. After inserting the doser in the bottle, screw the top on tightly to avoid spilling. Push the piece of clear plastic tubing over the end of the pump. Then press the plunger a few times to fill the pump and the clear plastic tube with medication. To administer SpectoGard Scour-Chek, insert the plastic tube in the pig's mouth and press the plunger to obtain the desired dose.

When not in use remove the plunger from the bottle. Put the clear plastic tube in the neck of the bottle and press the plunger a few times to remove the medication from the pump. Replace the original cap on the bottle and always rinse out the doser with water to prevent sticking.

LOT: _____

EXP: _____

1SPE022 8SPE003 Rev. 09/22



8 99855 40015 4

SPECTOGARD SCOUR-CHEK

spectinomycin solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-6002
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Spectinomycin (UNII: 93AK1U6QF) (Spectinomycin - UNII:93AK1U6QF)	Spectinomycin	50 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-6002-5	3785 mL in 1 BOTTLE		
2	NDC:61133-6002-7	1000 mL in 1 BOTTLE		
3	NDC:61133-6002-6	240 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200364	07/09/2004	

Labeler - Bimeda, Inc. (060492923)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Shandong Qilu King-phar Pharmaceutical Co., Ltd.		421524323	api manufacture

Revised: 3/2024

Bimeda, Inc.