

LICE-NIL LICE AND EGGS KILLER- piperonyl butoxide, pyrethrum extract oil
Sujanil Chemo Industries

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Lice-Nil Lice & Eggs Killer Oil

Drug Facts

Active ingredients

Piperonyl butoxide 4%

Pyrethrum extract (equivalent to 0.33% pyrethrins)

Purpose

Lice Treatment

Uses

treats head, pubic (crab), and body lice

Warnings

For external use only

Do not use

- near eyes
- inside nose, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are:

- allergic to ragweed. May cause breathing difficulty or an asthmatic attack.

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel.
- if product gets into eyes, flush with water right away.
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Shake well before using

**Important: Read warnings before use
adults and children 2 years and over:**

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at a time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

Treat

- apply thoroughly to dry hair or other affected area. For head lice, first apply behind ears and to back of neck.
- allow product to remain for minimum 10 minutes.
- wash area thoroughly with warm water and soap or shampoo.
- for head lice, towel dry hair and comb out tangles.

Remove lice and their eggs (nits)

- use a fine-tooth or special lice/nit comb. Remove any remaining nits by hand (using a throw-away glove).
- hair should remain slightly damp while removing nits
- if hair dries during combing, dampen slightly with water
- for head lice, part hair into sections. Do one section at a time starting on top of the head. Longer hair may take 1 to 2 hours.
- lift a 1- to 2- inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- pin back each strand of hair after combing.
- clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back.
- after combing, thoroughly recheck for lice/nits. Repeat combing if necessary.
- check daily for any lice/nits that you missed.
- a second treatment must be done in 7 to 10 days to kill any newly hatched lice.
- if infestation continues, see a doctor for other treatments.
- **children under 2 years: ask a doctor**

Other Information

- shake well before use
- store in cool and dry place
- keep carton for important product information
- see package insert for additional information.

Inactive ingredients

vegetable oils

Questions or comments?

855-KIL-LICE - 855-545-5423

www.sujanil.com or e-mail: sales@sujanil.com

Package Insert

DAILY CHECKING

TO PREVENT REINFESTATION:

- Carefully inspect infested family member ***daily*** for at least 2 weeks. If you see eggs or nits, repeat step 2.
- Check other family members for Head Lice. If anyone in the family is infested, follow Step 1 and Step 2. Repeat Step 3 with any articles or items with which they have come in contact.

Second Application

- Repeat the same procedures within 7 - 10 days for

Step 1: Treat and Step 2: Remove

- There may be some eggs that were not combed out after the first treatment. If infestation continues, see a doctor for other treatments. These eggs will hatch and need to be killed with the second treatment.

Did you know?

Head Lice

- Head lice lay small white eggs (nits) on hair shaft close to the scalp
- Nits are most easily found on back of neck or behind ears
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in hot water (above 54°C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

Pubic (Crab) Lice

- Pubic Lice may be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfestation.
- Pubic Lice are very small and look like brown or grey dots on skin.
- Pubic Lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface.
- Pubic Lice may be presented on the short hairs of groin, thighs, trunk and underarms and occasionally on the beard and mustache.
- Pubic Lice infested underwear can be disinfected by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes.

Body Lice

- Body Lice and their eggs (nits) are generally found in the seams of clothing particularly in waistline and armpit area.
- Body Lice feed on skin then return to clothing to lay their eggs.
- Body Lice infested clothing can be disinfected by machine washing in hot water (above 54°C (130°F)), then using the hottest dryer cycle for at least 20 minutes.
- Body Lice nits can remain dormant for up to 30 days and therefore clothing should not be sealed in a plastic bag

- *Save this package insert for future reference.*

STEP 1 TREAT

Lice-Nil Lice & Eggs Killer Oil

Steps to Success

with 1-2-3 Lice Control System

Complete Lice Control for your family and home.

SHAKE WELL BEFORE USING

DIRECTIONS: IMPORTANT: READ WARNINGS BEFORE USING. A SECOND TREATMENT MUST BE DONE IN 7-10 DAYS TO KILL ANY NEWLY HATCHED LICE.

Warnings

For external use only

Do not use:

- near eyes,
- inside nose, mouth or vagina,
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas

Ask a doctor before use if you are

- allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets in eyes, flush with water right away
- scalp itching or redness may occur

Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs
- Keep out of reach of children. if swallowed, get medical help or contact a Poison Control Center right away

See Usage Guidelines below to estimate how much product you will need.

Usage Guidelines

Hair Length	Approximate Amount for 1 Adult/Child:
Short (ear length or shorter)	First Application : 1 oz - 2 oz
	Second Application : 1 oz - 2 oz
	Total : 2 oz - 4 oz
Medium (shoulder length)	First Application : 2 oz - 3 oz
	Second Application : 2 oz - 3 oz
	Total : 4 oz - 6 oz
Long (past shoulder length)	First Application : 3 oz - 4 oz
	Second Application : 3 oz - 4 oz
	Total : 6 oz - 8 oz

Inspect

- check each household member with a magnifying glass in bright light for lice/nits (eggs)
- look for tiny nits near scalp, beginning at back of neck and behind ears
- examine small sections of hair at time
- unlike dandruff which moves when touched, nits stick to the hair
- if either lice or nits are found, treat with this product

NOTE:

- Use towels to protect child's eyes from treatment. Ask the child to keep eyes tightly closed.
- Apply '**Lice-Nil**' Lice & Eggs Killer Oil thoroughly to **DRY HAIR** or other affected area.
- Wetting the hair dilutes the treatment making it less effective.
- Also lice can hold their breath when immersed in water making it difficult for the active ingredient to penetrate the lice
- For head lice, thoroughly massage oil into scalp, behind the ears and to the back of the neck. Lice can crawl up and down the shaft of the hair very quickly so it is important to apply the treatment from the roots to the ends of the hair.
- Allow product to remain for minimum 10 minutes.
- Wash area thoroughly with warm water and soap or shampoo
- For head lice, towel dry hair and comb out tangles
- Proceed to Step 2.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.

The 'Licel-Nil' comb is proven 100% effective as demonstrated in laboratory studies performed by trained testers, individual results may vary.

STEP 2 REMOVE

NOTE:

- A fine - toothed comb or special lice / nit removing comb (included) must be used to remove lice, eggs, and nits from hair.
- **Before starting have on hand:**
 - extra towels
 - bobby pins or hair clips
 - facial tissues or paper towels
 - regular comb for parting and detangling hair
 - throw-away gloves
- Find a distraction for your child (e.g. books, TV, game)
- Comb out hair with a regular wide-tooth comb to take out snarls and tangles.
- When combing, use the fine-toothed lice/nit comb provided and make sure the teeth are facing you.
- Part hair into section. Do one section at a time. Starting on top of the head. Longer hair may take 1 to 2 hours.
- Lift a 1- to 2- inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp.
- Always comb away from the head.
- Pin back each strand of hair after combing.
- Wipe eggs and nits from the teeth of the comb with a tissue or paper towel often.
- Remove any remaining nits by hand (using a throw-away glove)
- Discard all the used tissue or cloths into bag, seal the bag and dispose of immediately to prevent lice from coming back.
- After combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- After you have combed and pinned the entire head, remove all bobby pins or hair clips and

thoroughly recheck for lice/nits. Repeat combing if necessary.

- Rinse hair thoroughly with warm water, If desired, you may also wash with regular shampoo.
- Check daily for any lice/nits that you missed.
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice.
- If infestation continues, see a doctor for other treatments.
- ***Children under 2 years:*** Ask a doctor

STEP 3 CONTROL

- Head lice lay small white eggs (nits) on the hair shaft close to the scalp
- Nits are most easily found on the back of the neck or behind the ears
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using the hottest dryer cycle for at least 20 minutes.
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in to water (above 54°C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

CHECKLIST

- Home and auto upholstery
- Pillows, pillowcases, mattresses
- Bed Linen
- Towels
- Stuffed animals toys
- Hats
- Headphones
- Headbands
- Bike/sports helmets
- Coats, gloves, scarves, earmuffs
- Play mats
- Clothing
- Barettes/Hair ornaments
- Hair rollers
- Combs and brushes
- Baseball caps

Principal Display Panel

Drug Facts

Active Ingredients	Purpose
Piperonyl butoxide 4%.....Lice treatment Pyrethrum extract.....Lice treatment (equivalent to 0.33% pyrethrins)	

Uses: Treats head lice/nits, pubic lice (crab) and body lice

Warnings

For external use only

- Do not use:**
- Near eyes
 - Inside nose, mouth or vagina
 - On lice in eyebrows or eyelashes. See a doctor if lice are present in these areas

Ask a doctor before use if you are:

- Allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product:

- Keep eyes tightly closed and protect eyes with a wash cloth or towel.
- If product gets in eyes, flush with water right away
- Scalp itching or redness may occur

Stop use and ask a doctor if:

- Breathing difficulty occurs
- Eye irritation occurs
- Skin or scalp irritation continues or infection occurs

Keep out of reach of children. If swallowed, get medical help or contact a poison Control Center right away.

Directions

- **Important:** Read warnings before use
- For Adults and Children 2 years and older

Inspect

- Check each household member with a magnifying glass in bright light for lice/nits (eggs)
- Look for tiny nits near scalp, beginning at back of neck and behind ears
- Examine small sections of hair at a time
- Unlike dandruff which moves when touched, nits stick to the hair
- If either lice or nits are found, treat with this product

Treat

- Apply thoroughly to dry hair or other affected area. For head lice, first apply on hair behind ears and back of neck, then continue to apply to the remainder of hair
- Leave product on hair for a minimum of 10 minutes
- Wash area thoroughly with warm water and shampoo or soap
- For head lice, towel dry hair and comb out tangles

Remove lice and their eggs (nits)

- Using a fine-tooth comb or special lice/nit comb, remove any remaining nits by hand (using a throw-away glove)
- Hair should remain slightly damp while removing nits
- If hair dries during combing, dampen slightly with water

Drug Facts (continued)

- For head lice, part hair into sections and inspect one section at a time, starting at the top of head. Longer hair may take 1-2 hours.
- Lift a 1-2-inch wide strand of hair. Place comb with a firm, even motion away from scalp.
- Pin back each strand of hair after combing
- Clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back
- After combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- Check daily for any lice/nits that may have been missed
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice
- If infestation continues, see a doctor for other treatments
- **Children under 2 years:** Consult with a doctor or physician

Other information

- Shake well before use.
- Store in cool and dry place.
- Keep carton for important product information.
- See package insert for a additional information.

Inactive ingredients:

Coconut and Sesame Oil

Questions or comments?

855-KIL-LICE - 855-545-5423
www.sujanil.com or e-mail: sales@sujanil.com

"Lice-Nil" Lice Treatment Oil will kill lice completely without leaving a chemical residue. The amount of lice treatment oil needed will vary depending on hair length. See below for guidelines:

Hair Length	Approximate Amount for 1 Adult/Child:
Short (ear length or shorter)	First Application : 1 oz - 2 oz
	Second Application : 1 oz - 2 oz
	Total : 2 oz - 4 oz
Medium (shoulder length)	First Application : 2 oz - 3 oz
	Second Application : 2 oz - 3 oz
	Total : 4 oz - 6 oz
Long (past shoulder length)	First Application : 3 oz - 4 oz
	Second Application : 3 oz - 4 oz
	Total : 6 oz - 8 oz

The "Lice-Nil" comb is proven 100% effective as demonstrated in laboratory studies performed by trained testers, individual results may vary.

20 Minute Application Kills Lice & Eggs Completely!

NDC No.: 76441-903-97



Proven & tested for more than 4 decades
with **Magnifier** included
NET WT : 4 FL OZ (118 ml)

A PRODUCT OF INDIA

'Lice-Nil' Lice & Eggs Eliminator Oil Lice Treatment

- Nothing proven more effective
- Leaves no chemical residue behind
- Easy to use oil

Lice-Nil 1-2-3 LICE ELIMINATION SYSTEM

Step 1 Head Lice

- Head lice lay small, white egg (nits) on the hair shaft near the scalp. Nits are most easily found on the back of the neck or behind the ears.
- Disinfect hats, hair ribbons, clothing, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then dry using the hottest dryer cycle for at least 20 minutes. Items that cannot be washed (pillows, stuffed toys, etc) should be dry-cleaned or sealed in a plastic bag for 4 weeks. After 4 weeks, remove them from the plastic bag outdoors and shake rigorously before using again.
- Items that cannot be washed, dry-cleaned or stored may be sprayed with a product designed for purpose.
- Soak all combs and brushes in hot water (above 54°C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture and car seats that may have been used by affected people.

Step 2 Pubic (Crab) Lice

- Pubic lice may be transmitted by sexual contact
- Sexual partners should be treated simultaneously to avoid reinfestation
- Pubic lice usually cause intense itching and lay small white eggs (nits) on the hair shaft, near the skin surface
- Pubic lice may be present on the groin, thighs, trunk, underarms and occasionally on the beard or mustache
- Disinfect underwear by machine washing in hot water (above 54°C (130°F)), then use hottest dryer cycle for at least 20 minutes

Step 3 Body Lice

- Body lice and their eggs (nits) are generally found in the seams of clothing, particularly in the waistline and armpit area
- Body lice feed on skin then return to clothing to lay their eggs
- Disinfect clothing by machine washing in hot water (above 54°C (130°F)), then use hottest dryer cycle for at least 20 minutes
- Do not seal clothing in plastic bag because nits can remain dormant for up to 30 days

Manufactured By:

sujanil chemo industries
425/27, T.M.V. Colony, Pune 411 037 (INDIA)

Imported by:
Global Tradelinks
4337, Austonway, Palm Harbour,
FL 34685, USA
Tel.: 1-727-359-1173, Fax: 1-813-886-3881,
E-mail: jay@sujanil.com

DAILY CHECKING



TO PREVENT REINFESTATION :

- Carefully inspect affected family member daily for at least 2 weeks. If you see eggs or nits, repeat step 2
- Check other family members for Head Lice. If anyone in the family is infested, follow Step 1 and Step 2. Repeat Step 3 with any articles or items which they have come into contact

Second Application

- Repeat the same process within 7-10 days for

Step 1 : Treat and Step 2 : Remove

There may be some eggs that were not combed out after the first treatment. If infestation continues, see a doctor for other treatments.

These eggs will hatch and need to be killed with the second treatment.

Did you know?

Head Lice

- Head lice lay small white eggs (nits) on hair shaft close to the scalp
- Nits are most easily found on back of neck or behind ears
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out

very hard before using again

- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in hot water (above 54°C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people

Pubic (Crab) Lice

- Pubic Lice may be transmitted by sexual contact. Sexual partners should be treated simultaneously to avoid reinfestation
- Pubic Lice are very small and look like brown or grey dots on skin
- Pubic Lice usually cause intense itching and lay small white eggs (nits) on the hair shaft generally close to the skin surface
- Pubic Lice may be present on the short hairs of groin, thighs, trunk and underarms and occasionally on the beard and mustache
- Pubic Lice infested underwear can be disinfected by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes

Body Lice

- Body Lice and their eggs (nits) are generally found in the seams of clothing particularly in the waistline and armpit area
- Body Lice feed on skin then return to clothing to lay their eggs
- Body Lice infested clothing can be disinfected by machine washing in hot water (above 54°C (130°F)), then using hottest dryer cycle for at least 20 minutes
- Body Lice nits can remain dormant for up to 30 days and therefore clothing should not be sealed in a plastic bag
- **Save this package insert for future reference.**



Maximum Strength
Lice-Nil
LICE AND EGGS ELIMINATOR OIL

Questions or comments?

855-KIL-LICE - 855-545-5423
www.sujanil.com or e-mail: sales@sujanil.com

Manufactured by:

sujanil chemo industries
425/27, T.M.V. Colony, Pune 411 037 (INDIA)

Imported by:

Global Tradelinks
4337, Austonway, Palm Harbour,
FL 34685, USA
Tel.: 1-727-359-1173, Fax: 1-813-886-3881,
E-mail: jay@sujanil.com

One 20 Minute Application Kills Lice & Eggs Completely!

NDC No.: 76441-903-97



Proven & tested for more than 4 decades
with **Magnifier** included
NET WT : 4 FL OZ (118 ml)

A PRODUCT OF INDIA

STEP 1 TREAT



Steps for Success with the Lice-Nil 1-2-3 Lice Control System

Complete Lice Control for your family and home.

SHAKE WELL BEFORE USING

Warnings: For external use only

Do not use:

- Near eyes
- Inside nose, mouth or vagina
- On lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

Ask a doctor before use if you are:

- Allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product:

- Keep eyes tightly closed and protect eyes with a washcloth or towel

- If product gets in eyes, flush with water right away
- Scalp itching or redness may occur

Stop use and ask a doctor if:

- Breathing difficulty occurs
- Eye irritation occurs
- Skin or scalp irritation continues or infection occurs
- Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away

DIRECTIONS:

IMPORTANT: READ WARNINGS BEFORE USING. A SECOND TREATMENT MUST BE DONE IN 7-10 DAYS TO KILL ANY NEWLY HATCHED LICE.

See Usage Guidelines below to estimate how much product you will need.

Inspect

- Check each household member with a magnifying glass in bright light for lice/nits (eggs)
- Look for tiny nits near scalp, beginning at back of neck and behind ears
- Examine small sections of hair at a time
- Unlike dandruff which moves when touched, nits stick to the hair
- If either lice or nits are found, treat with this product

NOTE:

- Use towels to protect child's eyes from treatment. Ask the child to keep eyes tightly closed
- Apply 'Lice-Nil' Lice Treatment Oil thoroughly to **DRY HAIR** or other affected area
- Wetting the hair dilutes the treatment making it less effective
- Also, lice can hold their breath when immersed in water, making it difficult for the active ingredient to penetrate the lice
- For head lice, thoroughly massage oil into scalp, behind the ears and to the back of the neck. Lice can crawl up and down the shaft of the hair very quickly so it is important to apply the treatment from the roots to the ends of the hair

- Allow product to remain on hair for a minimum 20 minutes
- Wash area thoroughly with warm water and shampoo or soap
- For head lice, towel dry hair and comb out tangles
- Proceed to Step 2
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice

Usage Guidelines

Hair Length	Approximate Amount for 1 Adult/Child
Short (ear length or shorter)	First Application : 1 oz - 2 oz
	Second Application : 1 oz - 2 oz
	Total : 2 oz - 4 oz
Medium (shoulder length)	First Application : 2 oz - 3 oz
	Second Application : 2 oz - 3 oz
	Total : 4 oz - 6 oz
Long (past shoulder length)	First Application : 3 oz - 4 oz
	Second Application : 3 oz - 4 oz
	Total : 6 oz - 8 oz

The 'Lice-Nil' comb is proven 100% effective as demonstrated in laboratory studies performed by trained testers, individual results may vary.

STEP 2 REMOVE



NOTE:

- A fine-tooth comb or a special lice/nit removing comb (included) must be used to remove lice, eggs and nits from hair
- Before starting, please have these items available:
 - extra towels
 - bobby pins or hair clips
 - facial tissues or paper towels
 - regular comb for parting and detangling hair
 - throw-away gloves
- Find a distraction for your child (ex. books, TV or games)
- Comb out hair with a regular wide-tooth comb to take out snags and tangles.

- When combing, use the fine-toothed lice/nit comb provided and make sure the teeth are facing you
- Part hair into section. Do one section at a time. Starting on top of the head. Longer hair may take 1-2 hours
- Lift a 1-2 inch wide strand of hair. Place comb as close to scalp as possible and comb with a firm, even motion away from scalp

- Always comb away from the head
- Pin back each strand of hair after combing
- Wipe eggs and nits from the teeth of the comb with a tissue or paper towel often
- Remove any remaining nits by hand (using a throw-away glove)
- Discard all the used tissues or cloths into bag, seal the bag and dispose of immediately to prevent lice from coming back
- After combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- After you have combed and pinned the entire head, remove all bobby pins or hair clips and thoroughly recheck for lice/nits. Repeat combing if necessary
- Rinse hair thoroughly with warm water. If desired, you may also wash with regular shampoo
- Check daily for any lice/nits that may have been missed
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice
- If infestation continues, see a doctor for other treatments
- Children under 2 years: Consult with a doctor or physician



STEP 3 CONTROL

- Head lice lay small white eggs (nits) on the hair shaft close to the scalp
- Nits are most easily found on the back of the neck or behind the ears
- Disinfect hats, hair ribbons, scarves, coats, towels and bed linens by machine washing in hot water (above 54° C (130°F)), then using the hottest dryer cycle for at least 20 minutes
- Items that cannot be washed (bedspreads, blankets, pillows, stuffed toys, etc.) should be dry-cleaned or sealed in a plastic bag for 4 weeks, then removed outdoors and shaken out very hard before using again
- Items that cannot be washed, dry-cleaned, or stored may be sprayed with a product designed for this purpose
- Soak all combs and brushes in water (above 54° C (130°F)) for at least 10 minutes
- Vacuum all carpets, mattresses, upholstered furniture, and car seats that may have been used by affected people.

CHECKLIST

- Home and auto upholstery
- Pillows, pillowcases, mattresses
- Bed Linen
- Towels
- Stuffed Animals and Toys
- Hats
- Headphones
- Headbands
- Bike/sports helmets
- Coats, gloves, scarves, earmuffs
- Play mats
- Clothing
- Barettes/Hair Accessories
- Hair rollers
- Combs and brushes
- Baseball caps

Active Ingredients

Purpose

Piperonyl butoxide 4%.....Lice treatment
Pyrethrum extract
(equivalent to 0.33% pyrethrins). Lice treatment

Uses: Treats head lice/nits, pubic lice (crab) and body lice

Inactive Ingredients: Coconut and Sesame Oil

Warning: For external use only

Do not use:

- Near eyes
- Inside nose, mouth or vagina
- On lice in eyebrows or eyelashes. See a doctor if lice are present in these areas

Ask a doctor before use if you are:

- Allergic to ragweed. May cause breathing difficulty or an asthmatic attack

When using this product:

- Keep eyes tightly closed and protect eyes with a wash cloth or towel
- If product gets in eyes, flush with water right away
- Scalp itching or redness may occur

Stop and ask a doctor if:

- Breathing difficulty occurs
- Eye irritation occurs
- Skin or scalp irritation continues or infection occurs

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Other Information:

- Shake well before use
- Store in cool and dry place
- Keep carton for important product information
- See package insert for additional information

Questions or comments?
855-KIL-LICE - 855-545-5423

www.sujanil.com or e-mail: sales@sujanil.com

20 Minute Application Kills Lice & Eggs Completely!

NDC No.: 76441-903-97



**100% NATURAL
ACTIVES**
NO HARMFUL CHEMICALS

Maximum Strength
Lice-Nil[®]
LICE AND EGGS ELIMINATOR OIL



**Proven & tested
for more than
4 decades**

NET WT : 4 FL OZ (118 ml)

**Comb
with Magnifier
included**

Directions :

- **Important:** Read warnings before use
- For Adults and Children 2 years and older

Inspect

- Check each household member with a magnifying glass in bright light for lice/nits (eggs)
- Look for tiny nits near scalp, beginning at back of neck and behind ears
- Examine small sections of hair at a time
- Unlike dandruff which moves when touched, nits stick to the hair
- If either lice or nits are found, treat with this product

Treat

- Apply thoroughly to dry hair or other affected area. For head lice, first apply on hair behind ears and back of neck, then continue to apply to the remainder of hair
- Leave product on hair for a minimum of 10 minutes
- Wash area thoroughly with warm water and shampoo or soap
- For head lice, towel dry hair and comb out tangles

Remove lice and their eggs (nits)

- Using a fine-tooth comb or special lice/nit comb, remove any remaining nits by hand (using a throw-away glove)
- Hair should remain slightly damp while removing nits
- If hair dries during combing, dampen slightly with water
- For head lice, part hair into sections and inspect one section at a time, starting at the top of head. Longer hair may take 1-2 hours
- Lift a 1-2-inch wide strand of hair. Place comb with a firm, even motion away from scalp
- Pin back each strand of hair after combing
- Clean comb often. Wipe nits away with tissue and discard in a plastic bag. Seal bag and discard to prevent lice from coming back
- After combing, thoroughly recheck for lice/nits. Repeat combing if necessary
- Check daily for any lice/nits that may have been missed
- A second treatment must be done in 7 to 10 days to kill any newly hatched lice
- If infestation continues, see a doctor for other treatments
- **Children under 2 years:** Consult with a doctor or physician

Batch No. :

Mfg. Date :

Expiry Date :

Manufactured By:

sujanil chem industries

425/27, T.M.V. Colony, Pune 411 037 (INDIA)

Imported by:

Global Tradelinks

4337, Austonway, Palm Harbour, FL 34685, USA

Tel. : 1-727-359-1173, Fax : 1-813-886-3881,

E-mail : jay@sujanil.com

A PRODUCT OF INDIA

LICE-NIL LICE AND EGGS KILLER

piperonyl butoxide, pyrethrum extract oil

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:76441-903
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PIPERONYL BUTOXIDE (UNII: LWK91TU9AH) (PIPERONYL BUTOXIDE - UNII:LWK91TU9AH)	PIPERONYL BUTOXIDE	4 g in 100 mL
PYRETHRUM EXTRACT (UNII: ZUM06L90GV) (PYRETHRUM EXTRACT - UNII:ZUM06L90GV)	PYRETHRUM EXTRACT	0.33 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SESAME OIL (UNII: QX10HYY4QV)	
COCONUT OIL (UNII: Q9L0O73W7L)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76441-903-97	1 in 1 BOX		
1		118 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part358G	06/18/2013	

Labeler - Sujanal Chemo Industries (650280816)

Establishment

Name	Address	ID/FEI	Business Operations
Sujanal Chemo Industries		915896189	manufacture(76441-903)

Revised: 11/2014

Sujanal Chemo Industries