

**EQUATE DAYTIME VAPOR ICE COLD AND FLU- acetaminophen,  
dextromethorphan hydrobromide, guaifenesin, phenylephrine  
hydrochloride solution  
Wal-Mart Stores Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Wal-Mart Day Time Vapor Ice® Cold & Flu Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- sinus congestion and pressure
- nasal congestion
- minor aches and pains
- headache
- sore throat
- fever
- cough due to minor throat and bronchial irritation
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

**do not use more than directed**

## Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

## Other information

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

## Questions?

**1-888-287-1915**

## Package/Label Principal Display Panel

equate™

Compare to Vicks® DayQuil® Severe+ VapoCOOL™ active ingredients

DAYTIME

SEVERE

VAPOR ICE® Cold & Flu

Minor Aches & Pains, Fever, Nasal Congestion & Sinus Pressure, Cough, Chest Congestion

- Acetaminophen – Pain reliever/Fever reducer
- Phenylephrine HCl – Nasal decongestant
- Dextromethorphan HBr – Cough suppressant
- Guaifenesin – Expectorant

12 FL OZ (355mL)

10% ALCOHOL

NDC 45035-019-90

**equate™**

Compare to  
Vicodin DayQuil®  
Severe+ VapoCOOL™  
active ingredients\*



**SEVERE**

**VAPOR ICE®**

# Cold & Flu

Minor Aches & Pains, Fever, Nasal Congestion & Sinus Pressure, Cough, Chest Congestion

- **Acetaminophen** - Pain reliever/Fever reducer
- **Phenylephrine HCl** - Nasal decongestant
- **Dextromethorphan HBr** - Cough suppressant
- **Guaifenesin** - Expectorant

**12 FL OZ (355mL)**

**10% ALCOHOL**

: 94M40 2E F2

## PARENTS:

Learn about teen medicine abuse

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

164152 \*

### Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

**Uses:** temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**PEEL BACK AT CORNER FOR MORE INFORMATION**

: 94M40 2E B1



### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

ADHESIVE AREA  
NO VARNISH • NO TYPE

### Drug Facts (continued)

When using this product do not use more than directed

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.



If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
---------------------------------	-------------------

ADHESIVE AREA  
NO VARNISH • NO TYPE



## EQUATE DAYTIME VAPOR ICE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-819
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	

<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-819-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/03/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/03/2019	

**Labeler** - Wal-Mart Stores Inc (051957769)

Revised: 10/2022

Wal-Mart Stores Inc