

DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl
Rite Aid Corporation

Drug Facts

Active ingredients for Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients for Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - headache
 - minor aches and pains
 - fever
 - sore throat

NIGHTTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing

Warnings

DAYTIME

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using these products

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

NIGHTTIME

Liver warning: This product contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

Do not use

DAYTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

NIGHTTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

taking the blood thinning drug warfarin

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product,

DAYTIME

do not exceed recommended dosage

NIGHTTIME

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts. These could be a signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

DAYTIME

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over; take 2 softgels with water every 4 hours.
- children under 12 years: do not use

NIGHTTIME

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours

- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

DAYTIME

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

NIGHTTIME

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan*, sorbitol, white ink

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps®

MULTI-SYMPTOM COLD & FLU RELIEF

DAYTIME

ACETAMINOPHEN 325 mg

DEXTROMETHORPHAN HBr 10 mg

PHENYLEPHRINE HCL 5 mg

PAIN RELIEVER / FEVER REDUCER

COUGH SUPPRESSANT

NASAL DECONGESTANT

NON-DROWSY

Temporarily relieves aches, fever & sore throat

Cough • Nasal congestion

Antihistamine-free

Alcohol-free

DAYTIME

SOFTGELS**

LIQUID FILLED CAPSULES

NIGHTTIME

ACETAMINOPHEN 325 mg

DEXTROMETHORPHAN HBr 15 mg

DOXYLAMINE SUCCINATE 6.25 mg

PAIN RELIEVER / FEVER REDUCER

COUGH SUPPRESSANT

ANTIHISTAMINE

Temporarily relieves aches, fever & sore throat

Cough,

Runny nose & sneezing

Alcohol-free

NIGHTTIME

SOFTGELS**

**LIQUID FILLED CAPSULES

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil, NyQuil®, and LiquiCaps® are registered trademarks of the Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DISTRIBUTED BY:

RITE AID

30 HUNTER LANE,

CAMP HILL, PA 17011

www.riteaid.com

Product Label

MULTI-SYMP TOM COLD & FLU RELIEF

● DAYTIME

ACETAMINOPHEN 325 mg
DEXTROMETHORPHAN HBr 10 mg
PHENYLEPHRINE HCl 5 mg
 PAIN RELIEVER / FEVER REDUCER
 COUGH SUPPRESSANT
 NASAL DECONGESTANT

NON-DROWSY

Temporarily relieves aches, fever & sore throat

Cough • Nasal congestion
 Antihistamine-free
 Alcohol-free

ACTUAL SIZE



When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

☾ NIGHTTIME

ACETAMINOPHEN 325 mg
DEXTROMETHORPHAN HBr 15 mg
DOXYLAMINE SUCCINATE 6.25 mg
 PAIN RELIEVER / FEVER REDUCER
 COUGH SUPPRESSANT
 ANTIHISTAMINE

Temporarily relieves aches, fever & sore throat

Cough
 Runny nose & sneezing
 Alcohol-free

ACTUAL SIZE

16 DAYTIME SOFTGELS**
8 NIGHTTIME SOFTGELS**
 **LIQUID FILLED CAPSULES

DISTRIBUTED BY:
 RITE AID
 30 HUNTER LANE,
 CAMP HILL, PA 17011
 www.riteaid.com



**SATISFACTION
 GUARANTEE**

If you're not satisfied, we'll happily refund your money.

PARENTS:
 Learn more here: www.riteaid.com
 www.StayHealthyUseItUp

PLD-C570A
 FC0005819

Lot No.:

Exp. Date:

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Daytime Cold & Flu Softgel

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product, do not exceed recommended dosage.
Stop use and ask a doctor if ■ nervousness, dizziness, or sleepiness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Nighttime Cold & Flu Softgel

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. ■ taking sedatives or tranquilizers ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery
Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
Keep out of reach of children. Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts
Active ingredients (in each softgel)
 Acetaminophen 325 mg
 Dextromethorphan HBr 10 mg
 Phenylephrine HCl 5 mg
Purposes
 Pain reliever/fever reducer
 Cough suppressant
 Nasal decongestant
Uses ■ temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ headache ■ minor aches and pains ■ fever ■ sore throat
Warnings ■ Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts
Active ingredients (in each softgel)
 Acetaminophen 325 mg
 Dextromethorphan HBr 15 mg
 Doxylamine succinate 6.25 mg
Purposes
 Pain reliever/fever reducer
 Cough suppressant
 Antihistamine
Uses ■ temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing
Warnings ■ Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, MyQuil®, and LiquiCaps® are registered trademarks of The Procter and Gamble Company.

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST
Inactive ingredients butylated hydroxyanisole, butylated glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink	Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink
Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat	Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat
Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use	Directions ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use
Drug Facts (continued) Daytime Cold & Flu Softgel	Drug Facts (continued) Nighttime Cold & Flu Softgel

RITE AID Multi-Symptom Cold & Flu Relief

DAYTIME NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0851
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0851-4	1 in 1 KIT; Type 0: Not a Combination Product	08/30/2019	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	8 BLISTER PACK	8
Part 2	16 BLISTER PACK	16

Part 1 of 2

NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:11822-3335
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	P30
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/30/2019	

Part 2 of 2

DAYTIME COLD AND FLU NON DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Item Code (Source) NDC:11822-3391

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O921CV9RU)	
SORBITOL (UNII: 506T60A25R)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	P19
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M012	08/30/2019	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/30/2019	

Labeler - Rite Aid Corporation (014578892)

Revised: 5/2024

Rite Aid Corporation