

ACID REDUCER- ranitidine tablet
Select Brand

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

select brand[®]

NDC 15127-340-24

Maximum Strength

Ranitidine Tablets, USP 150 mg

Acid Reducer


Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

†Compare to the active ingredient of Zantac 150[®]


24 Tablets

Distributed by: SELECT BRAND[®] DISTRIBUTORS

5093019/0112



6106905



5093019

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Questions? call 1-800-406-7984

Inactive ingredients
 titanium dioxide
 microcrystalline cellulose, polyethylene glycol, talc, hypromellose, iron oxide red, magnesium stearate, colloidal silicon dioxide, croscarmellose sodium.

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.
 ■ store at 20° - 25° C (68° - 77° F)
 ■ avoid excessive heat or humidity
 ■ this product is sodium and sugar free

Directions
 ■ adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water
 ■ to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
 ■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)
 ■ children under 12 years: ask a doctor

Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days

Ask a doctor before use if you have
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Warnings
 Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
 Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
 ■ with other acid reducers
 ■ if you have kidney disease, except under the advice and supervision of a doctor

Drug Facts (continued)
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition
 ■ heartburn with lightheadedness, sweating or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Drug Facts
Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg) Acid reducer
Purpose Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg) Acid reducer

Uses
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages
 ■ relieves heartburn associated with acid indigestion and sour stomach

select brand
 the lower price name brand

Maximum Strength
 Ranitidine Tablets, USP 150 mg
Acid Reducer

24
Tablets

select brand
 the lower price name brand

Maximum Strength
Ranitidine Tablets, USP 150 mg
Acid Reducer

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

† Compare to the active ingredient of Zantac 150®



24
Tablets

Expiration Date:

Batch No.:

NON VARNISH

† This product is not manufactured or distributed by



ACID REDUCER

ranitidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:15127-340
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	9R
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:15127-340-24	24 in 1 CARTON		
1		1 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200536	03/30/2012	

Labeler - Select Brand (043562370)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(15127-340)

Revised: 10/2012

Select Brand