


**ORPHENADRINE CITRATE- orphenadrine citrate tablet, extended release**  
**Preferred Pharmaceuticals, Inc**

-----  
**Orphenadrine Citrate Extended-Release Tablets**

**DESCRIPTION**

Orphenadrine citrate is the citrate salt of orphenadrine. It occurs as a white, crystalline powder having a bitter taste. It is practically odorless; sparingly soluble in water, slightly soluble in alcohol. The chemical name of orphenadrine citrate is ( $\pm$ )-*N,N*-Dimethyl-2-[(*o*-methyl- $\alpha$ -phenylbenzyl)oxy]ethylamine citrate (1:1) having molecular formula  $C_{18}H_{23}NO \cdot C_6H_8O_7$  and molecular weight of 461.51. It has the following structural formula:



Each tablet for oral administration contains 100 mg orphenadrine citrate.

Each Orphenadrine citrate extended-release tablet contains the following inactive ingredients: hydroxypropyl methylcellulose, lactose monohydrate and magnesium stearate.

**CLINICAL PHARMACOLOGY**

The mode of therapeutic action has not been clearly identified, but may be related to its analgesic properties. Orphenadrine citrate does not directly relax tense muscles in man. Orphenadrine citrate also possesses anti-cholinergic actions.

**INDICATIONS AND USAGE**

Orphenadrine citrate extended-release tablets are indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute painful musculoskeletal conditions.

**CONTRAINDICATIONS**

Orphenadrine citrate extended-release tablets are contraindicated in patients with glaucoma, pyloric or duodenal obstruction, stenosing peptic ulcers, prostatic hypertrophy or obstruction of the bladder neck, cardio-spasm (mega-esophagus) and myasthenia gravis.

Orphenadrine citrate tablets are contraindicated in patients who have demonstrated a previous hypersensitivity to the drug.

**WARNINGS**

Some patients may experience transient episodes of light-headedness, dizziness or syncope. Orphenadrine may impair the ability of the patient to engage in potentially hazardous activities such as operating machinery or driving a motor vehicle; ambulatory patients should therefore be cautioned

accordingly.

## **PRECAUTIONS**

Confusion, anxiety and tremors have been reported in few patients receiving propoxyphene and orphenadrine concomitantly. As these symptoms may be simply due to an additive effect, reduction of dosage and/or discontinuation of one or both agents is recommended in such cases.

Orphenadrine citrate should be used with caution in patients with tachycardia, cardiac decompensation, coronary insufficiency, cardiac arrhythmias.

Safety of continuous long-term therapy with orphenadrine has not been established. Therefore, if orphenadrine is prescribed for prolonged use, periodic monitoring of blood, urine and liver function values is recommended.

### **Pregnancy**

Pregnancy Category C

Animal reproduction studies have not been conducted with orphenadrine. It is also not known whether orphenadrine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Orphenadrine should be given to a pregnant woman only if clearly needed.

### **Pediatric Use**

Safety and effectiveness in pediatric patients have not been established.

## **ADVERSE REACTIONS**

Adverse reactions of orphenadrine are mainly due to the mild anti-cholinergic action of orphenadrine, and are usually associated with higher dosage. Dryness of the mouth is usually the first adverse effect to appear. When the daily dose is increased, possible adverse effects include: tachycardia, palpitation, urinary hesitancy or retention, blurred vision, dilatation of pupils, increased ocular tension, weakness, nausea, vomiting, headache, dizziness, constipation, drowsiness, hypersensitivity reactions, pruritus, hallucinations, agitation, tremor, gastric irritation, and rarely urticaria and other dermatoses. Infrequently, an elderly patient may experience some degree of mental confusion. These adverse reactions can usually be eliminated by reduction in dosage. Very rare cases of aplastic anemia associated with the use of orphenadrine tablets have been reported. No causal relationship has been established.

## **DRUG ABUSE AND DEPENDENCE**

Orphenadrine has been chronically abused for its euphoric effects. The mood elevating effects may occur at therapeutic doses of orphenadrine.

## **OVERDOSAGE**

Orphenadrine is toxic when overdosed and typically induces anticholinergic effects. In a review of orphenadrine toxicity, the minimum lethal dose was found to be 2 to 3 grams for adults; however, the range of toxicity is variable and unpredictable. Treatment for orphenadrine overdose is evacuation of stomach contents (when necessary), charcoal at repeated doses, intensive monitoring, and appropriate supportive treatment of any emergent anticholinergic effects.

## **DOSAGE AND ADMINISTRATION**

## **Adults**

Two tablets per day; one in the morning and one in the evening.

## **HOW SUPPLIED**

Orphenadrine citrate extended-release tablets 100 mg are round, white tablets, debossed GG 931 on one side and plain on the reverse side and are supplied as:

NDC 68788-9150-3 in bottles of 30 tablets

NDC 68788-9150-6 in bottles of 60 tablets

NDC 68788-9150-9 in bottles of 90 tablets

NDC 68788-9150-1 in bottles of 100 tablets

NDC 68788-9150-8 in bottles of 120 tablets

Store at 20°-25°C (68°-77°F) [see USP Controlled Room Temperature]. Dispense in a tight, light-resistant container.

01-2010M

8100

GIN-480-00

Manufactured by

Sandoz Inc.

Princeton, NJ 08540

for GAVIS Pharmaceuticals, LLC

400 Campus Drive

Somerset, NJ 08873

Repackaged By:

Preferred Pharmaceuticals, Inc

Anaheim, CA 92807

## **100 mg Label**

NDC 68788-9150

Orphenadrine

Citrate

Extended-Release

Tablets

100 mg

Rx only

GAVIS

PHARMACEUTICALS

**Orphenadrine Citrate  
Extended-Release  
Tablets 100mg**

Generic for: Norflex  
Each tablet contains: Orphenadrine Citrate  
Extended-Release Tablets 100mg

Pkg Size: Exp Date:  
Lot#: Batch#:  
In:  
Mfg: Sandoz, Inc., Princeton, NJ  
Prod#:

Warning:  
Keep Orphenadrine Citrate Extended-Release Tablets out of children's reach. Store at 20° to 25° (68° to 77°). See USP Controlled Substances. Tablets are not for use in children.



Directions English:

Take \_\_\_\_\_ tablet(s)  
every \_\_\_\_\_ hours.  
May cause drowsiness.  
Do not drink alcoholic  
beverages while taking  
this medicine

Instrucciones Espanol:

Toma \_\_\_\_\_ tableta(s)  
cada \_\_\_\_\_ horas.  
Puede causar  
somnolencia.  
No tome bebidas  
alcoholicas mientras  
toma esta medicina.

Orphenadrine Citrate  
Extended-Release Tablets 100mg  
NDC: 68788-9150-3  
Lot: \_\_\_\_\_  
Batch: \_\_\_\_\_  
Prod# (NDC): \_\_\_\_\_

Log

Orphenadrine Citrate  
Extended-Release Tablets 100mg  
NDC: 68788-9150-6  
Lot: \_\_\_\_\_  
Batch: \_\_\_\_\_  
Prod# (NDC): \_\_\_\_\_

Client

Orphenadrine Citrate  
Extended-Release Tablets 100mg  
NDC: 68788-9150-9  
Lot: \_\_\_\_\_  
Batch: \_\_\_\_\_  
Prod# (NDC): \_\_\_\_\_

Billing

Orphenadrine Citrate  
Extended-Release Tablets 100mg  
NDC: 68788-9150-1  
Lot: \_\_\_\_\_  
Batch: \_\_\_\_\_  
Prod# (NDC): \_\_\_\_\_

Pharm

**ORPHENADRINE CITRATE**

orphenadrine citrate tablet, extended release

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68788-9150(NDC:43386-480)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ORPHENADRINE CITRATE (UNII: X0A40N8I4S) (ORPHENADRINE - UNII:AL805O9OG9)	ORPHENADRINE CITRATE	100 mg

**Inactive Ingredients**

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	GG931
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-9150-3	30 in 1 BOTTLE		
2	NDC:68788-9150-6	60 in 1 BOTTLE		
3	NDC:68788-9150-9	90 in 1 BOTTLE		
4	NDC:68788-9150-1	100 in 1 BOTTLE		
5	NDC:68788-9150-8	120 in 1 BOTTLE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040284	06/19/1998	

**Labeler** - Preferred Pharmaceuticals, Inc (791119022)

## Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals, Inc		791119022	repack

Revised: 1/2010

Preferred Pharmaceuticals, Inc