

FAMOTIDINE- famotidine tablet
Family Dollar Stores, Inc.

Dr.Reddy's Laboratories Limited

Active ingredient (in each tablet)

Famotidine USP, 10 mg/20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **For Famotidine 10 mg:**
- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor
- **For Famotidine 20 mg:**
- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

Inactive ingredients

colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, synthetic red iron oxide (only in 10 mg), talc and titanium dioxide

Questions or comments?

call **1-888-375-3784**

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

PACKAGE LABEL PRINCIPAL DISPLAY PANEL SECTION



9 2 7 9 6 0 0 5 1

Drug Facts	
Active ingredient (in each tablet)	Purpose
Famotidine USP, 10 mg	Acid reducer
Uses	
<ul style="list-style-type: none"> relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages 	
Warnings	
<p>Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease 	
Directions	
<p>adults and children 12 years and over:</p> <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn do not use more than 2 tablets in 24 hours children under 12 years: ask a doctor 	
Other information	
<p>read the directions and warnings before use</p> <p>keep the carton. It contains important information.</p> <p>store at 20°-25°C (68°-77°F)</p> <p>protect from moisture</p>	
Inactive ingredients	
colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, synthetic red iron oxide, talc, and titanium dioxide	
Questions? call 1-888-375-3784	

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods or beverages

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Original Strength Peppid AC[®]. Peppid AC[®] is a registered trademark of Johnson & Johnson.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA

Made in India

REV 07/23

FAMILY Wellness[®]

ORIGINAL STRENGTH
FAMOTIDINE
TABLETS USP, 10 mg
Acid Reducer

COMPARE TO THE ACTIVE INGREDIENT IN PEPCID AC[®]

1 BOTTLE INSIDE



ORIGINAL STRENGTH FAMOTIDINE
TABLETS USP, 10 mg
Acid Reducer

JUST ONE TABLET!
Prevents & relieves heartburn due to acid indigestion

60 TABLETS

NDC 55319-426-60

LOT

EXP

NOT 100% SATISFIED? Return within 30 days to any Family Dollar store for a refund (with receipt) or exchange.

SKU 997772



0 52251 93341 7

FAMILY Wellness[®]

ORIGINAL STRENGTH
FAMOTIDINE
TABLETS USP, 10 mg
Acid Reducer

JUST ONE TABLET!

Prevents & relieves heartburn due to acid indigestion

60 TABLETS

NDC 55319-426-60

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Famotidine USP, 10 mg

Purpose Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Directions

adults and children 12 years and over:

- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

read the directions and warnings before use

keep the carton. It contains important information.

store at 20°-25°C (68°-77°F)

protect from moisture

Inactive ingredients colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, synthetic red iron oxide, talc, and titanium dioxide

Questions? call 1-888-375-3784

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA

Made in India

REV 07/23

150094477

LOT

EXP



MAXIMUM STRENGTH FAMOTIDINE

TABLETS USP, 20 mg Acid Reducer

JUST ONE TABLET!

Prevents & relieves heartburn
due to acid indigestion

25 TABLETS

NDC 55319-427-25

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each tablet) Famotidine USP, 20 mg
Purpose Acid reducer **Uses** ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages
Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers.
Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting
 ■ stomach pain ■ kidney disease
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
 Stop use and ask a doctor if ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours
 ■ children under 12 years: ask a doctor
Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°-25°C (68°-77°F) ■ protect from moisture **Inactive ingredients** colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, talc and titanium dioxide **Questions?** call 1-888-375-3784

Made in India 150094475 REV 07/23

LOT
EXP



Drug Facts (continued)
 ■ stomach pain
 ■ kidney disease
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
 Stop use and ask a doctor if
 ■ your heartburn continues or worsens
 ■ you need to take this product for more than 14 days
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ adults and children 12 years and over:
 ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
 ■ do not use more than 2 tablets in 24 hours
 ■ children under 12 years: ask a doctor

Other information
 ■ read the directions and warnings before use
 ■ keep the carton. It contains important information.
 ■ store at 20°-25°C (68°-77°F)
 ■ protect from moisture

Drug Facts (continued)
Inactive ingredients
 colloidal silicon dioxide, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, talc and titanium dioxide

Questions? call 1-888-375-3784

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods or beverages

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Maximum Strength Pepcid AC®. Pepcid AC® is a registered trademark of Johnson & Johnson.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
 500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA
 Made in India

NOT 100% SATISFIED?
 Return within 30 days to the store of purchase for a refund (with receipt) or exchange.



Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



MAXIMUM STRENGTH FAMOTIDINE

TABLETS USP, 20 mg Acid Reducer

JUST ONE TABLET!

Prevents & relieves
heartburn due to acid
indigestion



25 TABLETS NDC 55319-427-25

Drug Facts

Active ingredient (in each tablet)	Purpose
Famotidine USP, 20 mg	Acid reducer

Uses
 ■ relieves heartburn associated with acid indigestion and sour stomach
 ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
 Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ with other acid reducers
 Ask a doctor before use if you have
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating, or dizziness
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 ■ frequent chest pain
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting



Rev. date 11/23



LOT
EXP

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-426
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	C;118
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-426-60	1 in 1 CARTON	10/01/2023	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077367	11/01/2021	

FAMOTIDINE

famotidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-427
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-427-25	1 in 1 CARTON	10/01/2023	
1		25 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55319-427-50	1 in 1 CARTON	10/01/2023	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:55319-427-10	1 in 1 CARTON	10/01/2023	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077367	11/01/2021	

Labeler - Family Dollar Stores, Inc. (024472631)

Revised: 10/2024

Family Dollar Stores, Inc.