


## **CIPROFLOXACIN- ciprofloxacin hydrochloride solution solution/ drops DIRECT RX**

### **CIPROFLOXACIN OPTH SOLUTION**

#### **DESCRIPTION SECTION**

Ciprofloxacin Ophthalmic Solution is a synthetic, sterile, multiple dose, antimicrobial for topical use. Ciprofloxacin is a fluoroquinolone antibacterial active against a broad spectrum of gram positive and gram-negative ocular pathogens. It is available as the monohydrochloride monohydrate salt of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinoline-carboxylic acid. It is a faint to light yellow crystalline powder with a molecular weight of 385.8. Its empirical formula is  $C_{17}H_{18}FN_3O_3 \cdot HCl \cdot H_2O$  and its chemical structure is as follows:



Ciprofloxacin differs from other quinolones in that it has a fluorine atom at the 6-position, a piperazine moiety at the 7-position, and a cyclopropyl ring at the 1-position.

Each mL of Ciprofloxacin Ophthalmic Solution contains: Active: Ciprofloxacin HCl 3.5 mg equivalent to 3 mg base. Preservative: benzalkonium chloride 0.006%. Inactives: sodium acetate, acetic acid, mannitol 4.6%, edetate disodium 0.05%, hydrochloric acid and/or sodium hydroxide (to adjust pH) and Water for Injection. The pH is approximately 4.5 and the osmolality is approximately 300 mOsm.

#### **CLINICAL PHARMACOLOGY SECTION**

**Systemic Absorption:** A systemic absorption study was performed in which Ciprofloxacin Ophthalmic Solution was administered in each eye every two hours while awake for two days followed by every four hours while awake for an additional 5 days. The maximum reported plasma concentration of ciprofloxacin was less than 5 ng/mL. The mean concentration was usually less than 2.5 ng/mL.

**Microbiology:** Ciprofloxacin has in vitro activity against a wide range of gram-negative and gram-positive organisms. The bactericidal action of ciprofloxacin results from interference with the enzyme DNA gyrase which is needed for the synthesis of bacterial DNA. Ciprofloxacin has been shown to be active against most strains of the following microorganisms both in vitro and in clinical infections (SEE INDICATIONS AND USAGE section):

Gram-Positive:

*Staphylococcus aureus*

*Staphylococcus epidermidis*

*Streptococcus pneumoniae*

*Streptococcus* (Viridans Group)

Gram-Negative:

*Haemophilus influenzae*

*Pseudomonas aeruginosa*

*Serratia marcescens*

Ciprofloxacin has been shown to be active in vitro against most strains of the following organisms, however, the clinical significance of these data is unknown:

Gram-Positive:

*Enterococcus faecalis* (Many strains are only moderately susceptible)

*Staphylococcus haemolyticus*

*Staphylococcus hominis*

*Staphylococcus saprophyticus*

*Streptococcus pyogenes*

Gram-Negative

*Acinetobacter calcoaceticus* *Escherichia coli* *Proteus mirabilis*

subsp. *anitratum* *Haemophilus ducreyi* *Proteus vulgaris*

*Aeromonas caviae* *Haemophilus parainfluenzae* *Providencia rettgeri*

*Aeromonas hydrophila* *Klebsiella pneumoniae* *Providencia stuartii*

*Brucella melitensis* *Klebsiella oxytoca* *Salmonella enteritidis*

*Campylobacter coli* *Legionella pneumophila* *Salmonella typhi*

*Campylobacter jejuni* *Moraxella* (*Branhamella*) *Shigella sonnei*

*Citrobacter diversus* *catarrhalis* *Shigella flexneri*

*Citrobacter freundii* *Morganella morganii* *Vibrio cholerae*

*Edwardsiella tarda* *Neisseria gonorrhoeae* *Vibrio parahaemolyticus*

*Enterobacter aerogenes* *Neisseria meningitidis* *Vibrio vulnificus*

*Enterobacter cloacae* *Pasteurella multocida* *Yersinia enterocolitica*

Other Organisms: *Chlamydia trachomatis* (only moderately susceptible) and *Mycobacterium tuberculosis* (only moderately susceptible).

Most strains of *Pseudomonas cepacia* and some strains of *Pseudomonas maltophilia* are resistant to ciprofloxacin as are most anaerobic bacteria, including *Bacteroides fragilis* and *Clostridium difficile*. The minimal bactericidal concentration (MBC) generally does not exceed the minimal inhibitory concentration (MIC) by more than a factor of 2. Resistance

to ciprofloxacin in vitro usually develops slowly (multiple-step mutation).

Ciprofloxacin does not cross-react with other antimicrobial agents such as beta-lactams or aminoglycosides; therefore, organisms resistant to these drugs may be susceptible to ciprofloxacin.

Clinical Studies: Following therapy with Ciprofloxacin Ophthalmic Solution, 76% of the patients with corneal ulcers and positive bacterial cultures were clinically cured and complete re-epithelialization occurred in about 92% of the ulcers. In 3 and 7 day multicenter clinical trials, 52% of the patients with conjunctivitis and positive conjunctival cultures were clinically cured and 70-80% had all causative pathogens eradicated by the end of treatment.

## **INDICATIONS & USAGE SECTION**

Ciprofloxacin Ophthalmic Solution is indicated for the treatment of infections caused by susceptible strains of the designated microorganisms in the conditions listed below:

Corneal Ulcers: *Pseudomonas aeruginosa*

*Serratia marcescens* \*

*Staphylococcus aureus*

*Staphylococcus epidermidis*

*Streptococcus pneumoniae*

*Streptococcus* (Viridans Group) \*

Conjunctivitis: *Haemophilus influenzae*

*Staphylococcus aureus*

*Staphylococcus epidermidis*

*Streptococcus pneumoniae*

\* Efficacy for this organism was studied in fewer than 10 infections.

## **CONTRAINDICATIONS SECTION**

A history of hypersensitivity to ciprofloxacin or any other component of the medication is a contraindication to its use. A history of hypersensitivity to other quinolones may also contraindicate the use of ciprofloxacin.

## **WARNINGS SECTION**

**NOT FOR INJECTION INTO THE EYE.**

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions, some following the first dose, have been reported in patients receiving systemic quinolone therapy. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, tingling, pharyngeal or facial edema, dyspnea, urticaria, and itching. Only a few patients had a history of hypersensitivity reactions. Serious anaphylactic reactions require

immediate emergency treatment with epinephrine and other resuscitation measures, including oxygen, intravenous fluids, intravenous antihistamines, corticosteroids, pressor amines and airway management, as clinically indicated. Remove contact lenses before using.

## **PRECAUTIONS SECTION**

General: As with other antibacterial preparations, prolonged use of ciprofloxacin may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated. Whenever clinical judgment dictates, the patient should be examined with the aid of magnification, such as slit lamp biomicroscopy and, where appropriate, fluorescein staining. Ciprofloxacin should be discontinued at the first appearance of a skin rash or any other sign of hypersensitivity reaction. In clinical studies of patients with bacterial corneal ulcer, a white crystalline precipitate located in the superficial portion of the corneal defect was observed in 35 (16.6%) of 210 patients. The onset of the precipitate was within 24 hours to 7 days after starting therapy. In one patient, the precipitate was immediately irrigated out upon its appearance. In 17 patients, resolution of the precipitate was seen in 1 to 8 days (seven within the first 24-72 hours), in five patients, resolution was noted in 10-13 days. In nine patients, exact resolution days were unavailable; however, at follow-up examinations, 18-44 days after onset of the event, complete resolution of the precipitate was noted. In three patients, outcome information was unavailable. The precipitate did not preclude continued use of ciprofloxacin, nor did it adversely affect the clinical course of the ulcer or visual outcome. (SEE ADVERSE REACTIONS).

Information for patients: Do not touch dropper tip to any surface, as this may contaminate the solution.

Drug Interactions: Specific drug interaction studies have not been conducted with ophthalmic ciprofloxacin. However, the systemic administration of some quinolones has been shown to elevate plasma concentrations of theophylline, interfere with the metabolism of caffeine, enhance the effects of the oral anticoagulant, warfarin, and its derivatives, and has been associated with transient elevations in serum creatinine in patients receiving cyclosporine concomitantly.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Eight in vitro mutagenicity tests have been conducted with ciprofloxacin and the test results are listed below:

Salmonella/Microsome Test (Negative)

E. coli DNA Repair Assay (Negative)

Mouse Lymphoma Cell Forward Mutation Assay (Positive)

Chinese Hamster V79 Cell HGPRT Test (Negative)

Syrian Hamster Embryo Cell Transformation Assay (Negative)

Saccharomyces cerevisiae Point Mutation Assay (Negative)

Saccharomyces cerevisiae Mitotic Crossover and Gene Conversion Assay (Negative)

Rat Hepatocyte DNA Repair Assay (Positive)

Thus, two of the eight tests were positive, but the results of the following three in vivo test systems gave negative results:

Rat Hepatocyte DNA Repair Assay

Micronucleus Test (Mice)

Dominant Lethal Test (Mice).

Long term carcinogenicity studies in mice and rats have been completed. After daily oral dosing for up to two years, there is no evidence that ciprofloxacin had any carcinogenic or tumorigenic effects in these species.

**Pregnancy:** Pregnancy Category C. Reproduction studies have been performed in rats and mice at doses up to six times the usual daily human oral dose and have revealed no evidence of impaired fertility or harm to the fetus due to ciprofloxacin. In rabbits, as with most antimicrobial agents, ciprofloxacin (30 and 100 mg/kg orally) produced gastrointestinal disturbances resulting in maternal weight loss and an increased incidence of abortion. No teratogenicity was observed at either dose. After intravenous administration, at doses up to 20 mg/kg, no maternal toxicity was produced and no embryotoxicity or teratogenicity was observed. There are no adequate and well controlled studies in pregnant women. Ciprofloxacin Ophthalmic Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

**Nursing Mothers:** It is not known whether topically applied ciprofloxacin is excreted in human milk. However, it is known that orally administered ciprofloxacin is excreted in the milk of lactating rats and oral ciprofloxacin has been reported in human breast milk after a single 500 mg dose. Caution should be exercised when Ciprofloxacin Ophthalmic Solution is administered to a nursing mother.

**Pediatric Use:** Safety and effectiveness in pediatric patients below the age of 1 year have not been established. Although ciprofloxacin and other quinolones cause arthropathy in immature animals after oral administration, topical ocular administration of ciprofloxacin to immature animals did not cause any arthropathy and there is no evidence that the ophthalmic dosage form has any effect on the weight bearing joints.

**Geriatric Use:** No overall differences in safety or effectiveness have been observed between elderly and younger patients.

## **ADVERSE REACTIONS SECTION**

The most frequently reported drug related adverse reaction was local burning or discomfort. In corneal ulcer studies with frequent administration of the drug, white crystalline precipitates were seen in approximately 17% of patients (SEE PRECAUTIONS). Other reactions occurring in less than 10% of patients included lid margin crusting, crystals/scales, foreign body sensation, itching, conjunctival hyperemia and a bad taste following instillation. Additional events occurring in less than 1% of patients included corneal staining, keratopathy/keratitis, allergic reactions, lid edema, tearing, photophobia, corneal infiltrates, nausea and decreased vision.

to report suspected adverse reactions call 1-800-332-1088 <sup>1</sup>

---

<sup>1</sup> to report suspected adverse reactions call 1-800-332-1088

## **OVERDOSAGE SECTION**

A topical overdose of Ciprofloxacin Ophthalmic Solution may be flushed from the eye(s) with warm tap water.

## **DOSAGE & ADMINISTRATION SECTION**

**Corneal Ulcers:** The recommended dosage regimen for the treatment of corneal ulcers is two drops into the affected eye every 15 minutes for the first six hours and then two drops into the affected eye every 30 minutes for the remainder of the first day. On the second day, instill two drops in the affected eye hourly. On the third through the fourteenth day, place two drops in the affected eye every four hours. Treatment may be continued after 14 days if corneal re-epithelialization has not occurred.

**Bacterial Conjunctivitis:** The recommended dosage regimen for the treatment of bacterial conjunctivitis is one or two drops instilled into the conjunctival sac(s) every two hours while awake for two days and one or two drops every four hours while awake for the next five days.

**How Supplied:** As a sterile ophthalmic solution, in a plastic dispenser:

2.5 mL NDC16571-120-25

5 mL NDC16571-120-50

**STORAGE:** Store at 25°C (77°F); excursions permitted to 15 - 30°C (59 - 86°F) [See USP controlled room temperature]. Protect from light. Retain in carton until contents are used.

## **ANIMAL PHARMACOLOGY & OR TOXICOLOGY SECTION**

Ciprofloxacin and related drugs have been shown to cause arthropathy in immature animals of most species tested following oral administration. However, a one-month topical ocular study using immature Beagle dogs did not demonstrate any articular lesions.

Rx Only

Manufactured in India for:

Nexus Pharmaceuticals Inc.,

Distributed by: Pack Pharmaceuticals, LLC, Buffalo Grove, IL, 60089.

Close

**CIPROFLOXACIN OPTH SOLUTION**  
NDC 61919-491-25 2.5 mL  
Lot Exp Date 04/19  
Mfg NDC 61314-656-25

**CIPROFLOXACIN OPTH SOLUTION**  
NDC 61919-491-25 2.5 mL  
Lot Exp Date 04/19  
Mfg NDC 61314-656-25

**CIPROFLOXACIN OPTH SOLUTION**  
NDC 61919-491-25 2.5 mL  
Lot Exp Date 04/19  
Mfg NDC 61314-656-25

**CIPROFLOXACIN OPTH SOLUTION**  
NDC 61919-491-25 2.5 mL  
Lot Exp Date 04/19  
Mfg NDC 61314-656-25

**For EYES**

**M**

**NDC 61919-491-25**

Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed.  
RX ONLY-KEEP OUT OF REACH OF CHILDREN  
Dosage: See package insert. Store between 68-77 degrees F

**CIPROFLOXACIN OPTH SOLUTION**  
**0.3% 2.5 mL**

Generic For: **CIOXAN**  
Each mL contains: Active: ciprofloxacin hcl 3.5mg equivalent to 3mg base

Lot# Prod# 491-25  
Discard After: 04/19  
Packaged and Distributed By: **DIRECT**  
Alpharetta, GA 30005

**D**

Mfg For: Sandoz Inc.  
Princeton, NJ 08540  
NDC 61314-656-25

Mfg Lot: 3/14/2018

NDC 61919-491-05

**Ciprofloxacin Opth Soln**

**0.3% 5 mL**

Generic For: **Ciloxan**  
Each mL contains: Active: ciprofloxacin HCL 3.5 mg equivalent to 3 mg base

Lot# 23MY2221  
Prod# 4144-003-05  
Discard After: 10/31/23  
10/31/23  
BME7M Dawsonville, GA 30534

Packaged and Distributed By: **DIRECT**

Mfg Lot: 107WC  
BW 3/23/222 1437118  
Mfg Lot: 107WC  
NDC 61919-491-05 5 mL  
Lot 23MY2221 Exp 10/31/23  
Mfg NDC 61314-656-05

Ciprofloxacin Opth Soln 0.3%  
NDC 61919-491-05 5 mL  
Lot 23MY2221 Exp 10/31/23  
Mfg NDC 61314-656-05

Ciprofloxacin Opth Soln 0.3%  
NDC 61919-491-05 5 mL  
Lot 23MY2221 Exp 10/31/23  
Mfg NDC 61314-656-05

Ciprofloxacin Opth Soln 0.3%  
NDC 61919-491-05 5 mL  
Lot 23MY2221 Exp 10/31/23  
Mfg NDC 61314-656-05

Ciprofloxacin Opth Soln 0.3%  
NDC 61919-491-05 5 mL  
Lot 23MY2221 Exp 10/31/23  
Mfg NDC 61314-656-05

Mfg. For Sandoz Inc.  
Princeton, NJ 08540  
NDC 61314-656-05

## CIPROFLOXACIN

ciprofloxacin hydrochloride solution solution/ drops

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:61919-491(NDC:61314-656)
<b>Route of Administration</b>	OPHTHALMIC		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CIPROFLOXACIN HYDROCHLORIDE</b> (UNII: 4BA73M5E37) (CIPROFLOXACIN - UNII:5E8K9I0O4U)	CIPROFLOXACIN	3 mg in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SODIUM ACETATE</b> (UNII: 4550K0SC9B)	
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0K0OR)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>HYDROCHLORIC ACID</b> (UNII: QTT17582CB)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61919-491-25	1 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2018	
2	NDC:61919-491-50	1 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2018	
3	NDC:61919-491-05	5 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/15/2018	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA019992	03/15/2018	

**Labeler** - DIRECT RX (079254320)**Establishment**

Name	Address	ID/FEI	Business Operations
DIRECT RX		079254320	relabel(61919-491)

Revised: 6/2022

DIRECT RX