

**TOPCARE TRIPLE ANTIBIOTIC PLUS PAIN RELIEF- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment**

**Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**TopCare®**

**Triple Antibiotic Plus Pain Relief**

***Drug Facts***

<b><i>Active ingredients (in each gram)</i></b>	<b><i>Purpose</i></b>
Bacitracin zinc USP 500 units	First aid antibiotic
Neomycin sulfate USP 3.5 mg	First aid antibiotic
Polymyxin B sulfate USP 10,000 units	First aid antibiotic
Pramoxine HCl USP 10 mg	External analgesic

**Uses**

first aid to help prevent infection and for the temporary relief of pain or discomfort in minor:

- cuts
- scrapes
- burns

**Warnings**

**For external use only.**

**Do not use**

- if you are allergic to any of the ingredients
- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- you need to use longer than 1 week
- condition persists or gets worse
- symptoms persist for more than 1 week, or clear up and occur again within a few days
- rash or other allergic reaction develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 2 years of age and older:
  - clean the affected area

- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

**Other information**

- To open: unscrew cap, pull tab to remove foil seal
- store at 20° to 25°C (68° to 77°F)
- see carton or tube crimp for lot number and expiration date

**Inactive ingredient**

white petrolatum

**Questions?**

Call **1-888-423-0139**

DISTRIBUTED BY  
TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007

**PRINCIPAL DISPLAY PANEL - 28.4 g Tube Carton**

TopCare®  
health™

NDC 36800-056-24

MAXIMUM STRENGTH

Triple Antibiotic Ointment  
+ Pain Relief

BACITRACIN ZINC, NEOMYCIN SULFATE, POLYMYXIN B SULFATE & PRAMOXINE HCL  
First Aid Antibiotic / Pain Relieving Ointment

• OUR PHARMACISTS  
RECOMMEND •

NET WT 1 OZ (28.4 g)

COMPARE TO NEOSPORIN®  
+ PAIN RELIEF  
ACTIVE INGREDIENTS\*

# First Aid Antibiotic & Pain Relieving Ointment

Helps Prevent Infection in Minor Cuts,  
Scrapes & Burns  
Maximum Strength

LK-3727-4  
01-19-4  
17

**TopCare.**  
health™

**MAXIMUM STRENGTH**



## Triple Antibiotic Ointment

+ Pain Relief

**BACITRACIN ZINC, NEOMYCIN SULFATE, POLYMYXIN B SULFATE & PRAMOXINE HCL**

First Aid Antibiotic / Pain Relieving Ointment

NET WT 1 OZ (28.4 g)

**Triple Antibiotic Ointment + Pain Relief**  
BACITRACIN ZINC, NEOMYCIN SULFATE, POLYMYXIN B SULFATE & PRAMOXINE HCL

NDC 36800-0568-26



DISTRIBUTED BY  
TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007  
©TOPCO TARA0718  
QUESTIONS? 1-888-423-0139  
topcare@topco.com  
www.topcarebrand.com  
MADE IN CANADA



Scan here for  
more information



This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.

TS1

NO COPY ON THIS FLAP  
FOR LOT # AND EXPIRY  
DATE PRINT

\*All trademarks are property of their respective owners. The product is not affiliated with the makers/owners of Neosporin® + Pain Relief.

<p><b>Drug Facts (continued)</b></p> <p>• symptoms persist for more than 1 week, or clear up and occur again within a few days</p> <p>• rash or other allergic reaction develops</p> <p>• keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <p>• adults and children 2 years of age and older</p> <p>• clean the affected area</p> <p>• apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily</p> <p>• may be covered with a sterile bandage</p> <p>• children under 2 years of age: ask a doctor</p> <p><b>Other information</b></p> <p>• To open ureter cap, pull tab to remove foil seal</p> <p>• store at 20° to 25°C (68° to 77°F)</p> <p>• see carton or tube crimp for lot number and expiration date</p> <p><b>Inactive ingredient</b></p> <p>with petroleum</p> <p><b>Questions?</b> Call 1-888-423-0139</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each gram)</b></p> <p>Bacitracin zinc USP 500 units.....First aid antibiotic</p> <p>Neomycin sulfate USP 3.5 mg.....First aid antibiotic</p> <p>Polymyxin B sulfate USP 10,000 units.....First aid antibiotic</p> <p>Pramoxine HCl USP 10 mg.....External analgesic</p> <p><b>Purpose</b></p> <p>First aid antibiotic</p> <p><b>Uses</b></p> <p>First aid to help prevent infection and for the temporary relief of pain or discomfort in minor:</p> <p>• cuts • scrapes • burns</p> <p><b>Warnings</b></p> <p>For external use only.</p> <p>Do not use</p> <p>• if you are allergic to any of the ingredients</p> <p>• in the eyes</p> <p>• over large areas of the body</p> <p>Ask a doctor before use if you have</p> <p>• deep or puncture wounds • animal bites • serious burns</p> <p>Stop use and ask a doctor if</p> <p>• you need to use longer than 1 week</p> <p>• condition persists or gets worse</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

**Triple Antibiotic Ointment + Pain Relief**

**TopCare.**  
health™

## TOPCARE TRIPLE ANTIBIOTIC PLUS PAIN RELIEF

bacitracin zinc, neomycin sulfate, polymyxin b sulfate, and pramoxine hydrochloride ointment

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-056
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Bacitracin zinc</b> (UNII: 89Y4M234ES) (Bacitracin - UNII:58H6RWO52I)	Bacitracin	500 [iU] in 1 g
<b>Neomycin sulfate</b> (UNII: 057Y626693) (Neomycin - UNII:I16QD7X297)	Neomycin	3.5 mg in 1 g
<b>Polymyxin B sulfate</b> (UNII: 19371312D4) (Polymyxin B - UNII:J2VZ07J96K)	Polymyxin B	10000 [iU] in 1 g
<b>Pramoxine hydrochloride</b> (UNII: 88AYB867L5) (Pramoxine - UNII:068X84E056)	Pramoxine hydrochloride	10 mg in 1 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>petrolatum</b> (UNII: 4T6H12BN9U)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:36800-056-24	1 in 1 CARTON	03/31/2012	
1		28.4 g in 1 TUBE; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL	part333B	03/31/2012	

**Labeler** - Topco Associates LLC (006935977)**Registrant** - Taro Pharmaceuticals U.S.A., Inc (145186370)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Taro Pharmaceuticals Inc.		206263295	MANUFACTURE(36800-056)