DG HEALTH CHILDRENS IBUPROFEN- ibuprofen suspension Dolgencorp, LLC

Dolgencorp, LLC Children's Ibuprofen Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- · takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- · child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

give with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

· any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

^{**}or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze

Inactive ingredients

anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments?

1-888-309-9030

Principal Display Panel

SEE NEW WARNINGS

Compare to the active ingredient of Children's Motrin® Bubble Gum Flavor

For Ages 2 to 11 Years

Children's Ibuprofen

Oral Suspension

100 mg per 5 mL

Pain Reliever/Fever Reducer (NSAID)

LASTS UP TO 8 HOURS

Alcohol free

Gluten free

Bubble Gum Flavor

4 FL OZ (118 mL)



Bubble Gum Flavor

Pain Reliever∖Fever Reducer ∩ SAID) TO 0 wild be 12 mg Oral Suspension

i bu profen

For Ages 2 to 11 Years

De pestru

DG™|health

For Ages 2 to 11 Years

lbuprofen

Oral Suspension

100 mg per 5 mL

Pain Reliever/

Fever Reducer

4 FL 0Z (118 mL)

(NSAID)

Childrens

WARNINGS

Compare to the active ingredient of Children's Motrin[®] Bubble Gum

Bubble

Flavor

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL) Ibuprofen 100 mg (NSAID)*... .Pain reliever fever reducer

nonsteroidal anti-inflammatory drug

- Uses temporarily: relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache

Purposes

■ Disters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

That had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or

- takes a blood thinning (anticoagulant) or steroid drug
 takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 takes more or for a longer time than directed Heart ettack and stroke warning; NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use

Do not use

■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer

Drug Facts (continued)

■ right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child child has a history of stomach problems, such as heartburn
- such as hearthum

 child has problems or serious side effects
 from taking pain relievers or fever reducers

 child has not been drinking fluids

 child has lost a lot of fluid due to vorniting
 or diarrhea

 child has high blood pressure, heart disease,
 liver cirrhosis, kidney disease, or had a stroke

 child has asthma

 child is taking a diuretic

Ask a doctor or pharmacist before use if

the child is

under a doctor's care for any serious condition ■ taking any other drug

When using this product
give with food or milk if stomach upset occurs

- Stop use and ask a doctor if
 child experiences any of the following signs of stomach bleeding:
 feels faint
 comits blood
 has bleether bleed stools

- has bloody or black stools
 has stomach pain that does not get better
 child has symptoms of heart problems or
- chest pain

- trouble breathing
 weakness in one part or side of body
 surred speech
 leg swelling
 the child does not get any relief within first
 day (24 hours) of treatment
 fever or pain gets worse or lasts more than
- 3 days ■ redness or swelling is present in the
- nainful area

■ any new symptoms appea

Keep out of reach of children. In case of overdose, get medical help or contact a Poisor Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- or complete warnings for adult use

 do not give more than directed

Drug Facts (continued)

- shake well before using
- mL = milliliter
- mIL = milliliter
 ind right does on chart. If possible, use
 weight to does; otherwise use age.
 use only enclosed dosing cup. Do not
 use any other dosing device.
 if needed, repeat dose every 6-8 hours
 do not use more than 4 times a day
 replace original bottle cap to maintain
 debt designed.

- child resistance

 wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

as directed by a docto

Other information

- each 5 mL contains: sodium 2 mg
 do not use if printed neckband is
 broken or missing
 store at 20-25°C (68-77°F)
- do not freeze

Inactive ingredients anhydrous citric acid, FD&C red #40, glycerin, high fructose com syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments? 1-888-309-9030

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distribute of Children's Motrin® Bubble Gum Flavor. DISTRIBUTED BY OLD EAST MAIN CO. - 100%

GOODLETTSVILLE, Satisfaction Guaranteed!

(888) 309-9030

A0900

16626 VT C5

DG HEALTH CHILDRENS IBUPROFEN

ibuprofen suspension

Product Information

HUMAN OTC DRUG **Product Type** Item Code (Source) NDC:55910-166

Route of Administration

ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SORBITOL (UNII: 506T60A25R)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color	PINK (light)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-166- 26	1 in 1 CARTON	05/02/2016	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	05/02/2016	

Labeler - Dolgencorp, LLC (068331990)

Revised: 10/2022 Dolgencorp, LLC