

FAMILY WELLNESS ORAL PAIN RELIEF- benzocaine 20% gel
FAMILY DOLLAR SERVICES INC

Benzocaine 20% pain relieving gel

Active ingredients

Benzocaine 20%

Purpose

Oral pain reliever

Uses

temporarily relieves pain associated with the following mouth and gum irritations:

- toothache
- canker sores
- minor dental procedures
- dentures
- sore gums
- braces

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use:

- for teething
- in children under 2 years of age

When using this product:

- avoid contact with the eyes
- do not exceed recommended dosage
- do not use for more than 7 days unless directed by a doctor/dentist

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness lasts or worsens
- swelling, rash, or fever develops

Keep out of reach of children

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions

adults and children 2 years of age and over	apply a pea-size amount to the affected area up to 4 times daily or as directed by a doctor/dentist
children under 12 years of age	adult supervision should be given in the use of this product
children under 2 years of age	do not use

- For denture irritation:
- apply thin layer to the affected area
- do not reinsert dental work until irritation/pain is relieved
- rinse mouth well before reinserting

Other information

- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

benzyl alcohol, carbomer, D&C red no. 33, FD&C blue no. 1, FD&C yellow no. 6, glycerin, natural flavor, polyethylene glycol, polysorbate 80, propylene glycol, sodium saccharin, sorbic acid



COMPARE TO THE ACTIVE INGREDIENT OF ANBESOL® ORAL PAIN RELIEVER*

MAXIMUM STRENGTH ORAL PAIN RELIEF

GEL

FAST ACTING

Benzocaine 20% pain relieving gel



NDC 55391-110-14

MAXIMUM STRENGTH FAST ACTING ORAL PAIN RELIEF

- Toothache
- Gum Pain
- Canker Sores
- Denture Pain

MAXIMUM STRENGTH ORAL PAIN RELIEF

GEL

FAST ACTING

NET WT 0.5 OZ (14 g)



SKU 999543



DISTRIBUTED BY:
MIDWOOD BRANDS LLC
500 VOLVO PARKWAY
CHESAPEAKE, VA 23320 USA



NOT 100% SATISFIED?
Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

MADE IN INDIA

36-63-001-93625-1
V93625C

*This product is not manufactured or distributed by Foundation Consumer Brands, LLC, distributor of Anbesol Oral Pain Reliever.

Drug Facts (continued)

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Directions
 adults and children 2 years of age and older: apply a pea-size amount to the affected area up to 4 times daily or as directed by a doctor/dentist.
 children under 12 years of age: adult supervision should be given in the use of this product.
 children under 2 years of age: do not use.

For denture irritation:
 ■ apply thin layer to the affected area
 ■ do not reinsert dental work until irritation/pain is relieved
 ■ rinse mouth well before reinserting

Other information ■ store at 20-25° C (68-77°F) ■ do not refrigerate

Inactive ingredients benzyl alcohol, carbomer, glycerin, natural flavor, polyethylene glycol, polysorbate 80, propylene glycol, sodium saccharin, sorbic acid

Drug Facts
 Benzocaine 20%.....Oral pain reliever

Active ingredients

Purpose

Uses
 temporarily relieves pain associated with the following mouth and gum irritations: ■ toothache ■ canker sores ■ minor dental procedures: ■ dentures ■ sore gums ■ braces

Warnings
 methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:
 ■ pale, gray, or blue colored skin (cyanosis) ■ headache
 ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness
 ■ fatigue or lack of energy

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics.

Do not use: ■ in children under 2 years of age

When using this product:
 ■ avoid contact with the eyes
 ■ do not exceed recommended dosage
 ■ do not use for more than 7 days unless directed by a doctor/dentist

Stop use and ask doctor if
 ■ sore mouth symptoms do not improve in 7 days
 ■ irritation, pain, or redness lasts or worsens
 ■ swelling, rash, or fever develops

Do not use if the printed safety seal on the carton is broken or missing. READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

FAMILY WELLNESS ORAL PAIN RELIEF
benzocaine 20% gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-108
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	0.2 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE (UNII: 0A5MM307FC)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SORBIC ACID (UNII: X045WJ989B)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-108-14	1 in 1 CARTON	02/01/2024	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	02/01/2024	

Labeler - FAMILY DOLLAR SERVICES INC (024472631)