

CAREONE CHILDRENS PAIN AND FEVER- acetaminophen suspension
American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

American Sales Company Children's Pain & Fever Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1 1/2 tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2 1/2 tsp)
72-95	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the Active Ingredient in Children's Tylenol® Oral Suspension

CHILDREN'S PAIN & FEVER

Pain Reliever/Fever Reducer

Acetaminophen 160 mg per 5 mL

Oral Suspension

BUBBLE GUM FLAVOR

FOR AGES 2-11 YEARS

Fast, Effective Relief of Children's Pain and Fever

Alcohol Free

Aspirin Free

Ibuprofen Free

See New Warnings

4 FL OZ (118 mL)

FOR AGES 2 to 11 YEARS
BUBBLE GUM FLAVOR

0 mL Suspension
Acetaminophen 160 mg per 5 mL
Pain Reliever/Fever Reducer

PAIN & FEVER

CHILDREN'S

CARBONE

Drug Facts

Active Ingredient **Purpose**
(In each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg.....Pain reliever/fever reducer

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms. ▶

Drug Facts (continued)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1 ½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2 ½ tsp)
72-95	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other Information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing
- see bottom panel for lot number and expiration date

Inactive Ingredients anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

GLUTEN FREE

DISTRIBUTED BY FOODHOLD U.S.A., LLC
LANDOVER, MD 20795 • 1-877-846-9949
©2014 S&S Brands, LLC

Quality guaranteed or your money back.



LOT NO.

EXP. *Good to Avra*

100000 04 07

CAREONE CHILDRENS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-376
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QP11U3FV8)	
CALCIUM SULFATE (UNII: WAT0DDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics

Color	RED (opaque)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-376-26	1 in 1 CARTON		
1		118 mL in 1 BOTTLE		
2	NDC:41520-376-28	1 in 1 CARTON		
2		147 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/13/2004	

Labeler - American Sales Company (809183973)

Registrant - L. Perrigo Company (006013346)

Revised: 3/2015

American Sales Company