

LAXATIVE- bisacodyl tablet, sugar coated
Rite Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid 44-327

Active ingredient
(in each tablet)

Bisacodyl 5 mg

Purpose

Stimulant laxative

Uses

- relieves occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not use within 1 hour after taking an antacid or milk
- do not chew or crush tablet(s)
- you may have stomach discomfort, faintness or cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac, silica, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

RITE

AID®

PHARMACY

***Compare to the active ingredient in Dulcolax® Laxative Tablets**

laxative

bisacodyl USP 5 mg

stimulant laxative

gentle, predictable

overnight relief

delayed-release tablets

enteric coated

25 TABLETS

5 mg EACH

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark Dulcolax® Laxative Tablets

50844 REV0312E32756

DISTRIBUTED BY: RITE AID

30 HUNTER LANE

CAMP HILL, PA 17011

IF YOU'RE NOT SATISFIED, WE'LL HAPPILY REFUND YOUR MONEY.

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION



DISTRIBUTED BY: RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011

IF YOU'RE NOT
SATISFIED, WE'LL
HAPPILY REFUND
YOUR MONEY.

TAMPER EVIDENT: DO NOT USE IF
CARTON IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING

Drug Facts

Active ingredient (in each tablet) Purpose
Bisacodyl 5 mg Stimulant laxative

Uses

- relieves occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not use within 1 hour after taking an antacid or milk
- do not chew or crush tablet(s)
- you may have stomach discomfort, faintness or cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water
- adults and children 12 years and over: take 1 to 3 tablets in a single daily dose
- children 6 to under 12 years of age: take 1 tablet in a single daily dose
- children under 6 years of age: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients

acacia, ammonium hydroxide, calcium carbonate, camauba wax, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac, silica, simethicone,

ITEM 0372923



Drug Facts (continued)

sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate


Questions or comments? 1-800-426-9391

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark Dulcolax® Laxative Tablets.

50844 REV0312A32756

B-1702-327-56-R
REV0312A32756

5 mg EACH
25 TABLETS



stimulant laxative

bisacodyl USP 5 mg

laxative

gentle, predictable
overnight relief
delayed-release tablets
enteric coated



Rite Aid 44-327

LAXATIVE

bisacodyl tablet, sugar coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0327
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10 X0 709 Y6 I) (BISACODYL - UNII:10 X0 709 Y6 I)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
AMMONIA (UNII: 5138Q19F1X)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	

POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0327-6	1 in 1 CARTON	03/25/2002	
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0327-5	2 in 1 CARTON	03/25/2002	
2		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	03/25/2002	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(11822-0327)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(11822-0327)