

DAYTIME NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr, phenylephrine hcl and acetaminophen, dextromethorphan hbrdoxylamine succinate
Kroger

Kroger Daytime and Nighttime Cold & Flu Liquid (Apricot + Anise)

Nighttime Cold & Flu

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Pain reliever/ fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed

- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 24 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

alcohol, citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

Questions or comments ?

Call **1-800-632-6900**

Daytime Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adults take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child take more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

**When using this product,
do not use more than directed.**

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 56 mg
- store at no greater than 25°C and do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

Questions or comments ?

Call **1-800-632-6900**

PRINCIPAL DISPLAY PANEL - Convenience Pack

NDC 41224-790-02

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS' DAYQUIL AND NYQUIL: COUGH, COLD & FLU RELIEF LIQUID CO-PACK

COLD & FLU

DayTime

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

ALCOHOL FREE

HEADACHE, FEVER, SORE THROAT, MINOR ACHES & PAINS
COUGH
NASAL CONGESTION, SINUS PRESSURE

12 FL OZ (354 mL)

COLD & FLU

NightTime

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

ALCOHOL 10% | NIGHTTIME RELIEF

HEADACHE, FEVER, SORE THROAT, MINOR ACHES & PAINS
COUGH
SNEEZING, RUNNY NOSE

12 FL OZ (354 mL)

VALUE PACK

POWERFUL RELIEF

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg Pain reliever/fever reducer
Dextromethorphan HBr 20 mg Cough suppressant
Doxylamine succinate 12.5 mg Antihistamine

Uses

temporarily relieves common cold/flu symptoms: cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product.

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Decongestant

Uses

temporarily relieves common cold/flu symptoms: nasal congestion

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product.

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Decongestant

Uses

temporarily relieves common cold/flu symptoms: nasal congestion

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product.

NDC 41226-782-01

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS' NYQUIL: COUGH, COLD & FLU NIGHTTIME RELIEF LIQUID

COLD & FLU

NightTime

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

NIGHTTIME RELIEF
ALCOHOL 10%

HEADACHE, FEVER, SORE THROAT, MINOR ACHES & PAINS
COUGH
SNEEZING, RUNNY NOSE

12 FL OZ (354 mL)

COLD & FLU

DayTime

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

ALCOHOL FREE

HEADACHE, FEVER, SORE THROAT, MINOR ACHES & PAINS
COUGH
NASAL CONGESTION, SINUS PRESSURE

12 FL OZ (354 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING

← PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg Pain reliever/fever reducer
Dextromethorphan HBr 20 mg Cough suppressant
Doxylamine succinate 12.5 mg Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product.

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA

For More Product Information, Scan UPC Using Your Kroger App or Call 800-632-6900

OUR PHARMACIST RECOMMENDED

782L 0324

PARENTS: Learn about teen medicine risks. www.StopMedicineAbuse.org

LOT: Print/Varnish Omit Area

EXP.:

Drug Facts (continued)

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - hives
 - blistering
- if a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- Do not use**
 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- Ask a doctor before use if you have**
 - liver disease
 - glaucoma
 - cough that occurs with too much phlegm (mucus)
 - breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
 - trouble urinating due to enlarged prostate gland
- Ask a doctor or pharmacist before use if you are**
 - taking sedatives or tranquilizers
 - taking the blood thinning drug warfarin
- When using this product**
 - excitability may occur, especially in children
 - marked drowsiness may occur
 - avoid alcoholic drinks
 - be careful when driving a motor vehicle or operating machinery

Drug Facts (continued)

- alcohol, sedatives, and tranquilizers may increase drowsiness
- Stop use and ask a doctor if**
 - rain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- If pregnant or breast-feeding,** ask a health professional before use.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
- Directions**
 - take only as directed
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
 - adults & children 12 yrs & over 30 mL every 6 hrs
 - children 6 to under 12 yrs 15 mL every 4 hrs
 - children 4 to under 6 yrs ask a doctor
 - children under 4 yrs do not use
- Other information**
 - each 30 mL contains: sodium 24 mg
 - store at no greater than 25°C and do not refrigerate
- Inactive ingredients** alcohol, citric acid monohydrate, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, base, glycerin polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucrose, xanthan gum
- Questions or comments?** Call 1-800-632-6900

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com

NDC 41226-772-01

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS' DAYQUIL: COUGH, COLD & FLU DAYTIME RELIEF LIQUID

COLD & FLU

DayTime

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

ALCOHOL FREE

HEADACHE, FEVER, SORE THROAT, MINOR ACHES & PAINS
COUGH
NASAL CONGESTION, SINUS PRESSURE

12 FL OZ (354 mL)

COLD & FLU

NightTime

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

ALCOHOL 10%

HEADACHE, FEVER, SORE THROAT, MINOR ACHES & PAINS
COUGH
SNEEZING, RUNNY NOSE

12 FL OZ (354 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING

← PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Phenylephrine HCl 5 mg Decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product.

DISTRIBUTED BY THE KROGER CO., CINCINNATI, OHIO 45202

MADE IN INDIA

For More Product Information, Scan UPC Using Your Kroger App or Call 800-632-6900

OUR PHARMACIST RECOMMENDED

772L 0324

PARENTS: Learn about teen medicine risks. www.StopMedicineAbuse.org

LOT: Print/Varnish Omit Area

EXP.:

Drug Facts (continued)

- the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adults take 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - hives
 - blistering
- if a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 7 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- Do not use**
 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- Ask a doctor before use if you have**
 - liver disease
 - heart disease
 - high blood pressure
 - thyroid disease
 - diabetes
 - trouble urinating due to enlarged prostate gland
 - cough that occurs with too much phlegm (mucus)
 - persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 - a sodium-restricted diet
- Ask a doctor or pharmacist before use if you are**
 - taking the blood thinning drug warfarin
- When using this product**
 - excitability may occur, especially in children
 - marked drowsiness may occur
 - avoid alcoholic drinks
 - be careful when driving a motor vehicle or operating machinery

Drug Facts (continued)

- When using this product, do not use more than directed.
- Stop use and ask a doctor if**
 - you get nervous, dizzy or sleepless
 - nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.
- If pregnant or breast-feeding,** ask a health professional before use.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
- Directions**
 - take only as directed
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
 - adults & children 12 yrs & over 30 mL every 4 hrs
 - children 6 to under 12 yrs 15 mL every 4 hrs
 - children 4 to under 6 yrs ask a doctor
 - children under 4 yrs do not use
- Other information**
 - each 15 mL contains: sodium 56 mg
 - store at no greater than 25°C and do not refrigerate
- Inactive ingredients** alcohol, citric acid monohydrate, D&C yellow no. 6, FD&C green no. 3, FD&C yellow no. 6, base, glycerin polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucrose, xanthan gum
- Questions or comments?** Call 1-800-632-6900

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

Our Brands. Our Guarantee. Love It or Your Money Back. www.kroger.com

DAYTIME NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl and acetaminophen, dextromethorphan hbrdoxylamine succinate kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41226-790
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-790-02	1 in 1 PACKAGE; Type 0: Not a Combination Product	04/30/2024	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	354 mL
Part 2	1 BOTTLE	354 mL

Part 1 of 2

DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl liquid

Product Information

Item Code (Source)	NDC:41226-772
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	orange (clear orange color)	Score	
Shape		Size	
Flavor	APRICOT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-772-01	354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2024	

Part 2 of 2

NIGHTTIME COLD AND FLU

acetaminophen, dextromethrophan hbr, doxylamine succinate liquid

Product Information

Item Code (Source)	NDC:41226-782
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	green (clear green color)	Score	
Shape		Size	
Flavor	ANISE (Anise, Menthol)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41226-782-01	354 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2024	

Labeler - Kroger (006999528)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(41226-790)

Revised: 5/2024

Kroger