MAGNESIUM CITRATE- leader magnesium citrate oral solution - lemon liquid CARDINAL HEALTH, INC. 063997360

Leader Magnesium Citrate Oral Solution - Lemon

Active Ingredient

Magnesium citrate 1.745g per fl oz

Purpose

Saline Laxtive

Use

for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

kidney disease
a magnesium restricted diet
abdorominal pain, nausea, or vomiting
noticed a sudden change in bowel habits that persists over a period of 2 weeks
already used a laxative for a period longer than 1 week

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any other drug.

Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

STOP USE AND ASK A DOCTOR IF

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

IF PREGNANT OR BREAST-FEEDING

ask health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center right away.

Other Information

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

shake well before using drink a full glass (8 ounces) of liquid with each dose may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age - 2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours children under 2 years of age - ask a doctor

Other Information

each fl oz contains: magnesium 290 mg

each fl oz contains: sodium 1 mg

Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions?

1-800-200-6313



Drug Facts

Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g

Uses: ■ For relief of occasional constipation

(irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings: Ask a doctor before use if you have

- Kidney disease a magnesium restricted diet Abdominal pain, nausea, or vomiting Noticed a sudden change in bowel habits that persists
- over a period of 2 weeks ■ Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are
■ Taking any other drug.
■ Take this product 2 or more hours before or after other

drugs. Laxa'tives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use if tamper evident twist-off cap is missing, broken, or separated from neckring

Drug Facts (continued)

Store at temperatures between 46° and 86°F (8° and 3°C)

Directions: ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose may be taken as a single daily dose or in divided doses 6.5 to 10 fl oz maximum 10 fl oz in 24 hours adults and children 12 years of age and over 3 to 7 fl oz maximum 7 fl oz in 24 hours children 6 to under 12 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours children 2 to under 6 years of age children under 2 years ask a doctor of age

Other Information:

- each fl oz contains: magnesium 290 mg ■ each fl oz contains: sodium 1 mg
- Inactive ingredients: benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions? 1-833-551-0932

DIST. BY CAH DUBLIN, OH 43017 ©2023 Cardinal Health. www.myleader.com 1-800-200-6313



MAGNESIUM CITRATE

leader magnesium citrate oral solution - lemon liquid

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:70000-0661

Route of Administration ORAL

Active Ingredient/Active Moiety Basis of **Ingredient Name** Strength Strength MAGNESIUM CITRATE (UNII: RHO2601T9V) (MAGNESIUM CATION -**MAGNESIUM** 1.745 g UNII:T6V3LHY838) **CITRATE** in 29.6 mL

Inactive Ingredients			
Ingredient Name	Strength		
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)			
BENZOIC ACID (UNII: 85KN0B0MIM)			
WATER (UNII: 059QF0KO0R)			
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

Packaging			
# Hom Code	Dackage Description	Marketing Start	Marketing End

#	item code	Раскаде резсприон	Date	Date	
1	NDC:70000- 0661-1	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product 02/06/2024			
Marketing Information					
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
01	TC Monograph Di	rug M007	02/06/2024		

Labeler - CARDINAL HEALTH, INC. 063997360 (063997360)

Registrant - Pharma Nobis, LLC (118856114)

Establishment				
Name	Address	ID/FEI	Business Operations	
Pharma Nobis, LLC		118564114	label(70000-0661) , manufacture(70000-0661) , analysis(70000-0661) , pack(70000-0661)	

Revised: 2/2024 CARDINAL HEALTH, INC. 063997360